

3,4,5-Trimethyl-2,4,6-triphenyl-4H-thiopyran

Hossein Rahmani,^a Hooshang Pirelahi^b and Seik Weng Ng^{c*}

^aInstitute of Chemical Industries, Iranian Research Organization for Science and Technology, PO Box 15815-358, Tehran, Iran, ^bDepartment of Chemistry, College of Science, University of Tehran, PO Box 13145-143, Tehran, Iran, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 18 February 2009; accepted 19 February 2009

Key indicators: single-crystal X-ray study; $T = 115\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.046; wR factor = 0.123; data-to-parameter ratio = 18.0.

The six-membered thiopyran ring in the title compound, $\text{C}_{26}\text{H}_{24}\text{S}$, adopts a boat conformation, with the S atom displaced by 0.478 (2) Å and the 3-methylene C atom by 0.644 (2) Å from the plane of the other four sp^2 -hybridized C atoms. The methyl group on the methylene carbon lies in a pseudo-equatorial position and the phenyl ring in a pseudo-axial position.

Related literature

For a similar compound, see: Rahmani *et al.* (2009). For the synthesis, see: Rahmani & Pirelahi (1997).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{24}\text{S}$
 $M_r = 368.51$
Monoclinic, $P2_1/n$
 $a = 8.4525 (1)$ Å
 $b = 14.4732 (2)$ Å
 $c = 16.2971 (3)$ Å
 $\beta = 103.156 (1)$ °

$V = 1941.37 (5)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.17\text{ mm}^{-1}$
 $T = 115\text{ K}$
 $0.35 \times 0.35 \times 0.20\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.840$, $T_{\max} = 0.966$

17659 measured reflections
4445 independent reflections
3576 reflections with $I > \sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.123$
 $S = 1.05$
4445 reflections

247 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.39\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Iranian Research Organization for Science and Technology and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2579).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Rahmani, H. & Pirelahi, H. (1997). *Sulfur Lett.* **20**, 159–163.
- Rahmani, H., Pirelahi, H. & Ng, S. W. (2009). *Acta Cryst. E* **65**, o603..
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supporting information

Acta Cryst. (2009). E65, o604 [doi:10.1107/S1600536809005923]

3,4,5-Trimethyl-2,4,6-triphenyl-4H-thiopyran

Hossein Rahmani, Hooshang Pirelahi and Seik Weng Ng

S1. Experimental

4-Methyl-2,4,6-triphenyl-4H-thiopyran was synthesized by the reaction of methyl magnesium bromide and 3,5-dimethyl-2,4,6-triphenyl thiopyrylium perchlorate in dry ether under an argon atmosphere according to a reported method (Rahmani & Pirelahi, 1997). The product was isolated by TLC on neutral alumina (petroleum ether 40–60 °C) and purified by recrystallization from ethanol.

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 to 0.98 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2 to 1.5 $U(\text{C})$.




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{26}\text{H}_{24}\text{S}$; probability levels are set at 70% and H-atoms are drawn as spheres of arbitrary radius.

3,4,5-Trimethyl-2,4,6-triphenyl-4H-thiopyran*Crystal data*

C₂₆H₂₄S
 $M_r = 368.51$
 Monoclinic, P2₁/n
 Hall symbol: -P 2yn
 $a = 8.4525 (1)$ Å
 $b = 14.4732 (2)$ Å
 $c = 16.2971 (3)$ Å
 $\beta = 103.156 (1)^\circ$
 $V = 1941.37 (5)$ Å³
 $Z = 4$

$F(000) = 784$
 $D_x = 1.261$ Mg m⁻³
 Mo K α radiation, $\lambda = 0.71073$ Å
 Cell parameters from 5510 reflections
 $\theta = 2.5\text{--}28.2^\circ$
 $\mu = 0.17$ mm⁻¹
 $T = 115$ K
 Block, colorless
 $0.35 \times 0.35 \times 0.20$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.840$, $T_{\max} = 0.966$

17659 measured reflections
 4445 independent reflections
 3576 reflections with $I > \sigma(I)$
 $R_{\text{int}} = 0.037$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -10 \rightarrow 10$
 $k = -18 \rightarrow 18$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.123$
 $S = 1.05$
 4445 reflections
 247 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.059P)^2 + 1.0605P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.39$ e Å⁻³
 $\Delta\rho_{\min} = -0.36$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.66266 (5)	0.60193 (3)	0.30451 (3)	0.01781 (12)
C1	0.56658 (19)	0.69496 (11)	0.34341 (10)	0.0168 (3)
C2	0.4182 (2)	0.68372 (11)	0.35910 (10)	0.0170 (3)
C3	0.32243 (19)	0.59530 (11)	0.32806 (11)	0.0165 (3)
C4	0.4297 (2)	0.51230 (12)	0.36251 (10)	0.0174 (3)
C5	0.5796 (2)	0.50682 (11)	0.34781 (10)	0.0166 (3)
C6	0.3438 (2)	0.75486 (13)	0.40643 (12)	0.0231 (4)
H6A	0.4224	0.8045	0.4257	0.035*
H6B	0.3145	0.7257	0.4552	0.035*
H6C	0.2462	0.7806	0.3692	0.035*
C7	0.1580 (2)	0.59255 (13)	0.35402 (12)	0.0223 (4)
H7A	0.1768	0.5943	0.4156	0.033*
H7B	0.1002	0.5356	0.3329	0.033*

C10—H10	0.9500	C25—C26	1.386 (2)
C11—C12	1.389 (3)	C25—H25	0.9500
C11—H11	0.9500	C26—H26	0.9500
C1—S1—C5	101.20 (8)	C13—C12—C11	119.49 (17)
C2—C1—C21	126.50 (15)	C13—C12—H12	120.3
C2—C1—S1	120.10 (13)	C11—C12—H12	120.3
C21—C1—S1	113.39 (12)	C12—C13—C14	120.77 (17)
C1—C2—C6	122.02 (15)	C12—C13—H13	119.6
C1—C2—C3	118.81 (14)	C14—C13—H13	119.6
C6—C2—C3	119.17 (14)	C13—C14—C9	120.34 (17)
C4—C3—C2	107.98 (13)	C13—C14—H14	119.8
C4—C3—C7	111.96 (14)	C9—C14—H14	119.8
C2—C3—C7	111.78 (13)	C16—C15—C20	117.66 (16)
C4—C3—C15	110.63 (13)	C16—C15—C3	120.41 (15)
C2—C3—C15	109.40 (13)	C20—C15—C3	121.70 (15)
C7—C3—C15	105.07 (13)	C17—C16—C15	121.32 (16)
C5—C4—C8	121.52 (15)	C17—C16—H16	119.3
C5—C4—C3	118.93 (14)	C15—C16—H16	119.3
C8—C4—C3	119.54 (14)	C18—C17—C16	120.20 (17)
C4—C5—C9	125.67 (15)	C18—C17—H17	119.9
C4—C5—S1	120.16 (13)	C16—C17—H17	119.9
C9—C5—S1	114.17 (12)	C19—C18—C17	119.29 (17)
C2—C6—H6A	109.5	C19—C18—H18	120.4
C2—C6—H6B	109.5	C17—C18—H18	120.4
H6A—C6—H6B	109.5	C18—C19—C20	120.40 (16)
C2—C6—H6C	109.5	C18—C19—H19	119.8
H6A—C6—H6C	109.5	C20—C19—H19	119.8
H6B—C6—H6C	109.5	C19—C20—C15	121.13 (16)
C3—C7—H7A	109.5	C19—C20—H20	119.4
C3—C7—H7B	109.5	C15—C20—H20	119.4
H7A—C7—H7B	109.5	C26—C21—C22	118.38 (16)
C3—C7—H7C	109.5	C26—C21—C1	119.97 (15)
H7A—C7—H7C	109.5	C22—C21—C1	121.63 (15)
H7B—C7—H7C	109.5	C23—C22—C21	120.38 (16)
C4—C8—H8A	109.5	C23—C22—H22	119.8
C4—C8—H8B	109.5	C21—C22—H22	119.8
H8A—C8—H8B	109.5	C24—C23—C22	120.70 (17)
C4—C8—H8C	109.5	C24—C23—H23	119.6
H8A—C8—H8C	109.5	C22—C23—H23	119.6
H8B—C8—H8C	109.5	C23—C24—C25	119.19 (17)
C10—C9—C14	118.23 (16)	C23—C24—H24	120.4
C10—C9—C5	120.13 (15)	C25—C24—H24	120.4
C14—C9—C5	121.62 (15)	C26—C25—C24	120.39 (17)
C11—C10—C9	121.18 (17)	C26—C25—H25	119.8
C11—C10—H10	119.4	C24—C25—H25	119.8
C9—C10—H10	119.4	C25—C26—C21	120.93 (16)
C10—C11—C12	119.97 (18)	C25—C26—H26	119.5

C10—C11—H11	120.0	C21—C26—H26	119.5
C12—C11—H11	120.0		
C5—S1—C1—C2	-29.19 (15)	C10—C11—C12—C13	0.3 (3)
C5—S1—C1—C21	151.99 (12)	C11—C12—C13—C14	-0.1 (3)
C21—C1—C2—C6	-12.4 (3)	C12—C13—C14—C9	0.6 (3)
S1—C1—C2—C6	168.91 (13)	C10—C9—C14—C13	-1.3 (2)
C21—C1—C2—C3	167.85 (15)	C5—C9—C14—C13	-179.74 (16)
S1—C1—C2—C3	-10.8 (2)	C4—C3—C15—C16	-156.56 (15)
C1—C2—C3—C4	54.54 (19)	C2—C3—C15—C16	-37.7 (2)
C6—C2—C3—C4	-125.18 (16)	C7—C3—C15—C16	82.42 (18)
C1—C2—C3—C7	178.12 (15)	C4—C3—C15—C20	29.1 (2)
C6—C2—C3—C7	-1.6 (2)	C2—C3—C15—C20	147.93 (15)
C1—C2—C3—C15	-65.93 (19)	C7—C3—C15—C20	-91.92 (18)
C6—C2—C3—C15	114.35 (16)	C20—C15—C16—C17	-0.1 (3)
C2—C3—C4—C5	-54.04 (19)	C3—C15—C16—C17	-174.63 (16)
C7—C3—C4—C5	-177.52 (15)	C15—C16—C17—C18	-0.1 (3)
C15—C3—C4—C5	65.65 (19)	C16—C17—C18—C19	0.2 (3)
C2—C3—C4—C8	125.17 (16)	C17—C18—C19—C20	-0.1 (3)
C7—C3—C4—C8	1.7 (2)	C18—C19—C20—C15	-0.1 (3)
C15—C3—C4—C8	-115.13 (16)	C16—C15—C20—C19	0.2 (3)
C8—C4—C5—C9	11.4 (3)	C3—C15—C20—C19	174.69 (15)
C3—C4—C5—C9	-169.35 (15)	C2—C1—C21—C26	-49.4 (2)
C8—C4—C5—S1	-169.24 (13)	S1—C1—C21—C26	129.31 (14)
C3—C4—C5—S1	10.0 (2)	C2—C1—C21—C22	132.24 (18)
C1—S1—C5—C4	29.71 (15)	S1—C1—C21—C22	-49.03 (19)
C1—S1—C5—C9	-150.90 (12)	C26—C21—C22—C23	2.0 (2)
C4—C5—C9—C10	50.0 (2)	C1—C21—C22—C23	-179.63 (16)
S1—C5—C9—C10	-129.34 (15)	C21—C22—C23—C24	-1.1 (3)
C4—C5—C9—C14	-131.58 (18)	C22—C23—C24—C25	0.2 (3)
S1—C5—C9—C14	49.07 (19)	C23—C24—C25—C26	-0.2 (3)
C14—C9—C10—C11	1.5 (3)	C24—C25—C26—C21	1.1 (3)
C5—C9—C10—C11	179.99 (16)	C22—C21—C26—C25	-2.0 (3)
C9—C10—C11—C12	-1.0 (3)	C1—C21—C26—C25	179.61 (16)