

Bis[4-bromo-2-(cyclopentylimino-methyl)phenolato]copper(II)

Bang-Hong Cai

Department of Chemistry, Jiaying University, Meizhou Guangdong 514015, People's Republic of China

Correspondence e-mail: banghong_cai@163.com


Received 20 February 2009; accepted 23 February 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.039; wR factor = 0.098; data-to-parameter ratio = 18.6.

The title compound, $[\text{Cu}(\text{C}_{12}\text{H}_{13}\text{BrNO})_2]$, was prepared by the reaction of 5-bromosalicylaldehyde, cyclopentylamine and copper(II) acetate in an ethanol solution. The Cu^{II} atom lies on an inversion center and is four-coordinated in a square-planar geometry by two N and two O atoms from two 4-bromo-2-(cyclopentyliminomethyl)phenolate Schiff base ligands.

Related literature

For background on Schiff base complexes, see: Costes *et al.* (2002); Erxleben (2001); Lacroix *et al.* (1996); Odoko *et al.* (2006); Ali *et al.* (2006). For related copper(II) complexes, see: Wang *et al.* (2007); Datta *et al.* (2008); Yusnita *et al.* (2008); Wang & Zheng (2007). For a related zinc(II) complex, see: Cai (2009).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{12}\text{H}_{13}\text{BrNO})_2]$
 $M_r = 597.83$
 Monoclinic, $P2_1/c$
 $a = 9.190$ (2) Å
 $b = 10.960$ (2) Å
 $c = 12.166$ (2) Å
 $\beta = 109.73$ (3)°

$V = 1153.5$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 4.44$ mm⁻¹
 $T = 298$ K
 $0.27 \times 0.23 \times 0.23$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.381$, $T_{\text{max}} = 0.429$
 (expected range = 0.320–0.361)

9709 measured reflections
 2636 independent reflections
 2003 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.098$
 $S = 1.02$
 2636 reflections

142 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.93$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.42$ e Å⁻³

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The author acknowledges financial support from Jiaying University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2099).

References

- Ali, H. M., Puvaneswary, S. & Ng, S. W. (2006). *Acta Cryst.* **E62**, m2737–m2738.
 Bruker (2002). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cai, B.-H. (2009). *Acta Cryst.* **E65**, m142.
 Costes, J.-P., Clemente-Juan, J. M., Dahan, F., Dumestre, F. & Tuchagues, J.-P. (2002). *Inorg. Chem.* **41**, 2886–2891.
 Datta, A., Huang, J.-H. & Lee, H. M. (2008). *Acta Cryst.* **E64**, m1497.
 Erxleben, A. (2001). *Inorg. Chem.* **40**, 208–213.
 Lacroix, P. G., Di Bella, S. & Ledoux, I. (1996). *Chem. Mater.* **8**, 541–545.
 Odoko, M., Tsuchida, N. & Okabe, N. (2006). *Acta Cryst.* **E62**, m710–m711.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, L., Dong, J.-F., Li, L.-Z., Li, L.-W. & Wang, D.-Q. (2007). *Acta Cryst.* **E63**, m1059–m1060.
 Wang, L.-G. & Zheng, Y.-F. (2007). *Acta Cryst.* **E63**, m390–m391.
 Yusnita, J., Ali, H. M., Puvaneswary, S., Robinson, W. T. & Ng, S. W. (2008). *Acta Cryst.* **E64**, m1039.

supplementary materials

Acta Cryst. (2009). E65, m339 [doi:10.1107/S1600536809006606]

Bis[4-bromo-2-(cyclopentyliminomethyl)phenolato]copper(II)

B.-H. Cai

Comment

Schiff bases are interesting ligands which form a large number of complexes with metal atoms (Costes *et al.*, 2002; Erxleben, 2001; Lacroix *et al.*, 1996; Odoko *et al.*, 2006; Ali *et al.*, 2006). The author has recently reported on the crystal structure of a zinc(II) complex with the Schiff base (2-morpholin-4-ylethyl)-(1-pyridin-2-ylmethylidene)amine (Cai, 2009). As a continuous of our work in this area, we report here on the crystal structure of the title copper(II) complex (Fig. 1), derived from the Schiff base 4-bromo-2-(cyclopentyliminomethyl)phenol.

In the centrosymmetric title complex the Cu^{II} atom, is located on an inversion center, and is four-coordinate in a square planar geometry with two nitrogen and two oxygen atoms from two Schiff base ligands. All the coordinate bond lengths are typical and comparable with those in similar copper(II) complexes (Wang *et al.*, 2007; Datta *et al.*, 2008; Yusnita *et al.*, 2008; Wang & Zheng, 2007).

Experimental

5-Bromosalicylaldehyde (0.2 mmol, 40.3 mg), cyclopentylamine (0.2 mmol, 17.0 mg), and copper(II) acetate monohydrate (0.1 mmol, 20.0 mg) were mixed in 20 ml of ethanol. The mixture was stirred for 2 h at rt, giving a blue solution. Single-crystals were formed by gradual evaporation of the solution in air after several days.

Refinement

H atoms were placed in calculated positions and treated as riding: C–H = 0.93–0.98 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids [unlabelled atoms are related to the labelled atoms by the symmetry operation $-x, 2-y, -z$].

Bis[4-bromo-2-(cyclopentyliminomethyl)phenolato]copper(II)

Crystal data

[Cu(C ₁₂ H ₁₃ BrNO) ₂]	$F_{000} = 598$
$M_r = 597.83$	$D_x = 1.721 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.190 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 10.960 (2) \text{ \AA}$	Cell parameters from 2505 reflections
$c = 12.166 (2) \text{ \AA}$	$\theta = 2.3\text{--}25.0^\circ$
$\beta = 109.73 (3)^\circ$	$\mu = 4.44 \text{ mm}^{-1}$
$V = 1153.5 (4) \text{ \AA}^3$	$T = 298 \text{ K}$
$Z = 2$	Block, blue
	$0.27 \times 0.23 \times 0.23 \text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	2636 independent reflections
Radiation source: fine-focus sealed tube	2003 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.035$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.381$, $T_{\text{max}} = 0.429$	$k = -14 \rightarrow 14$
9709 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.098$	$w = 1/[\sigma^2(F_o^2) + (0.0427P)^2 + 0.9221P]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
2636 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
142 parameters	$\Delta\rho_{\text{max}} = 0.93 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.41 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.0000	1.0000	0.0000	0.03400 (16)
Br1	-0.34783 (5)	0.60773 (4)	0.33661 (4)	0.06123 (17)
O1	-0.0883 (3)	1.00623 (19)	0.1197 (2)	0.0446 (6)
N1	0.0534 (3)	0.8203 (2)	0.0328 (2)	0.0341 (6)
C1	-0.1071 (4)	0.7941 (3)	0.1548 (3)	0.0344 (7)
C2	-0.1456 (4)	0.9175 (3)	0.1630 (3)	0.0348 (7)
C3	-0.2475 (4)	0.9435 (3)	0.2236 (3)	0.0435 (8)
H3	-0.2753	1.0242	0.2294	0.052*
C4	-0.3073 (4)	0.8541 (3)	0.2744 (3)	0.0465 (9)
H4	-0.3752	0.8741	0.3136	0.056*
C5	-0.2659 (4)	0.7337 (3)	0.2671 (3)	0.0422 (8)
C6	-0.1676 (4)	0.7034 (3)	0.2090 (3)	0.0391 (7)
H6	-0.1404	0.6222	0.2052	0.047*
C7	-0.0063 (4)	0.7551 (3)	0.0935 (3)	0.0374 (7)
H7	0.0184	0.6726	0.0987	0.045*
C8	0.1560 (4)	0.7602 (3)	-0.0219 (3)	0.0386 (7)
H8	0.1265	0.7915	-0.1019	0.046*
C9	0.3252 (4)	0.7923 (4)	0.0374 (4)	0.0603 (11)
H9A	0.3467	0.8740	0.0163	0.072*
H9B	0.3536	0.7874	0.1216	0.072*
C10	0.4115 (5)	0.6976 (5)	-0.0078 (6)	0.0901 (18)
H10A	0.4543	0.7346	-0.0626	0.108*
H10B	0.4955	0.6632	0.0563	0.108*
C11	0.2977 (5)	0.6002 (4)	-0.0671 (4)	0.0642 (12)
H11A	0.3412	0.5199	-0.0429	0.077*
H11B	0.2696	0.6065	-0.1512	0.077*
C12	0.1586 (5)	0.6206 (3)	-0.0307 (4)	0.0507 (9)
H12A	0.1713	0.5824	0.0438	0.061*
H12B	0.0651	0.5900	-0.0889	0.061*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0412 (3)	0.0242 (3)	0.0424 (3)	0.0024 (2)	0.0218 (3)	0.0028 (2)
Br1	0.0702 (3)	0.0547 (3)	0.0720 (3)	-0.0114 (2)	0.0414 (2)	0.0125 (2)
O1	0.0659 (16)	0.0251 (11)	0.0570 (15)	0.0009 (11)	0.0395 (13)	0.0012 (10)
N1	0.0374 (14)	0.0263 (13)	0.0425 (15)	0.0034 (11)	0.0185 (12)	0.0018 (11)

supplementary materials

C1	0.0377 (17)	0.0294 (16)	0.0387 (17)	0.0018 (13)	0.0163 (14)	0.0037 (13)
C2	0.0401 (18)	0.0290 (15)	0.0373 (17)	0.0023 (13)	0.0159 (14)	0.0031 (13)
C3	0.056 (2)	0.0336 (17)	0.050 (2)	0.0058 (16)	0.0300 (18)	-0.0006 (16)
C4	0.050 (2)	0.049 (2)	0.049 (2)	0.0019 (17)	0.0287 (18)	0.0009 (17)
C5	0.0443 (19)	0.0424 (19)	0.0442 (19)	-0.0079 (15)	0.0205 (16)	0.0070 (15)
C6	0.0461 (19)	0.0289 (16)	0.0441 (19)	-0.0008 (14)	0.0175 (15)	0.0036 (14)
C7	0.0406 (17)	0.0270 (16)	0.0470 (19)	0.0064 (13)	0.0177 (15)	0.0040 (14)
C8	0.0421 (18)	0.0292 (16)	0.0489 (19)	0.0060 (14)	0.0212 (16)	0.0012 (14)
C9	0.043 (2)	0.056 (2)	0.085 (3)	-0.0012 (19)	0.026 (2)	-0.024 (2)
C10	0.051 (3)	0.082 (3)	0.147 (5)	-0.006 (2)	0.046 (3)	-0.051 (4)
C11	0.066 (3)	0.049 (2)	0.085 (3)	0.012 (2)	0.036 (2)	-0.010 (2)
C12	0.059 (2)	0.0349 (19)	0.064 (2)	0.0030 (17)	0.028 (2)	-0.0058 (17)

Geometric parameters (Å, °)

Cu1—O1 ⁱ	1.892 (2)	C6—H6	0.9300
Cu1—O1	1.892 (2)	C7—H7	0.9300
Cu1—N1	2.036 (2)	C8—C9	1.518 (5)
Cu1—N1 ⁱ	2.036 (2)	C8—C12	1.534 (4)
Br1—C5	1.902 (3)	C8—H8	0.9800
O1—C2	1.299 (4)	C9—C10	1.518 (5)
N1—C7	1.278 (4)	C9—H9A	0.9700
N1—C8	1.480 (4)	C9—H9B	0.9700
C1—C6	1.407 (4)	C10—C11	1.498 (6)
C1—C2	1.411 (4)	C10—H10A	0.9700
C1—C7	1.436 (4)	C10—H10B	0.9700
C2—C3	1.403 (4)	C11—C12	1.504 (6)
C3—C4	1.369 (5)	C11—H11A	0.9700
C3—H3	0.9300	C11—H11B	0.9700
C4—C5	1.385 (5)	C12—H12A	0.9700
C4—H4	0.9300	C12—H12B	0.9700
C5—C6	1.362 (5)		
O1 ⁱ —Cu1—O1	180.0	N1—C8—C9	112.9 (3)
O1 ⁱ —Cu1—N1	88.74 (10)	N1—C8—C12	120.2 (3)
O1—Cu1—N1	91.26 (10)	C9—C8—C12	103.1 (3)
O1 ⁱ —Cu1—N1 ⁱ	91.26 (10)	N1—C8—H8	106.6
O1—Cu1—N1 ⁱ	88.74 (10)	C9—C8—H8	106.6
N1—Cu1—N1 ⁱ	180.0	C12—C8—H8	106.6
C2—O1—Cu1	128.5 (2)	C10—C9—C8	104.2 (3)
C7—N1—C8	118.3 (3)	C10—C9—H9A	110.9
C7—N1—Cu1	122.1 (2)	C8—C9—H9A	110.9
C8—N1—Cu1	119.42 (19)	C10—C9—H9B	110.9
C6—C1—C2	119.7 (3)	C8—C9—H9B	110.9
C6—C1—C7	117.4 (3)	H9A—C9—H9B	108.9
C2—C1—C7	122.9 (3)	C11—C10—C9	107.3 (3)
O1—C2—C3	119.7 (3)	C11—C10—H10A	110.3
O1—C2—C1	122.9 (3)	C9—C10—H10A	110.3

C3—C2—C1	117.3 (3)	C11—C10—H10B	110.3
C4—C3—C2	122.3 (3)	C9—C10—H10B	110.3
C4—C3—H3	118.9	H10A—C10—H10B	108.5
C2—C3—H3	118.9	C10—C11—C12	106.1 (3)
C3—C4—C5	119.5 (3)	C10—C11—H11A	110.5
C3—C4—H4	120.3	C12—C11—H11A	110.5
C5—C4—H4	120.3	C10—C11—H11B	110.5
C6—C5—C4	120.7 (3)	C12—C11—H11B	110.5
C6—C5—Br1	119.0 (3)	H11A—C11—H11B	108.7
C4—C5—Br1	120.3 (3)	C11—C12—C8	101.8 (3)
C5—C6—C1	120.5 (3)	C11—C12—H12A	111.4
C5—C6—H6	119.7	C8—C12—H12A	111.4
C1—C6—H6	119.7	C11—C12—H12B	111.4
N1—C7—C1	127.8 (3)	C8—C12—H12B	111.4
N1—C7—H7	116.1	H12A—C12—H12B	109.3
C1—C7—H7	116.1		

Symmetry codes: (i) $-x, -y+2, -z$.

