

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Benzyloxy-1-naphthaldehyde

Rong Gao,^a Wen-Hong Li,^a Peng Liu^b and Ping-An Wang^{b*}

^aDepartment of Chemical Engineering, Northwest University, Taibai North Road 229, 710069 Xi-An, People's Republic of China, and ^bDepartment of Chemistry, School of Pharmacy, Fourth Military Medical University, Changle West Road 17, 710032 Xi-An, People's Republic of China

Correspondence e-mail: ping_an1718@yahoo.com.cn


Received 25 December 2008; accepted 7 February 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.084; wR factor = 0.285; data-to-parameter ratio = 12.5.

In the title compound, $\text{C}_{18}\text{H}_{14}\text{O}_2$, the dihedral angle between the phenyl and naphthyl ring systems is $21.8(3)^\circ$. The packing of molecules in the crystal structure is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the preparation of 2-benzyloxy-1-naphthaldehyde, see: Quideau *et al.* (2001). For synthetic use of the title compound, see: Knight & Little (2001).



Experimental

Crystal data

 $\text{C}_{18}\text{H}_{14}\text{O}_2$ $M_r = 262.29$ Monoclinic, $P2_1/c$ $a = 10.427(7)$ Å $b = 8.128(6)$ Å $c = 15.787(11)$ Å $\beta = 94.746(11)^\circ$ $V = 1333.3(16)$ Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.08$ mm⁻¹ $T = 296$ K $0.39 \times 0.26 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2005)

 $T_{\min} = 0.968$, $T_{\max} = 0.987$

5088 measured reflections

2262 independent reflections

1354 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.038$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.084$ $wR(F^2) = 0.285$ $S = 1.04$

2262 reflections

181 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.41$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.53$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C12}-\text{H12A}\cdots\text{O1}^{\dagger}$	0.97	2.48	3.381 (4)	155
$\text{C14}-\text{H14}\cdots\text{O1}^{\dagger}$	0.93	2.72	3.544 (5)	148

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *Mercury* (Macrae *et al.*, 2006) and *CAMERON* (Watkin *et al.*, 1996).

We thank the Natural Science Foundation of China (grant No. 20802092) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2303).

References

- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2002). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Knight, D. W. & Little, P. B. (2001). *J. Chem. Soc. Perkin Trans. 1*, pp. 1771–1777.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Quideau, S., Pouységú, L., Oxoby, M. & Looney, M. A. (2001). *Tetrahedron*, **57**, 319–329.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory, Oxford, England.

supplementary materials

Acta Cryst. (2009). E65, o534 [doi:10.1107/S1600536809004486]

2-Benzoyloxy-1-naphthaldehyde

R. Gao, W.-H. Li, P. Liu and P.-A. Wang

Comment

The title compound, 2-benzoyloxy-1-naphthaldehyde, was obtained by benzylation of 2-hydroxy-1-naphthaldehyde with benzyl bromide (Quideau *et al.*, 2001) and used for alkylation of position 4 in the naphthyl ring system. It has also been used for the intramolecular trapping of benzyne to yield some novel xanthenes (Knight & Little, 2001).

In the title compound, C₁₈H₁₄O₂, the dihedral angle between the phenyl and naphthyl ring systems is 21.8 (3)°. The packing of molecules in the crystal structure is stabilized by weak intermolecular C—H···O hydrogen bonds.

Experimental

To a stirred solution of commercially available 2-hydroxy-1-naphthaldehyde (4.30 g, 25.0 mmol) in *N,N*-dimethylformamide (100.0 cm³) was added potassium carbonate (3.82 g, 27.6 mmol) and benzyl bromide (3.0 cm³, 25.0 mmol), and the mixture was heated for 4 h at 90–100°C. The solution was filtered through celite and the solvent removed *in vacuo*. The residue was dissolved with Et₂O (160 cm³), washed with 1 M NaOH (110 cm³), brine (2 × 110 cm³), and dried over Na₂SO₄. Evaporation of the solvent afforded the title compound as a light yellow powder (6.0 g, 91%). The melting point and the spectroscopic data of the title compound were consisted with the reported literature (Quideau *et al.*, 2001).

Refinement

All H atoms were placed in calculated positions and refined as riding, with C—H = 0.93–0.97 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The values of $R[F^2 > 2\sigma(F^2)]$ and $wR(F^2)$ are 0.084 and 0.285, respectively; these high values may be due to the poor quality of the crystals.

Figures




Fig. 1. The molecular structure of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. The packing of the title compound, viewed down the *b* axis. Dotted lines indicate hydrogen bonds.

2-Benzoyloxy-1-naphthaldehyde

Crystal data

$C_{18}H_{14}O_2$	$F_{000} = 552$
$M_r = 262.29$	$D_x = 1.307 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Melting point: 393(1) K
Hall symbol: -P 2ybc	Mo $K\alpha$ radiation
$a = 10.427 (7) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 8.128 (6) \text{ \AA}$	Cell parameters from 1554 reflections
$c = 15.787 (11) \text{ \AA}$	$\theta = 2.6\text{--}24.3^\circ$
$\beta = 94.746 (11)^\circ$	$\mu = 0.08 \text{ mm}^{-1}$
$V = 1333.3 (16) \text{ \AA}^3$	$T = 296 \text{ K}$
$Z = 4$	Block, colourless
	$0.39 \times 0.26 \times 0.16 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	2262 independent reflections
Radiation source: fine-focus sealed tube	1354 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.038$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 25.1^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -7 \rightarrow 12$
$T_{\text{min}} = 0.968$, $T_{\text{max}} = 0.987$	$k = -9 \rightarrow 6$
5088 measured reflections	$l = -18 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.084$	H-atom parameters constrained
$wR(F^2) = 0.285$	$w = 1/[\sigma^2(F_o^2) + (0.18P)^2 + 0.612P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
2262 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
181 parameters	$\Delta\rho_{\text{max}} = 0.41 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.53 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6120 (3)	0.2057 (4)	0.65206 (15)	0.0658 (10)
O2	0.6321 (2)	0.1063 (4)	0.41483 (14)	0.0531 (8)
C1	0.4002 (3)	0.3376 (5)	0.5290 (2)	0.0411 (9)
C2	0.3808 (4)	0.4033 (5)	0.6101 (2)	0.0518 (11)
H2	0.4423	0.3837	0.6551	0.062*
C3	0.2749 (4)	0.4942 (6)	0.6241 (3)	0.0630 (12)
H3	0.2641	0.5335	0.6784	0.076*
C4	0.1821 (4)	0.5289 (7)	0.5572 (3)	0.0716 (14)
H4	0.1103	0.5918	0.5671	0.086*
C5	0.1967 (4)	0.4710 (5)	0.4783 (3)	0.0577 (11)
H5	0.1349	0.4952	0.4343	0.069*
C6	0.3053 (3)	0.3737 (6)	0.4616 (2)	0.0504 (11)
C7	0.3228 (3)	0.3153 (5)	0.3794 (2)	0.0523 (11)
H7	0.2608	0.3395	0.3354	0.063*
C8	0.4275 (3)	0.2244 (6)	0.3622 (2)	0.0539 (11)
H8	0.4354	0.1851	0.3075	0.065*
C9	0.5238 (3)	0.1904 (5)	0.4280 (2)	0.0431 (10)
C10	0.5100 (3)	0.2424 (5)	0.5110 (2)	0.0392 (9)
C11	0.6122 (3)	0.1927 (6)	0.5756 (2)	0.0514 (11)
H11	0.6853	0.1459	0.5556	0.062*
C12	0.6621 (3)	0.0652 (6)	0.3305 (2)	0.0575 (12)
H12A	0.6381	0.1548	0.2918	0.069*
H12B	0.6154	-0.0326	0.3108	0.069*
C13	0.8049 (3)	0.0350 (5)	0.3337 (2)	0.0468 (10)
C14	0.8757 (4)	0.1189 (6)	0.2765 (2)	0.0571 (12)
H14	0.8348	0.1902	0.2368	0.069*
C15	1.0079 (4)	0.0954 (7)	0.2792 (3)	0.0713 (15)
H15	1.0549	0.1484	0.2397	0.086*
C16	1.0699 (4)	-0.0047 (6)	0.3390 (3)	0.0635 (13)
H16	1.1589	-0.0163	0.3412	0.076*
C17	1.0006 (4)	-0.0884 (6)	0.3960 (3)	0.0630 (12)
H17	1.0423	-0.1582	0.4360	0.076*
C18	0.8681 (4)	-0.0675 (6)	0.3932 (2)	0.0579 (12)

supplementary materials

H18 0.8214 -0.1232 0.4319 0.069*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0596 (17)	0.099 (3)	0.0374 (14)	0.0078 (16)	-0.0065 (11)	-0.0031 (14)
O2	0.0498 (15)	0.075 (2)	0.0352 (13)	0.0178 (14)	0.0062 (10)	-0.0012 (12)
C1	0.0352 (18)	0.047 (3)	0.0415 (17)	-0.0085 (16)	0.0054 (14)	0.0049 (17)
C2	0.045 (2)	0.062 (3)	0.050 (2)	-0.003 (2)	0.0112 (16)	-0.0008 (19)
C3	0.058 (2)	0.066 (3)	0.068 (3)	-0.003 (2)	0.023 (2)	-0.001 (2)
C4	0.046 (2)	0.082 (4)	0.090 (3)	0.008 (2)	0.024 (2)	0.003 (3)
C5	0.043 (2)	0.056 (3)	0.075 (3)	-0.0013 (19)	0.0041 (18)	0.008 (2)
C6	0.0307 (17)	0.068 (3)	0.052 (2)	-0.0035 (17)	0.0040 (15)	0.011 (2)
C7	0.040 (2)	0.067 (3)	0.047 (2)	-0.0013 (19)	-0.0074 (15)	0.010 (2)
C8	0.045 (2)	0.082 (3)	0.0340 (18)	-0.002 (2)	-0.0007 (15)	0.0032 (18)
C9	0.0345 (17)	0.055 (3)	0.0393 (18)	-0.0040 (17)	0.0035 (13)	0.0078 (17)
C10	0.0328 (17)	0.047 (2)	0.0373 (17)	-0.0047 (15)	0.0012 (13)	0.0028 (16)
C11	0.0386 (19)	0.076 (3)	0.0386 (19)	0.0002 (19)	-0.0005 (14)	0.0038 (19)
C12	0.043 (2)	0.096 (4)	0.0337 (17)	-0.003 (2)	0.0069 (14)	-0.004 (2)
C13	0.0395 (18)	0.066 (3)	0.0355 (16)	-0.0014 (18)	0.0055 (14)	-0.0021 (18)
C14	0.045 (2)	0.081 (4)	0.047 (2)	0.005 (2)	0.0070 (16)	0.015 (2)
C15	0.044 (2)	0.107 (4)	0.064 (3)	0.003 (2)	0.0155 (19)	0.017 (3)
C16	0.043 (2)	0.087 (4)	0.061 (2)	0.010 (2)	0.0049 (18)	0.002 (2)
C17	0.056 (2)	0.075 (3)	0.057 (2)	0.015 (2)	-0.0031 (18)	0.008 (2)
C18	0.054 (2)	0.071 (3)	0.050 (2)	-0.002 (2)	0.0091 (17)	0.010 (2)

Geometric parameters (\AA , $^\circ$)

O1—C11	1.211 (4)	C8—H8	0.9300
O2—C9	1.351 (4)	C9—C10	1.395 (5)
O2—C12	1.433 (4)	C10—C11	1.471 (5)
C1—C2	1.417 (5)	C11—H11	0.9300
C1—C6	1.422 (5)	C12—C13	1.506 (5)
C1—C10	1.430 (5)	C12—H12A	0.9700
C2—C3	1.362 (6)	C12—H12B	0.9700
C2—H2	0.9300	C13—C18	1.382 (6)
C3—C4	1.400 (6)	C13—C14	1.391 (5)
C3—H3	0.9300	C14—C15	1.389 (5)
C4—C5	1.351 (6)	C14—H14	0.9300
C4—H4	0.9300	C15—C16	1.367 (6)
C5—C6	1.424 (5)	C15—H15	0.9300
C5—H5	0.9300	C16—C17	1.380 (6)
C6—C7	1.407 (5)	C16—H16	0.9300
C7—C8	1.365 (5)	C17—C18	1.389 (5)
C7—H7	0.9300	C17—H17	0.9300
C8—C9	1.412 (5)	C18—H18	0.9300
C9—O2—C12	120.7 (3)	C9—C10—C11	116.3 (3)
C2—C1—C6	117.1 (3)	C1—C10—C11	123.8 (3)

C2—C1—C10	124.1 (3)	O1—C11—C10	127.3 (4)
C6—C1—C10	118.8 (3)	O1—C11—H11	116.3
C3—C2—C1	121.9 (4)	C10—C11—H11	116.3
C3—C2—H2	119.0	O2—C12—C13	107.3 (3)
C1—C2—H2	119.0	O2—C12—H12A	110.2
C2—C3—C4	120.4 (4)	C13—C12—H12A	110.2
C2—C3—H3	119.8	O2—C12—H12B	110.2
C4—C3—H3	119.8	C13—C12—H12B	110.2
C5—C4—C3	120.1 (4)	H12A—C12—H12B	108.5
C5—C4—H4	120.0	C18—C13—C14	119.2 (3)
C3—C4—H4	120.0	C18—C13—C12	122.4 (3)
C4—C5—C6	121.1 (4)	C14—C13—C12	118.3 (3)
C4—C5—H5	119.5	C15—C14—C13	119.4 (4)
C6—C5—H5	119.5	C15—C14—H14	120.3
C7—C6—C1	119.0 (3)	C13—C14—H14	120.3
C7—C6—C5	121.6 (3)	C16—C15—C14	121.0 (4)
C1—C6—C5	119.4 (4)	C16—C15—H15	119.5
C8—C7—C6	122.2 (3)	C14—C15—H15	119.5
C8—C7—H7	118.9	C15—C16—C17	120.0 (4)
C6—C7—H7	118.9	C15—C16—H16	120.0
C7—C8—C9	119.4 (3)	C17—C16—H16	120.0
C7—C8—H8	120.3	C16—C17—C18	119.5 (4)
C9—C8—H8	120.3	C16—C17—H17	120.2
O2—C9—C10	116.8 (3)	C18—C17—H17	120.2
O2—C9—C8	122.5 (3)	C13—C18—C17	120.8 (4)
C10—C9—C8	120.6 (3)	C13—C18—H18	119.6
C9—C10—C1	119.9 (3)	C17—C18—H18	119.6
C6—C1—C2—C3	1.4 (6)	O2—C9—C10—C11	-2.9 (5)
C10—C1—C2—C3	179.6 (4)	C8—C9—C10—C11	176.6 (4)
C1—C2—C3—C4	-1.4 (7)	C2—C1—C10—C9	-176.6 (4)
C2—C3—C4—C5	0.5 (7)	C6—C1—C10—C9	1.5 (5)
C3—C4—C5—C6	0.4 (7)	C2—C1—C10—C11	4.0 (6)
C2—C1—C6—C7	178.0 (4)	C6—C1—C10—C11	-177.9 (3)
C10—C1—C6—C7	-0.2 (6)	C9—C10—C11—O1	-169.9 (4)
C2—C1—C6—C5	-0.5 (5)	C1—C10—C11—O1	9.6 (7)
C10—C1—C6—C5	-178.8 (3)	C9—O2—C12—C13	158.2 (3)
C4—C5—C6—C7	-178.8 (4)	O2—C12—C13—C18	49.5 (6)
C4—C5—C6—C1	-0.3 (6)	O2—C12—C13—C14	-127.9 (4)
C1—C6—C7—C8	0.3 (6)	C18—C13—C14—C15	1.4 (6)
C5—C6—C7—C8	178.8 (4)	C12—C13—C14—C15	178.8 (4)
C6—C7—C8—C9	-1.6 (6)	C13—C14—C15—C16	-2.2 (7)
C12—O2—C9—C10	-172.4 (3)	C14—C15—C16—C17	2.2 (8)
C12—O2—C9—C8	8.1 (6)	C15—C16—C17—C18	-1.3 (7)
C7—C8—C9—O2	-177.6 (4)	C14—C13—C18—C17	-0.6 (7)
C7—C8—C9—C10	2.9 (6)	C12—C13—C18—C17	-177.9 (4)
O2—C9—C10—C1	177.6 (3)	C16—C17—C18—C13	0.5 (7)
C8—C9—C10—C1	-2.8 (6)		

supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C12—H12A···O1 ⁱ	0.97	2.48	3.381 (4)	155
C14—H14···O1 ⁱ	0.93	2.72	3.544 (5)	148

Symmetry codes: (i) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

