

Tetraqua(2,2'-diamino-4,4'-bi-1,3-thiazole- κ^2N^3,N^3')nickel(II) bis(pyridine-2,6-dicarboxylato- κ^3O^2,N,O^6)nickel(II) trihydrate

 Bing-Xin Liu,^{a*} Yan-Ping Yu,^a Zen Cao^a and Liang-Jun Zhang^b

^aDepartment of Chemistry, Shanghai University, Shanghai 200444, People's Republic of China, and ^bDepartment of Petroleum and Chemical Industry, Guangxi Vocational and Technical Institute of Industry, People's Republic of China
Correspondence e-mail: r5744011@yahoo.com.cn


Received 23 October 2008; accepted 9 February 2009

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.034; wR factor = 0.084; data-to-parameter ratio = 12.3.

The crystal structure of the title compound, $[Ni(C_6H_6N_4S_2)(H_2O)_4][Ni(C_7H_3NO_4)_2] \cdot 3H_2O$, consists of Ni^{II} complex cations, Ni^{II} complex anions and lattice water molecules. The Ni^{II} ions in both the complex cation and anion assume a distorted octahedral coordination geometry. O—H...O, N—H...O and C—H...S hydrogen bonds occur in the crystal structure.

Related literature

For general background, see: Waring (1981); Fisher *et al.* (1985). For a related structure, see: Liu *et al.* (2003); Zhang *et al.* (2006). For synthesis, see: Erlenmeyer (1948).



Experimental

Crystal data

$[Ni(C_6H_6N_4S_2)(H_2O)_4][Ni(C_7H_3NO_4)_2] \cdot 3H_2O$
 $M_r = 772.01$
 Triclinic, $P\bar{1}$
 $a = 11.4756$ (13) Å
 $b = 11.5609$ (13) Å
 $c = 13.2667$ (15) Å
 $\alpha = 65.3590$ (10)°
 $\beta = 82.1140$ (11)°
 $\gamma = 66.0180$ (10)°

$V = 1460.6$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

$\mu = 1.51$ mm⁻¹
 $T = 295$ K
 $0.23 \times 0.18 \times 0.15$ mm

Data collection

Bruker APEX CCD diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{min} = 0.710$, $T_{max} = 0.795$

7580 measured reflections
 5052 independent reflections
 4181 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.084$
 $S = 1.05$
 5052 reflections

412 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.31$ e Å⁻³
 $\Delta\rho_{min} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1—H1A...O14 ⁱ	0.86	1.75	2.606 (3)	174
O1—H1B...O21 ⁱⁱ	0.82	1.92	2.747 (3)	176
O2—H2A...O12 ⁱⁱⁱ	0.77	2.01	2.772 (3)	169
O2—H2B...O22 ⁱⁱ	0.76	2.12	2.874 (3)	170
O3—H3A...O3W ⁱⁱⁱ	0.78	1.98	2.742 (3)	169
O3—H3B...O11 ⁱⁱⁱ	0.75	1.94	2.683 (3)	173
O4—H4A...O24 ^{iv}	0.83	1.89	2.717 (4)	179
O4—H4B...O1W	0.74	2.16	2.820 (4)	149
O1W—H1WA...O23 ^{iv}	0.91	2.01	2.907 (4)	166
O1W—H1WB...O24 ⁱⁱⁱ	0.75	2.41	3.150 (5)	169
O2W—H2WB...O3W ^v	0.72	2.41	3.104 (4)	162
O2W—H2WA...O14	0.79	2.02	2.815 (4)	174
O3W—H3WA...O22 ^{vi}	0.73	2.24	2.934 (3)	158
O3W—H3WB...O21	0.88	2.10	2.892 (3)	149
N32—H32A...O12	0.98	1.97	2.940 (5)	169
N32—H32B...O2	0.95	2.19	2.996 (5)	142
N34—H34A...O2W ⁱⁱ	0.92	1.98	2.886 (4)	168
N34—H34B...O3	0.92	2.12	2.952 (4)	149
C12—H12...S31	0.93	2.71	3.631 (4)	170

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x, y, z + 1$; (iii) $-x, -y + 1, -z + 1$; (iv) $x, y + 1, z$; (v) $-x + 1, -y, -z$; (vi) $-x, -y + 1, -z$; (vii) $x, y + 1, z + 1$.

Data collection: SMART (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SIR92 (Altomare *et al.*, 1993); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

This project was supported by the Educational Development Foundation of Shanghai Educational Committee, China (grant No. AB0448).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2461).

References

- Altomare, A., Casciarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
 Bruker (2004). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Erlenmeyer, H. (1948). *Helv. Chim. Acta*, **31**, 206–210.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Fisher, L. M., Kurod, R. & Sakai, T. (1985). *Biochemistry*, **24**, 3199–3207.
 Liu, J.-G., Xu, D.-J., Sun, W.-L., Wu, Z.-Y., Xu, Y.-Z., Wu, J.-Y. & Chiang, M. Y. (2003). *J. Coord. Chem.* **56**, 71–76.

Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Waring, M. J. (1981). *Annu. Rev. Biochem.* **50**, 159–192.

Zhang, L.-J., Liu, B.-X., Ge, H.-Q. & Xu, D.-J. (2006). *Acta Cryst.* **E62**, m2180–m2182.

supplementary materials

Acta Cryst. (2009). E65, m277-m278 [doi:10.1107/S1600536809004589]

Tetraaqua(2,2'-diamino-4,4'-bi-1,3-thiazole- $\kappa^2N^3,N^{3'}$)nickel(II) bis(pyridine-2,6-dicarboxylato- κ^3O^2,N,O^6)nickel(II) trihydrate

B.-X. Liu, Y.-P. Yu, Z. Cao and L.-J. Zhang

Comment

Transition metal complexes of 2,2'-diamino-4,4'-bi-1,3-thiazole (DABT) have shown potential application in some fields, for example, a Co^{II} complex and a Ni^{II} complex with the DABT ligand have been found to be effective inhibitors of DNA synthesis of tumor cell (Waring, 1981; Fisher *et al.*, 1985). As part of serial structural investigation of metal complexes with DABT, the title Ni^{II} complex was prepared in the laboratory and its X-ray structure is presented here.

The crystal of title compound consists of Ni^{II} complex cations, Ni^{II} complex anions and lattice water molecules (Fig. 1). Within the complex cation, the Ni^{II} ion is coordinated by a DABT ligand and four water molecules in a distorted octahedral geometry. The thiazole rings of DABT are approximately coplanar with the dihedral angle of 6.2 (2)° between thiazole rings. The average of Ni—N bond distance of 2.082 (2) Å is comparable to the Ni—N bond distance of 2.103 (4) Å found in [Ni(C₆H₆N₄S₂)₂(H₂O)₂][Ni(C₇H₃NO₄)₂].5(H₂O) (Zhang *et al.*, 2006) and 2.113 (2) Å found in [Ni(C₈H₄O₄)(C₆H₆N₄S₂)₂].3.5H₂O (Liu *et al.*, 2003).

In the complex anion, the Ni^{II} ion is chelated by two pyridinedicarboxylate (pdc) dianions with a distorted octahedral geometry. Two planar pdc ligands are nearly perpendicular to each other, the dihedral angle being 88.46 (8)°.

The extensive hydrogen bonding between lattice water molecules, complex cations and complex anions helps to stabilize the crystal structure (Table 1).

Experimental

The microcrystals of DABT were obtained in the manner reported by Erlenmeyer (1948). An aqueous solution (20 ml) containing DABT (0.20 g, 1 mmol) and NiCl₂ (0.13 g, 1 mmol) was mixed with another aqueous solution (10 ml) of 2,6-pyridinedicarboxylic acid (0.17 g, 1 mmol) and NaOH (0.08 g, 2 mmol). The mixture was refluxed for 6 h. The solution was filtered after cooling to room temperature. Green single crystals were obtained from the filtrate after 30 d.

Refinement

Aromatic H atoms were placed in calculated positions with C—H = 0.93 Å and were included in the final cycles of refinement in riding mode with $U_{iso}(H) = 1.2U_{eq}(C)$. H atoms on N atom of amino group of DABT and on O atom of water molecules were located in a difference Fourier map and also included in the final cycles of refinement in riding mode with $U_{iso}(H) = 1.5U_{eq}(O)$ or $1.2U_{eq}(N)$.

Figures




Fig. 1. The molecular structure of (I) with 30% probability displacement ellipsoids (arbitrary spheres for H atoms), dashed lines showing the hydrogen bonding.

Tetraaqua(2,2'-diamino-4,4'-bi-1,3-thiazole- $\kappa^2N^3,N^{3'}$)nickel(II) bis(pyridine-2,6-dicarboxylato- κ^3O^2,N,O^6)nickel(II) trihydrate

Crystal data

$[\text{Ni}(\text{C}_6\text{H}_6\text{N}_4\text{S}_2)(\text{H}_2\text{O})_4][\text{Ni}(\text{C}_7\text{H}_3\text{NO}_4)_2] \cdot 3\text{H}_2\text{O}$

$M_r = 772.01$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 11.4756$ (13) Å

$b = 11.5609$ (13) Å

$c = 13.2667$ (15) Å

$\alpha = 65.3590$ (10)°

$\beta = 82.1140$ (11)°

$\gamma = 66.0180$ (10)°

$V = 1460.6$ (3) Å³

$Z = 2$

$F_{000} = 792$

$D_x = 1.755$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 4980 reflections

$\theta = 2.0$ – 25.0 °

$\mu = 1.51$ mm⁻¹

$T = 295$ K

Prism, green

$0.23 \times 0.18 \times 0.15$ mm

Data collection

Bruker APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$ K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.710$, $T_{\max} = 0.795$

7580 measured reflections

5052 independent reflections

4181 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

$\theta_{\text{max}} = 25.0$ °

$\theta_{\text{min}} = 2.1$ °

$h = -13 \rightarrow 7$

$k = -13 \rightarrow 13$

$l = -15 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.084$

$S = 1.05$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0368P)^2 + 0.7271P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

5052 reflections $\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$
 412 parameters $\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.16770 (3)	0.07786 (4)	0.17826 (3)	0.02475 (11)
Ni2	0.23271 (3)	0.51033 (4)	0.71455 (3)	0.02385 (11)
O1	0.2614 (2)	0.3223 (2)	0.84334 (17)	0.0390 (6)
H1A	0.3396	0.2657	0.8611	0.059*
H1B	0.2134	0.3025	0.8940	0.059*
O2	0.06240 (19)	0.4957 (2)	0.68468 (17)	0.0343 (5)
H2A	-0.0007	0.5604	0.6691	0.051*
H2B	0.0430	0.4451	0.7356	0.051*
O3	0.13816 (18)	0.6134 (2)	0.81574 (16)	0.0319 (5)
H3A	0.1392	0.5667	0.8783	0.048*
H3B	0.0692	0.6533	0.7976	0.048*
O4	0.1730 (2)	0.7000 (2)	0.58611 (17)	0.0414 (6)
H4A	0.1238	0.7351	0.5318	0.062*
H4B	0.1773	0.7579	0.5939	0.062*
O11	0.10601 (19)	0.2234 (2)	0.25192 (17)	0.0337 (5)
O12	0.1791 (2)	0.2960 (2)	0.35197 (18)	0.0389 (5)
O13	0.29701 (19)	-0.0576 (2)	0.11358 (17)	0.0326 (5)
O14	0.50368 (19)	-0.1545 (2)	0.08943 (17)	0.0349 (5)
O21	0.10956 (19)	0.2430 (2)	0.01530 (16)	0.0311 (5)
O22	-0.0369 (2)	0.3366 (2)	-0.12130 (17)	0.0382 (5)
O23	0.1450 (2)	-0.0753 (2)	0.32817 (17)	0.0362 (5)
O24	0.0132 (3)	-0.1844 (3)	0.4064 (2)	0.0612 (8)
N11	0.3291 (2)	0.0700 (2)	0.21889 (19)	0.0232 (5)
N21	0.0100 (2)	0.0718 (2)	0.14401 (19)	0.0239 (5)
N31	0.3493 (2)	0.4052 (2)	0.62139 (19)	0.0290 (6)
N32	0.2175 (3)	0.3633 (4)	0.5323 (3)	0.0644 (10)
H32A	0.2169	0.3353	0.4722	0.077*
H32B	0.1419	0.4222	0.5525	0.077*

supplementary materials

N33	0.4079 (2)	0.5113 (2)	0.74188 (19)	0.0257 (5)
N34	0.3714 (3)	0.6488 (3)	0.8419 (2)	0.0467 (8)
H34A	0.4167	0.6570	0.8887	0.056*
H34B	0.2868	0.6592	0.8476	0.056*
S31	0.47085 (9)	0.26783 (9)	0.50388 (7)	0.0432 (2)
S32	0.61255 (8)	0.50673 (9)	0.80414 (8)	0.0412 (2)
C11	0.3283 (3)	0.1372 (3)	0.2798 (2)	0.0248 (6)
C12	0.4413 (3)	0.1217 (3)	0.3184 (3)	0.0343 (7)
H12	0.4412	0.1698	0.3597	0.041*
C13	0.5547 (3)	0.0337 (3)	0.2948 (3)	0.0360 (8)
H13	0.6320	0.0210	0.3209	0.043*
C14	0.5528 (3)	-0.0360 (3)	0.2316 (3)	0.0297 (7)
H14	0.6284	-0.0953	0.2148	0.036*
C15	0.4374 (3)	-0.0157 (3)	0.1945 (2)	0.0238 (6)
C16	0.1940 (3)	0.2267 (3)	0.2975 (2)	0.0279 (7)
C17	0.4124 (3)	-0.0821 (3)	0.1264 (2)	0.0263 (6)
C21	-0.0484 (3)	0.1524 (3)	0.0445 (2)	0.0238 (6)
C22	-0.1516 (3)	0.1387 (3)	0.0152 (2)	0.0316 (7)
H22	-0.1932	0.1958	-0.0540	0.038*
C23	-0.1914 (3)	0.0377 (4)	0.0917 (3)	0.0384 (8)
H23	-0.2598	0.0254	0.0737	0.046*
C24	-0.1295 (3)	-0.0449 (3)	0.1947 (3)	0.0356 (8)
H24	-0.1557	-0.1127	0.2467	0.043*
C25	-0.0284 (3)	-0.0246 (3)	0.2186 (2)	0.0279 (7)
C26	0.0113 (3)	0.2535 (3)	-0.0281 (2)	0.0267 (7)
C27	0.0489 (3)	-0.1018 (3)	0.3268 (3)	0.0361 (8)
C31	0.4781 (3)	0.3740 (3)	0.6331 (2)	0.0270 (7)
C32	0.5566 (3)	0.3003 (3)	0.5773 (3)	0.0382 (8)
H32	0.6450	0.2712	0.5783	0.046*
C33	0.3309 (3)	0.3552 (3)	0.5550 (3)	0.0364 (8)
C34	0.5109 (3)	0.4281 (3)	0.7014 (2)	0.0281 (7)
C35	0.6263 (3)	0.4128 (3)	0.7279 (3)	0.0372 (8)
H35	0.7033	0.3589	0.7079	0.045*
C36	0.4474 (3)	0.5613 (3)	0.7966 (3)	0.0307 (7)
O1W	0.1949 (3)	0.9552 (3)	0.5228 (2)	0.0749 (9)
H1WA	0.1810	0.9596	0.4548	0.112*
H1WB	0.1411	1.0142	0.5309	0.112*
O2W	0.5287 (2)	-0.3045 (3)	-0.0391 (2)	0.0537 (7)
H2WB	0.5946	-0.3508	-0.0270	0.081*
H2WA	0.5167	-0.2587	-0.0049	0.081*
O3W	0.1763 (2)	0.4471 (2)	0.03785 (18)	0.0483 (6)
H3WA	0.1475	0.4837	0.0745	0.072*
H3WB	0.1492	0.3796	0.0574	0.072*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0183 (2)	0.0276 (2)	0.0284 (2)	-0.00704 (16)	-0.00210 (15)	-0.01206 (16)

Ni2	0.0214 (2)	0.0248 (2)	0.0232 (2)	-0.00605 (16)	-0.00041 (15)	-0.00997 (16)
O1	0.0254 (12)	0.0335 (12)	0.0358 (12)	-0.0041 (10)	0.0056 (10)	-0.0017 (10)
O2	0.0263 (11)	0.0338 (12)	0.0377 (12)	-0.0071 (10)	-0.0021 (9)	-0.0130 (10)
O3	0.0244 (11)	0.0359 (12)	0.0290 (11)	-0.0038 (9)	-0.0016 (9)	-0.0138 (9)
O4	0.0510 (15)	0.0320 (12)	0.0317 (12)	-0.0134 (11)	-0.0126 (11)	-0.0025 (10)
O11	0.0204 (11)	0.0401 (12)	0.0423 (13)	-0.0039 (10)	-0.0023 (10)	-0.0245 (10)
O12	0.0338 (13)	0.0422 (13)	0.0471 (14)	-0.0064 (10)	-0.0004 (10)	-0.0312 (11)
O13	0.0244 (12)	0.0385 (12)	0.0417 (13)	-0.0083 (10)	-0.0025 (9)	-0.0247 (10)
O14	0.0282 (12)	0.0351 (12)	0.0417 (13)	-0.0035 (10)	0.0006 (10)	-0.0239 (10)
O21	0.0299 (12)	0.0327 (11)	0.0303 (11)	-0.0160 (10)	-0.0016 (9)	-0.0077 (9)
O22	0.0388 (13)	0.0392 (12)	0.0274 (12)	-0.0171 (11)	-0.0044 (10)	-0.0013 (10)
O23	0.0287 (12)	0.0432 (13)	0.0308 (12)	-0.0153 (10)	-0.0064 (9)	-0.0060 (10)
O24	0.0633 (18)	0.0762 (19)	0.0344 (14)	-0.0482 (16)	-0.0134 (12)	0.0120 (13)
N11	0.0191 (12)	0.0251 (12)	0.0261 (13)	-0.0069 (10)	-0.0008 (10)	-0.0117 (10)
N21	0.0195 (12)	0.0266 (13)	0.0228 (13)	-0.0079 (10)	-0.0002 (10)	-0.0080 (10)
N31	0.0305 (14)	0.0314 (14)	0.0260 (13)	-0.0096 (12)	0.0009 (11)	-0.0147 (11)
N32	0.048 (2)	0.098 (3)	0.077 (2)	-0.0195 (19)	-0.0009 (18)	-0.070 (2)
N33	0.0242 (13)	0.0241 (12)	0.0262 (13)	-0.0080 (11)	0.0006 (10)	-0.0090 (10)
N34	0.0357 (16)	0.0591 (19)	0.067 (2)	-0.0165 (15)	0.0037 (15)	-0.0468 (17)
S31	0.0550 (6)	0.0434 (5)	0.0343 (5)	-0.0156 (4)	0.0099 (4)	-0.0242 (4)
S32	0.0301 (5)	0.0451 (5)	0.0524 (5)	-0.0159 (4)	-0.0051 (4)	-0.0198 (4)
C11	0.0259 (16)	0.0219 (14)	0.0279 (15)	-0.0099 (12)	-0.0002 (12)	-0.0103 (12)
C12	0.0332 (18)	0.0415 (18)	0.0411 (19)	-0.0174 (15)	-0.0003 (15)	-0.0250 (16)
C13	0.0225 (16)	0.0424 (19)	0.050 (2)	-0.0126 (15)	-0.0024 (14)	-0.0234 (16)
C14	0.0181 (15)	0.0298 (16)	0.0389 (18)	-0.0074 (13)	0.0012 (13)	-0.0136 (14)
C15	0.0224 (15)	0.0229 (14)	0.0261 (15)	-0.0097 (12)	0.0024 (12)	-0.0092 (12)
C16	0.0262 (16)	0.0251 (15)	0.0264 (16)	-0.0051 (13)	-0.0019 (13)	-0.0088 (13)
C17	0.0281 (17)	0.0217 (15)	0.0257 (16)	-0.0074 (13)	-0.0010 (13)	-0.0079 (12)
C21	0.0186 (15)	0.0260 (15)	0.0254 (15)	-0.0061 (12)	-0.0004 (12)	-0.0111 (12)
C22	0.0252 (16)	0.0406 (18)	0.0270 (16)	-0.0118 (14)	-0.0045 (13)	-0.0107 (14)
C23	0.0301 (18)	0.057 (2)	0.0347 (18)	-0.0252 (17)	-0.0010 (14)	-0.0162 (16)
C24	0.0324 (18)	0.0446 (19)	0.0312 (17)	-0.0242 (16)	0.0031 (14)	-0.0081 (15)
C25	0.0269 (16)	0.0309 (16)	0.0237 (15)	-0.0112 (13)	-0.0014 (13)	-0.0081 (13)
C26	0.0226 (16)	0.0248 (15)	0.0304 (17)	-0.0061 (13)	0.0004 (13)	-0.0116 (13)
C27	0.0306 (18)	0.0411 (19)	0.0297 (18)	-0.0144 (15)	-0.0036 (14)	-0.0058 (15)
C31	0.0276 (16)	0.0228 (15)	0.0264 (16)	-0.0089 (13)	0.0051 (13)	-0.0079 (12)
C32	0.0375 (19)	0.0330 (17)	0.0402 (19)	-0.0105 (15)	0.0101 (15)	-0.0167 (15)
C33	0.042 (2)	0.0367 (18)	0.0312 (17)	-0.0121 (16)	0.0003 (15)	-0.0164 (15)
C34	0.0264 (16)	0.0223 (15)	0.0301 (16)	-0.0086 (13)	0.0020 (13)	-0.0066 (12)
C35	0.0270 (18)	0.0326 (17)	0.047 (2)	-0.0073 (14)	0.0024 (15)	-0.0152 (15)
C36	0.0271 (17)	0.0320 (17)	0.0353 (17)	-0.0139 (14)	0.0002 (13)	-0.0130 (14)
O1W	0.108 (3)	0.0634 (19)	0.0524 (17)	-0.0390 (18)	-0.0031 (17)	-0.0143 (15)
O2W	0.0598 (17)	0.0577 (16)	0.0551 (16)	-0.0190 (14)	-0.0104 (13)	-0.0327 (13)
O3W	0.0644 (17)	0.0424 (14)	0.0417 (14)	-0.0231 (13)	0.0052 (12)	-0.0187 (11)

Geometric parameters (Å, °)

Ni1—N11	1.956 (2)	N33—C34	1.394 (4)
Ni1—N21	1.960 (2)	N34—C36	1.339 (4)

supplementary materials

Ni1—O13	2.067 (2)	N34—H34A	0.9184
Ni1—O23	2.110 (2)	N34—H34B	0.9241
Ni1—O11	2.121 (2)	S31—C32	1.711 (4)
Ni1—O21	2.164 (2)	S31—C33	1.736 (3)
Ni2—O4	2.041 (2)	S32—C35	1.720 (3)
Ni2—O1	2.056 (2)	S32—C36	1.740 (3)
Ni2—N31	2.067 (2)	C11—C12	1.375 (4)
Ni2—O3	2.0695 (19)	C11—C16	1.519 (4)
Ni2—N33	2.096 (2)	C12—C13	1.380 (4)
Ni2—O2	2.131 (2)	C12—H12	0.9300
O1—H1A	0.8609	C13—C14	1.390 (4)
O1—H1B	0.8248	C13—H13	0.9300
O2—H2A	0.7732	C14—C15	1.371 (4)
O2—H2B	0.7640	C14—H14	0.9300
O3—H3A	0.7774	C15—C17	1.523 (4)
O3—H3B	0.7478	C21—C22	1.381 (4)
O4—H4A	0.8275	C21—C26	1.515 (4)
O4—H4B	0.7402	C22—C23	1.386 (4)
O11—C16	1.267 (4)	C22—H22	0.9300
O12—C16	1.236 (3)	C23—C24	1.383 (4)
O13—C17	1.257 (3)	C23—H23	0.9300
O14—C17	1.235 (3)	C24—C25	1.373 (4)
O21—C26	1.276 (3)	C24—H24	0.9300
O22—C26	1.234 (3)	C25—C27	1.510 (4)
O23—C27	1.263 (4)	C31—C32	1.344 (4)
O24—C27	1.247 (4)	C31—C34	1.453 (4)
N11—C11	1.332 (4)	C32—H32	0.9300
N11—C15	1.337 (3)	C34—C35	1.340 (4)
N21—C21	1.330 (3)	C35—H35	0.9300
N21—C25	1.333 (4)	O1W—H1WA	0.9140
N31—C33	1.314 (4)	O1W—H1WB	0.7473
N31—C31	1.387 (4)	O2W—H2WB	0.7194
N32—C33	1.335 (4)	O2W—H2WA	0.7934
N32—H32A	0.9783	O3W—H3WA	0.7332
N32—H32B	0.9467	O3W—H3WB	0.8837
N33—C36	1.312 (4)		
N11—Ni1—N21	175.99 (9)	C35—S32—C36	89.37 (15)
N11—Ni1—O13	79.07 (9)	N11—C11—C12	120.2 (3)
N21—Ni1—O13	99.01 (9)	N11—C11—C16	112.4 (2)
N11—Ni1—O23	97.79 (9)	C12—C11—C16	127.4 (3)
N21—Ni1—O23	78.84 (9)	C11—C12—C13	119.0 (3)
O13—Ni1—O23	95.95 (9)	C11—C12—H12	120.5
N11—Ni1—O11	77.57 (8)	C13—C12—H12	120.5
N21—Ni1—O11	104.53 (9)	C12—C13—C14	119.6 (3)
O13—Ni1—O11	156.33 (8)	C12—C13—H13	120.2
O23—Ni1—O11	90.94 (9)	C14—C13—H13	120.2
N11—Ni1—O21	106.47 (9)	C15—C14—C13	118.9 (3)
N21—Ni1—O21	77.05 (8)	C15—C14—H14	120.6
O13—Ni1—O21	92.33 (8)	C13—C14—H14	120.6

O23—Ni1—O21	155.44 (8)	N11—C15—C14	120.2 (3)
O11—Ni1—O21	90.65 (8)	N11—C15—C17	111.9 (2)
O4—Ni2—O1	170.27 (9)	C14—C15—C17	127.9 (3)
O4—Ni2—N31	95.55 (9)	O12—C16—O11	126.0 (3)
O1—Ni2—N31	88.13 (9)	O12—C16—C11	119.3 (3)
O4—Ni2—O3	86.23 (8)	O11—C16—C11	114.7 (2)
O1—Ni2—O3	91.32 (8)	O14—C17—O13	125.6 (3)
N31—Ni2—O3	172.35 (9)	O14—C17—C15	119.2 (3)
O4—Ni2—N33	94.86 (9)	O13—C17—C15	115.2 (2)
O1—Ni2—N33	94.67 (9)	N21—C21—C22	120.8 (3)
N31—Ni2—N33	79.68 (9)	N21—C21—C26	113.1 (2)
O3—Ni2—N33	92.77 (9)	C22—C21—C26	126.1 (3)
O4—Ni2—O2	87.35 (9)	C21—C22—C23	118.2 (3)
O1—Ni2—O2	83.27 (8)	C21—C22—H22	120.9
N31—Ni2—O2	96.50 (9)	C23—C22—H22	120.9
O3—Ni2—O2	91.00 (8)	C24—C23—C22	120.1 (3)
N33—Ni2—O2	175.75 (8)	C24—C23—H23	119.9
Ni2—O1—H1A	116.2	C22—C23—H23	119.9
Ni2—O1—H1B	128.2	C25—C24—C23	118.5 (3)
H1A—O1—H1B	111.8	C25—C24—H24	120.7
Ni2—O2—H2A	119.6	C23—C24—H24	120.7
Ni2—O2—H2B	112.6	N21—C25—C24	120.9 (3)
H2A—O2—H2B	100.9	N21—C25—C27	112.8 (3)
Ni2—O3—H3A	115.4	C24—C25—C27	126.3 (3)
Ni2—O3—H3B	109.9	O22—C26—O21	126.0 (3)
H3A—O3—H3B	105.8	O22—C26—C21	119.1 (3)
Ni2—O4—H4A	132.5	O21—C26—C21	114.8 (2)
Ni2—O4—H4B	118.2	O24—C27—O23	125.7 (3)
H4A—O4—H4B	106.3	O24—C27—C25	118.3 (3)
C16—O11—Ni1	115.03 (18)	O23—C27—C25	116.1 (3)
C17—O13—Ni1	115.52 (18)	C32—C31—N31	115.6 (3)
C26—O21—Ni1	114.41 (18)	C32—C31—C34	128.2 (3)
C27—O23—Ni1	113.85 (19)	N31—C31—C34	116.2 (2)
C11—N11—C15	122.1 (2)	C31—C32—S31	110.3 (3)
C11—N11—Ni1	119.51 (19)	C31—C32—H32	124.8
C15—N11—Ni1	117.94 (19)	S31—C32—H32	124.8
C21—N21—C25	121.5 (2)	N31—C33—N32	124.3 (3)
C21—N21—Ni1	120.25 (19)	N31—C33—S31	113.6 (3)
C25—N21—Ni1	117.90 (19)	N32—C33—S31	122.0 (3)
C33—N31—C31	110.7 (3)	C35—C34—N33	115.4 (3)
C33—N31—Ni2	134.9 (2)	C35—C34—C31	129.0 (3)
C31—N31—Ni2	114.33 (19)	N33—C34—C31	115.5 (3)
C33—N32—H32A	114.5	C34—C35—S32	110.6 (2)
C33—N32—H32B	119.9	C34—C35—H35	124.7
H32A—N32—H32B	121.6	S32—C35—H35	124.7
C36—N33—C34	110.7 (3)	N33—C36—N34	125.1 (3)
C36—N33—Ni2	135.7 (2)	N33—C36—S32	113.9 (2)
C34—N33—Ni2	113.26 (19)	N34—C36—S32	121.0 (2)
C36—N34—H34A	111.4	H1WA—O1W—H1WB	107.0

supplementary materials

C36—N34—H34B	116.1	H2WB—O2W—H2WA	104.5
H34A—N34—H34B	126.2	H3WA—O3W—H3WB	107.6
C32—S31—C33	89.77 (16)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1A \cdots O14 ⁱ	0.86	1.75	2.606 (3)	174
O1—H1B \cdots O21 ⁱⁱ	0.82	1.92	2.747 (3)	176
O2—H2A \cdots O12 ⁱⁱⁱ	0.77	2.01	2.772 (3)	169
O2—H2B \cdots O22 ⁱⁱ	0.76	2.12	2.874 (3)	170
O3—H3A \cdots O3W ⁱⁱ	0.78	1.98	2.742 (3)	169
O3—H3B \cdots O11 ⁱⁱⁱ	0.75	1.94	2.683 (3)	173
O4—H4A \cdots O24 ^{iv}	0.83	1.89	2.717 (4)	179
O4—H4B \cdots O1W	0.74	2.16	2.820 (4)	149
O1W—H1WA \cdots O23 ^{iv}	0.91	2.01	2.907 (4)	166
O1W—H1WB \cdots O24 ⁱⁱⁱ	0.75	2.41	3.150 (5)	169
O2W—H2WB \cdots O3W ^v	0.72	2.41	3.104 (4)	162
O2W—H2WA \cdots O14	0.79	2.02	2.815 (4)	174
O3W—H3WA \cdots O22 ^{vi}	0.73	2.24	2.934 (3)	158
O3W—H3WB \cdots O21	0.88	2.10	2.892 (3)	149
N32—H32A \cdots O12	0.98	1.97	2.940 (5)	169
N32—H32B \cdots O2	0.95	2.19	2.996 (5)	142
N34—H34A \cdots O2W ^{vii}	0.92	1.98	2.886 (4)	168
N34—H34B \cdots O3	0.92	2.12	2.952 (4)	149
C12—H12 \cdots S31	0.93	2.71	3.631 (4)	170

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x, y, z+1$; (iii) $-x, -y+1, -z+1$; (iv) $x, y+1, z$; (v) $-x+1, -y, -z$; (vi) $-x, -y+1, -z$; (vii) $x, y+1, z+1$.

Fig. 1

