

Tris(3,5-dimethyl-1*H*-pyrazole- κN^2)- (pyridine-2,6-dicarboxylato- $\kappa^3 O^2, N, O^6$)- cobalt(II) monohydrate

Kun-Hua Lin,^a Zhe-Yin Yu,^{a*} Yan-Hua Zhong^a and Min Shao^b

^aDepartment of Chemistry, College of Science, Shanghai University, Shanghai 200444, People's Republic of China, and ^bInstrumental Analysis and Research Center, Shanghai University, Shanghai 200444, People's Republic of China
Correspondence e-mail: yuzheyin@shu.edu.cn


Received 16 February 2009; accepted 23 March 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.030; wR factor = 0.077; data-to-parameter ratio = 12.6.

The reaction of $\text{Co}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ with pyridine-2,6-dicarboxylic acid and 3,5-dimethyl-1*H*-pyrazole in a 1:1:3 molar ratio affords the title complex, $[\text{Co}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_5\text{H}_8\text{N}_2)_3] \cdot \text{H}_2\text{O}$. The Co^{II} atom is coordinated by one pyridine-2,6-dicarboxylate chelating ligand and three 3,5-dimethyl-1*H*-pyrazole ligands in a distorted octahedral geometry. Hydrogen-bonding interactions involving the coordinated carboxylate group, 3,5-dimethyl-1*H*-pyrazole and water help to consolidate the crystal structure

Related literature

For the use of complexes with pyrazole-based ligands in studying the relationship between the structure and the activity of the active site of metalloproteins, see: Haanstra *et al.* (1990). For the coordination modes of pyrazole complexes, see: Grotjahn *et al.* (2003).



Experimental

Crystal data

$[\text{Co}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_5\text{H}_8\text{N}_2)_3] \cdot \text{H}_2\text{O}$
 $M_r = 530.45$

Triclinic, $P\bar{1}$
 $a = 8.4220(8)\text{ \AA}$

$b = 11.9936(12)\text{ \AA}$	$Z = 2$
$c = 13.1418(13)\text{ \AA}$	$\text{Mo } K\alpha$ radiation
$\alpha = 75.1290(10)^\circ$	$\mu = 0.76\text{ mm}^{-1}$
$\beta = 84.7720(10)^\circ$	$T = 296\text{ K}$
$\gamma = 70.0940(10)^\circ$	$0.30 \times 0.30 \times 0.25\text{ mm}$
$V = 1206.3(2)\text{ \AA}^3$	

Data collection

Bruker SMART CCD area-detector diffractometer	6298 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4183 independent reflections
$T_{\min} = 0.804$, $T_{\max} = 0.833$	3769 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.014$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.077$	$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$
4183 reflections	
331 parameters	
2 restraints	

Table 1
Selected geometric parameters (\AA , $^\circ$).

Co1—N1	2.0407 (16)	Co1—O3	2.1522 (14)
Co1—N4	2.0798 (16)	Co1—N2	2.2336 (17)
Co1—O1	2.1453 (14)	Co1—N6	2.2477 (17)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3A \cdots O1W ⁱ	0.86	2.22	2.926 (2)	139
N3—H3A \cdots O1	0.86	2.61	3.048 (2)	113
N5—H5 \cdots O2 ⁱⁱ	0.86	2.10	2.945 (2)	168
N7—H7 \cdots O2 ⁱⁱ	0.86	2.08	2.838 (2)	146
N7—H7 \cdots O3	0.86	2.42	2.906 (2)	116
O1W—H1WA \cdots O4 ⁱⁱⁱ	0.844 (17)	1.967 (18)	2.797 (2)	168 (3)
O1W—H1WB \cdots O3 ^{iv}	0.828 (17)	2.204 (19)	3.009 (2)	164 (3)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$; (iii) $x, y + 1, z$; (iv) $-x, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2127).

References

- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Grotjahn, D. B., Van, S., Combs, D., Daniel, A., Schneider, C., Incarvito, C. D., Lam, K.-C., Rossi, G., Rheingold, A. L., Rideout, M., Meyer, C., Hernandez, G. & Mejorado, L. (2003). *Inorg. Chem.* **42**, 3347–3355.
- Haanstra, W. G., Van der Donk, W. A. J. W., Driessen, W. L., Reedijk, J., Wood, J. S. & Drew, M. G. B. (1990). *J. Chem. Soc. Dalton Trans.* **10**, 3123–3128.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2009). E65, m449 [doi:10.1107/S1600536809010538]

Tris(3,5-dimethyl-1*H*-pyrazole- κN^2)(pyridine-2,6-dicarboxylato- $\kappa^3 O^2,N,O^6$)cobalt(II) monohydrate

Kun-Hua Lin, Zhe-Yin Yu, Yan-Hua Zhong and Min Shao

S1. Comment

Complexes with pyrazole-based ligands are a frequent subject of chemical investigations giving an opportunity for a better understanding of the relationship between the structure and the activity of the active site of metalloproteins (Haanstra *et al.*, 1990). Nowadays, attention is paid to the design of various pyrazole ligands, and some coordination modes of pyrazole complexes were reported (Grotjahn *et al.*, 2003). In our systematic studies on transition metal complexes with the pyrazole derivatives, the title compound was prepared and its X-ray structure is presented here.


The molecular structure of the title compound is shown in Fig. 1. The compound assumes a distorted octahedron geometry, formed by three 3,5-Dimethyl-1-*H*-pyrazole molecules and a pyridine-2,6-dicarboxylate. Tridentate ligand pyridine-2,6-dicarboxylate dianion chelates to the Co atom by a N atom of pyridine ring and two O atoms of carboxyl groups with a meridional configuration. Monodentate ligand 3,5-Dimethyl-1-*H*-pyrazole coordinated to the Co atom by N atoms of pyrazole rings. The bond distances of Co1—N1 and Co1—N4 are 2.0407 (16) Å and 2.0798 (16) Å (Table 1), which are shorter than the the bond distances of Co1—N2 and Co1—N6 with 2.2336 (17) Å and 2.2477 (17) Å.

S2. Experimental


An ethanol solution (6 ml) containing 3,5-Dimethyl-1-*H*-pyrazole(0.1153 g, 1.2 mmol) and Co(NO₃)₂.3H₂O(0.0870 g, 0.3 mmol) was mixed with an aqueous solution (6 ml) of pyridine-2,6-dicarboxylic acid(0.0501 g, 0.3 mmol) and NaOH (0.0240 g, 0.6 mmol). The mixture was refluxed for 6 h. The solution was filtered after cooling to room temperature. Pink single crystals suitable for X-ray diffraction were obtained from the filtrate after 11 d.

S3. Refinement

The H atoms of water molecule were located in a difference Fourier map and refined freely. Methyl H atoms were placed in caculated positions with C—H distances = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. Other H atoms were placed in caculated positions with C—H distances = 0.93 Å and N—H distances = 0.86 Å, and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C},\text{N})$.

**Figure 1**

Molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

A packing diagram of (I).

Tris(3,5-dimethyl-1*H*-pyrazole- κ^2 N²)(pyridine-2,6-dicarboxylato- κ^3 O²,N,O⁶)cobalt(II) monohydrate*Crystal data*

$[Co(C_7H_3NO_4)(C_5H_8N_2)_3] \cdot H_2O$	$Z = 2$
$M_r = 530.45$	$F(000) = 554$
Triclinic, $P\bar{1}$	$D_x = 1.460 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 8.4220 (8) \text{ \AA}$	Cell parameters from 4060 reflections
$b = 11.9936 (12) \text{ \AA}$	$\theta = 2.7\text{--}27.5^\circ$
$c = 13.1418 (13) \text{ \AA}$	$\mu = 0.76 \text{ mm}^{-1}$
$\alpha = 75.129 (1)^\circ$	$T = 296 \text{ K}$
$\beta = 84.772 (1)^\circ$	Block, pink
$\gamma = 70.094 (1)^\circ$	$0.30 \times 0.30 \times 0.25 \text{ mm}$
$V = 1206.3 (2) \text{ \AA}^3$	

Data collection

Bruker SMART CCD area-detector diffractometer	6298 measured reflections
Radiation source: fine-focus sealed tube	4183 independent reflections
Graphite monochromator	3769 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.014$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\max} = 25.1^\circ, \theta_{\min} = 2.1^\circ$
$T_{\min} = 0.804, T_{\max} = 0.833$	$h = -10 \rightarrow 9$
	$k = -14 \rightarrow 14$
	$l = -13 \rightarrow 15$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.030$	$w = 1/[\sigma^2(F_o^2) + (0.0334P)^2 + 0.7243P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.077$	$(\Delta/\sigma)_{\max} = 0.002$
$S = 1.03$	$\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
4183 reflections	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
331 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
2 restraints	Extinction coefficient: 0.0176 (11)
Primary atom site location: structure-invariant direct methods	
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.33263 (3)	0.28250 (2)	0.246751 (19)	0.02818 (10)
C1	0.5981 (2)	0.24913 (18)	0.39330 (15)	0.0295 (4)
C2	0.6852 (3)	0.2085 (2)	0.48700 (17)	0.0392 (5)
H2	0.7859	0.2228	0.4914	0.047*
C3	0.6195 (3)	0.1461 (2)	0.57368 (17)	0.0457 (6)
H3	0.6770	0.1171	0.6373	0.055*
C4	0.4680 (3)	0.1260 (2)	0.56719 (17)	0.0398 (5)
H4	0.4220	0.0853	0.6259	0.048*
C5	0.3884 (3)	0.16841 (17)	0.47115 (15)	0.0308 (4)
C6	0.6491 (2)	0.31692 (18)	0.28856 (16)	0.0310 (4)
C7	0.2218 (3)	0.15601 (18)	0.44770 (17)	0.0339 (5)
C8	0.7310 (3)	-0.0243 (2)	0.15919 (17)	0.0369 (5)
C9	0.5995 (3)	-0.0532 (2)	0.13323 (17)	0.0400 (5)
H9	0.6068	-0.1179	0.1042	0.048*
C10	0.4521 (3)	0.03249 (19)	0.15849 (15)	0.0330 (4)
C11	0.2751 (3)	0.0397 (2)	0.14480 (18)	0.0418 (5)
H11A	0.2067	0.0675	0.2017	0.063*
H11B	0.2726	-0.0397	0.1446	0.063*
H11C	0.2318	0.0960	0.0792	0.063*
C12	0.9178 (3)	-0.0810 (2)	0.1495 (2)	0.0549 (7)
H12A	0.9611	-0.0264	0.0971	0.082*
H12B	0.9424	-0.1565	0.1292	0.082*
H12C	0.9699	-0.0965	0.2160	0.082*
C13	0.2400 (2)	0.34373 (19)	0.00957 (15)	0.0327 (4)
C14	0.1089 (3)	0.3533 (2)	-0.05258 (16)	0.0361 (5)
H14	0.1106	0.3631	-0.1252	0.043*
C15	-0.0231 (3)	0.34544 (19)	0.01417 (16)	0.0340 (5)
C16	-0.1946 (3)	0.3455 (3)	-0.0043 (2)	0.0491 (6)
H16A	-0.1863	0.2693	-0.0190	0.074*
H16B	-0.2450	0.4114	-0.0632	0.074*
H16C	-0.2632	0.3558	0.0572	0.074*
C17	0.4138 (3)	0.3448 (3)	-0.02341 (18)	0.0479 (6)
H17A	0.4612	0.3677	0.0286	0.072*
H17B	0.4089	0.4027	-0.0899	0.072*
H17C	0.4832	0.2649	-0.0302	0.072*
C18	-0.0812 (3)	0.5490 (2)	0.36524 (17)	0.0381 (5)
C19	0.0209 (3)	0.6189 (2)	0.35379 (19)	0.0439 (5)
H19	-0.0057	0.6941	0.3704	0.053*
C20	0.1734 (3)	0.5549 (2)	0.31215 (17)	0.0375 (5)
C21	0.3282 (3)	0.5905 (2)	0.2884 (2)	0.0546 (6)
H21A	0.3927	0.5666	0.3516	0.082*
H21B	0.2966	0.6774	0.2615	0.082*
H21C	0.3950	0.5504	0.2368	0.082*
C22	-0.2576 (3)	0.5668 (3)	0.4048 (2)	0.0580 (7)
H22A	-0.3351	0.6281	0.3534	0.087*

H22B	-0.2709	0.5926	0.4694	0.087*
H22C	-0.2804	0.4913	0.4172	0.087*
N1	0.45453 (19)	0.22688 (14)	0.38745 (12)	0.0272 (3)
N2	0.4898 (2)	0.11200 (15)	0.19911 (13)	0.0322 (4)
N3	0.6621 (2)	0.07452 (16)	0.19804 (14)	0.0342 (4)
H3A	0.7201	0.1105	0.2200	0.041*
N4	0.1920 (2)	0.33111 (15)	0.11058 (13)	0.0309 (4)
N5	0.0297 (2)	0.33282 (16)	0.11091 (13)	0.0316 (4)
H5	-0.0318	0.3265	0.1666	0.038*
N6	0.1671 (2)	0.45011 (15)	0.29795 (13)	0.0330 (4)
N7	0.0095 (2)	0.44975 (16)	0.33138 (13)	0.0343 (4)
H7	-0.0280	0.3916	0.3308	0.041*
O1	0.55090 (18)	0.33974 (14)	0.21281 (11)	0.0362 (3)
O2	0.77947 (18)	0.34440 (15)	0.28302 (12)	0.0418 (4)
O3	0.18031 (17)	0.18992 (13)	0.35129 (11)	0.0353 (3)
O4	0.1395 (2)	0.11702 (16)	0.52179 (13)	0.0552 (5)
O1W	0.0578 (2)	0.93763 (17)	0.67852 (14)	0.0487 (4)
H1WA	0.073 (4)	0.989 (2)	0.6251 (17)	0.063 (9)*
H1WB	0.006 (4)	0.898 (3)	0.661 (2)	0.071 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02400 (16)	0.03763 (17)	0.02563 (15)	-0.01357 (12)	-0.00029 (10)	-0.00758 (11)
C1	0.0247 (10)	0.0335 (11)	0.0329 (10)	-0.0085 (8)	-0.0003 (8)	-0.0137 (8)
C2	0.0330 (12)	0.0478 (13)	0.0395 (12)	-0.0118 (10)	-0.0058 (9)	-0.0153 (10)
C3	0.0481 (14)	0.0498 (14)	0.0333 (12)	-0.0075 (11)	-0.0119 (10)	-0.0076 (10)
C4	0.0483 (13)	0.0369 (12)	0.0299 (11)	-0.0115 (10)	0.0013 (9)	-0.0046 (9)
C5	0.0355 (11)	0.0271 (10)	0.0296 (10)	-0.0097 (8)	0.0037 (8)	-0.0086 (8)
C6	0.0268 (11)	0.0354 (11)	0.0362 (11)	-0.0132 (9)	0.0039 (8)	-0.0153 (9)
C7	0.0353 (11)	0.0289 (10)	0.0399 (12)	-0.0137 (9)	0.0074 (9)	-0.0110 (9)
C8	0.0319 (11)	0.0424 (12)	0.0374 (11)	-0.0109 (9)	0.0036 (9)	-0.0145 (9)
C9	0.0385 (12)	0.0447 (13)	0.0433 (12)	-0.0144 (10)	0.0036 (10)	-0.0225 (10)
C10	0.0330 (11)	0.0400 (11)	0.0289 (10)	-0.0154 (9)	0.0009 (8)	-0.0092 (9)
C11	0.0355 (12)	0.0514 (14)	0.0454 (13)	-0.0200 (10)	0.0013 (10)	-0.0161 (11)
C12	0.0348 (13)	0.0628 (16)	0.0710 (17)	-0.0104 (12)	0.0068 (12)	-0.0330 (14)
C13	0.0281 (11)	0.0414 (12)	0.0303 (10)	-0.0142 (9)	0.0014 (8)	-0.0083 (9)
C14	0.0319 (11)	0.0495 (13)	0.0277 (10)	-0.0148 (10)	-0.0011 (8)	-0.0086 (9)
C15	0.0268 (11)	0.0412 (12)	0.0344 (11)	-0.0106 (9)	-0.0042 (8)	-0.0090 (9)
C16	0.0282 (12)	0.0743 (17)	0.0508 (14)	-0.0197 (12)	-0.0037 (10)	-0.0206 (12)
C17	0.0347 (12)	0.0744 (17)	0.0414 (13)	-0.0274 (12)	0.0070 (10)	-0.0151 (12)
C18	0.0337 (11)	0.0404 (12)	0.0400 (12)	-0.0073 (9)	-0.0020 (9)	-0.0155 (10)
C19	0.0463 (14)	0.0381 (12)	0.0518 (14)	-0.0129 (10)	0.0001 (11)	-0.0203 (10)
C20	0.0400 (12)	0.0408 (12)	0.0369 (11)	-0.0180 (10)	-0.0015 (9)	-0.0113 (9)
C21	0.0520 (15)	0.0570 (16)	0.0689 (17)	-0.0317 (13)	0.0074 (13)	-0.0232 (13)
C22	0.0368 (13)	0.0637 (17)	0.0790 (19)	-0.0108 (12)	0.0106 (13)	-0.0379 (15)
N1	0.0257 (8)	0.0297 (8)	0.0276 (8)	-0.0098 (7)	0.0012 (7)	-0.0090 (7)
N2	0.0254 (9)	0.0382 (9)	0.0344 (9)	-0.0105 (7)	0.0008 (7)	-0.0113 (7)

N3	0.0262 (9)	0.0398 (10)	0.0410 (10)	-0.0130 (8)	0.0003 (7)	-0.0147 (8)
N4	0.0228 (8)	0.0404 (10)	0.0309 (9)	-0.0122 (7)	-0.0008 (7)	-0.0083 (7)
N5	0.0230 (8)	0.0447 (10)	0.0288 (9)	-0.0138 (7)	0.0028 (7)	-0.0090 (7)
N6	0.0285 (9)	0.0383 (10)	0.0350 (9)	-0.0130 (7)	0.0016 (7)	-0.0117 (7)
N7	0.0284 (9)	0.0369 (10)	0.0430 (10)	-0.0132 (8)	0.0024 (7)	-0.0167 (8)
O1	0.0334 (8)	0.0495 (9)	0.0309 (7)	-0.0221 (7)	0.0011 (6)	-0.0073 (6)
O2	0.0316 (8)	0.0559 (10)	0.0486 (9)	-0.0259 (7)	0.0041 (7)	-0.0168 (7)
O3	0.0315 (8)	0.0417 (8)	0.0382 (8)	-0.0186 (6)	0.0023 (6)	-0.0108 (6)
O4	0.0571 (11)	0.0660 (11)	0.0486 (10)	-0.0381 (9)	0.0155 (8)	-0.0056 (8)
O1W	0.0543 (11)	0.0579 (11)	0.0447 (10)	-0.0330 (9)	0.0035 (8)	-0.0123 (9)

Geometric parameters (\AA , $^{\circ}$)

Co1—N1	2.0407 (16)	C13—N4	1.337 (3)
Co1—N4	2.0798 (16)	C13—C14	1.389 (3)
Co1—O1	2.1453 (14)	C13—C17	1.492 (3)
Co1—O3	2.1522 (14)	C14—C15	1.366 (3)
Co1—N2	2.2336 (17)	C14—H14	0.9300
Co1—N6	2.2477 (17)	C15—N5	1.340 (3)
C1—N1	1.337 (2)	C15—C16	1.486 (3)
C1—C2	1.380 (3)	C16—H16A	0.9600
C1—C6	1.515 (3)	C16—H16B	0.9600
C2—C3	1.376 (3)	C16—H16C	0.9600
C2—H2	0.9300	C17—H17A	0.9600
C3—C4	1.390 (3)	C17—H17B	0.9600
C3—H3	0.9300	C17—H17C	0.9600
C4—C5	1.376 (3)	C18—N7	1.337 (3)
C4—H4	0.9300	C18—C19	1.367 (3)
C5—N1	1.331 (2)	C18—C22	1.491 (3)
C5—C7	1.526 (3)	C19—C20	1.398 (3)
C6—O2	1.241 (2)	C19—H19	0.9300
C6—O1	1.267 (2)	C20—N6	1.336 (3)
C7—O4	1.229 (2)	C20—C21	1.489 (3)
C7—O3	1.270 (2)	C21—H21A	0.9600
C8—N3	1.338 (3)	C21—H21B	0.9600
C8—C9	1.362 (3)	C21—H21C	0.9600
C8—C12	1.494 (3)	C22—H22A	0.9600
C9—C10	1.393 (3)	C22—H22B	0.9600
C9—H9	0.9300	C22—H22C	0.9600
C10—N2	1.339 (3)	N2—N3	1.365 (2)
C10—C11	1.488 (3)	N3—H3A	0.8600
C11—H11A	0.9600	N4—N5	1.359 (2)
C11—H11B	0.9600	N5—H5	0.8600
C11—H11C	0.9600	N6—N7	1.361 (2)
C12—H12A	0.9600	N7—H7	0.8600
C12—H12B	0.9600	O1W—H1WA	0.844 (17)
C12—H12C	0.9600	O1W—H1WB	0.828 (17)

N1—Co1—N4	174.52 (6)	C13—C14—H14	126.9
N1—Co1—O1	75.49 (6)	N5—C15—C14	106.49 (17)
N4—Co1—O1	109.98 (6)	N5—C15—C16	121.66 (19)
N1—Co1—O3	76.60 (6)	C14—C15—C16	131.8 (2)
N4—Co1—O3	97.92 (6)	C15—C16—H16A	109.5
O1—Co1—O3	152.00 (5)	C15—C16—H16B	109.5
N1—Co1—N2	91.87 (6)	H16A—C16—H16B	109.5
N4—Co1—N2	88.21 (6)	C15—C16—H16C	109.5
O1—Co1—N2	86.36 (6)	H16A—C16—H16C	109.5
O3—Co1—N2	92.37 (6)	H16B—C16—H16C	109.5
N1—Co1—N6	88.03 (6)	C13—C17—H17A	109.5
N4—Co1—N6	91.67 (6)	C13—C17—H17B	109.5
O1—Co1—N6	95.87 (6)	H17A—C17—H17B	109.5
O3—Co1—N6	85.33 (6)	C13—C17—H17C	109.5
N2—Co1—N6	177.66 (6)	H17A—C17—H17C	109.5
N1—C1—C2	120.37 (19)	H17B—C17—H17C	109.5
N1—C1—C6	112.56 (16)	N7—C18—C19	105.87 (19)
C2—C1—C6	127.04 (18)	N7—C18—C22	121.8 (2)
C3—C2—C1	118.3 (2)	C19—C18—C22	132.4 (2)
C3—C2—H2	120.8	C18—C19—C20	106.28 (19)
C1—C2—H2	120.8	C18—C19—H19	126.9
C2—C3—C4	120.7 (2)	C20—C19—H19	126.9
C2—C3—H3	119.6	N6—C20—C19	110.63 (19)
C4—C3—H3	119.6	N6—C20—C21	122.0 (2)
C5—C4—C3	117.9 (2)	C19—C20—C21	127.3 (2)
C5—C4—H4	121.1	C20—C21—H21A	109.5
C3—C4—H4	121.1	C20—C21—H21B	109.5
N1—C5—C4	120.88 (19)	H21A—C21—H21B	109.5
N1—C5—C7	113.15 (17)	C20—C21—H21C	109.5
C4—C5—C7	125.96 (19)	H21A—C21—H21C	109.5
O2—C6—O1	125.78 (19)	H21B—C21—H21C	109.5
O2—C6—C1	119.27 (18)	C18—C22—H22A	109.5
O1—C6—C1	114.94 (16)	C18—C22—H22B	109.5
O4—C7—O3	126.1 (2)	H22A—C22—H22B	109.5
O4—C7—C5	118.49 (19)	C18—C22—H22C	109.5
O3—C7—C5	115.42 (17)	H22A—C22—H22C	109.5
N3—C8—C9	106.06 (18)	H22B—C22—H22C	109.5
N3—C8—C12	122.1 (2)	C5—N1—C1	121.79 (17)
C9—C8—C12	131.8 (2)	C5—N1—Co1	118.58 (13)
C8—C9—C10	106.82 (19)	C1—N1—Co1	119.63 (13)
C8—C9—H9	126.6	C10—N2—N3	104.33 (16)
C10—C9—H9	126.6	C10—N2—Co1	133.05 (14)
N2—C10—C9	110.16 (18)	N3—N2—Co1	122.35 (12)
N2—C10—C11	122.52 (19)	C8—N3—N2	112.62 (16)
C9—C10—C11	127.32 (19)	C8—N3—H3A	123.7
C10—C11—H11A	109.5	N2—N3—H3A	123.7
C10—C11—H11B	109.5	C13—N4—N5	104.78 (15)
H11A—C11—H11B	109.5	C13—N4—Co1	130.69 (13)

C10—C11—H11C	109.5	N5—N4—Co1	123.41 (12)
H11A—C11—H11C	109.5	C15—N5—N4	112.05 (16)
H11B—C11—H11C	109.5	C15—N5—H5	124.0
C8—C12—H12A	109.5	N4—N5—H5	124.0
C8—C12—H12B	109.5	C20—N6—N7	104.03 (16)
H12A—C12—H12B	109.5	C20—N6—Co1	140.31 (14)
C8—C12—H12C	109.5	N7—N6—Co1	115.57 (12)
H12A—C12—H12C	109.5	C18—N7—N6	113.19 (17)
H12B—C12—H12C	109.5	C18—N7—H7	123.4
N4—C13—C14	110.39 (17)	N6—N7—H7	123.4
N4—C13—C17	121.13 (18)	C6—O1—Co1	117.38 (12)
C14—C13—C17	128.48 (19)	C7—O3—Co1	115.59 (12)
C15—C14—C13	106.29 (18)	H1WA—O1W—H1WB	110 (3)
C15—C14—H14	126.9		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3A···O1W ⁱ	0.86	2.22	2.926 (2)	139
N3—H3A···O1	0.86	2.61	3.048 (2)	113
N5—H5···O2 ⁱⁱ	0.86	2.10	2.945 (2)	168
N7—H7···O2 ⁱⁱ	0.86	2.08	2.838 (2)	146
N7—H7···O3	0.86	2.42	2.906 (2)	116
O1W—H1WA···O4 ⁱⁱⁱ	0.84 (2)	1.97 (2)	2.797 (2)	168 (3)
O1W—H1WB···O3 ^{iv}	0.83 (2)	2.20 (2)	3.009 (2)	164 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x-1, y, z$; (iii) $x, y+1, z$; (iv) $-x, -y+1, -z+1$.