

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

# Diaqua[*N,N'*-bis(3-carboxyprop-2-enoyl)pyridine-2,6-dicarbohydrazidato(2-)]cadmium(II) *N,N*-dimethylformamide disolvate

Quanfu Cao and Dacheng Li\*

College of Chemistry and Chemical Engineering, Liaocheng University, Liaocheng 252059, People's Republic of China

Correspondence e-mail: lidacheng62@lcu.edu.cn


Received 18 February 2009; accepted 25 March 2009

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.027;  $wR$  factor = 0.075; data-to-parameter ratio = 13.0.

In the title complex,  $[\text{Cd}(\text{C}_{15}\text{H}_{11}\text{N}_5\text{O}_8)(\text{H}_2\text{O})_2] \cdot 2\text{C}_3\text{H}_7\text{NO}$ , the  $\text{Cd}^{\text{II}}$  ion is located on a twofold rotation axis and is seven-coordinated in a distorted pentagonal-bipyramidal manner. The asymmetric unit comprises one metal ion, one doubly deprotonated *N,N'*-bis(3-carboxyprop-2-enoyl)pyridine-2,6-dicarbohydrazide ligand, two coordinating water molecules and two dimethylformamide solvent molecules. In the crystal, a two-dimensional network is formed through  $\text{N}-\text{H} \cdots \text{O}$  and  $\text{O}-\text{H} \cdots \text{O}$  hydrogen bonds.

## Related literature

For polydimensional supermolecular architectures formed by aromatic hydrazides through hydrogen bonds and  $\pi-\pi$  interactions, see: Bacchi *et al.* (1993); Bermejo *et al.* (1999). The condensation products of 2,6-picolylhydrazide with anhydrides have been found to adopt a pentagonal-bipyramidal coordination in various metal complexes, see: Pelizzi *et al.* (1987); Wang *et al.* (2005). For the chelating behaviour of *N,N'*-acetyl-2,6-picolylhydrazide with  $\text{Fe}^{3+}$ , see: Cao *et al.* (2008). For our continuing study of aroylhydrazides, see: Dou *et al.* (2006). For  $\text{Cd}-\text{O}(\text{carbonyl})$  bond lengths in other seven-coordinated pentagonal-bipyramidal cadmium complexes, see: Charles *et al.* (1983).



## Experimental

### Crystal data

$[\text{Cd}(\text{C}_{15}\text{H}_{11}\text{N}_5\text{O}_8)(\text{H}_2\text{O})_2] \cdot 2\text{C}_3\text{H}_7\text{NO}$   
 $M_r = 683.91$   
 Monoclinic,  $C2/c$   
 $a = 18.6176$  (2) Å  
 $b = 12.6065$  (8) Å  
 $c = 12.0038$  (6) Å  
 $\beta = 99.51^\circ$   
 $V = 2778.6$  (2) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.86$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.20 \times 0.18 \times 0.17$  mm

### Data collection

Siemens SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\text{min}} = 0.847$ ,  $T_{\text{max}} = 0.868$   
 6846 measured reflections  
 2448 independent reflections  
 2071 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$   
 $wR(F^2) = 0.075$   
 $S = 1.00$   
 2448 reflections  
 189 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.73$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.48$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cd1—N2 <sup>i</sup>	2.287 (2)	Cd1—O2 <sup>i</sup>	2.444 (19)
Cd1—O5 <sup>i</sup>	2.3412 (19)	N2—N3	1.369 (3)
Cd1—N1	2.387 (3)		

Symmetry code: (i)  $-x + 1, y, -z + \frac{1}{2}$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O5—H5A $\cdots$ O4 <sup>ii</sup>	0.85	1.97	2.802 (3)	165
O5—H5B $\cdots$ O1 <sup>iii</sup>	0.85	1.84	2.685 (3)	174
N3—H3A $\cdots$ O6	0.86	1.97	2.808 (3)	163
O3—H3 $\cdots$ O2	0.82	1.68	2.498 (3)	175

Symmetry codes: (ii)  $-x + 1, -y, -z + 1$ ; (iii)  $-x + 1, -y + 1, -z + 1$ .

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We gratefully acknowledge financial support by the Natural Science Foundation of China (grant No. 20671048).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2207).

## References

- Bacchi, A., Battaglia, L. P., Carcelli, M., Pelizzi, C., Pelizzi, G., Solinas, C. & Zoroddu, M. A. (1993). *J. Chem. Soc. Dalton Trans.* pp. 775–779.  
 Bermejo, M. R., Fondo, M., González, A. M., Hoyos, O. L., Sousa, A., McAuliffe, C. A., Hussain, W., Pritchard, R. & Novotorsev, V. M. (1999). *J. Chem. Soc. Dalton Trans.* pp. 2211–2218.  
 Cao, Q.-F., Dou, J.-M., Li, D.-C. & Wang, D.-Q. (2008). *Acta Cryst.* E64, m47.

- Charles, N. G., Griffith, E. A. H., Rodesiler, P. F. & Amma, E. L. (1983). *Inorg. Chem.* **22**, 2717–2723.
- Dou, J. M., Liu, M. L., Li, D. C. & Wang, D. Q. (2006). *Eur. J. Inorg. Chem.* pp. 4866–4871.
- Pelizzi, C., Pelizzi, G. & Vitali, F. (1987). *J. Chem. Soc. Dalton Trans.* pp. 177–181.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wang, C. X., Du, C. X., Li, Y. H. & Wu, Y. J. (2005). *Inorg. Chem. Commun.* **8**, 379–381.

**supplementary materials**

*Acta Cryst.* (2009). E65, m464-m465 [ doi:10.1107/S1600536809011003 ]

**Diaqua[*N,N'*-bis(3-carboxyprop-2-enoyl)pyridine-2,6-dicarbohydrazidato(2-)]cadmium(II) *N,N'*-dimethylformamide disolvate**

**Q. Cao and D. Li**

**Comment**

Containing N, O and other coordinating sites, aromatic hydrazides can form poly-dimensional supermolecular architectures through hydrogen bonds and  $\pi$ - $\pi$  interactions (Bacchi *et al.*, 1993, Bermejo *et al.*, 1999). The condensation products of 2,6-picolylhydrazide with anhydrides have been found to adopt a pentagonal-bipyramidal stereochemistry in various metal complexes, in which they may participate as neutral and/or dianionic ligands (Pelizzi, *et al.*, 1987, Wang *et al.*, 2005). Previously we have examined the chelating behaviour of *N,N'*-acetyl-2,6-picolylhydrazide with  $\text{Fe}^{3+}$  (Cao, *et al.*, 2008). As a part of continuing study of our research on aroylhydrazide in our laboratory (Dou, *et al.*, 2006), we synthesized *N,N'*-bis(3-carboxy-*cis*-propenoyl)-2,6-picolylidihydrazide and obtained its Cd(II) complex (I).

The molecular structure of the complex (Fig. 1) and its characteristic geometry parameters (Table 1) reveal one cadmium ion which is located on the 2-fold rotation axis, one deprotonated ligand, two coordinated  $\text{H}_2\text{O}$  molecules and two solvent DMF molecules. The divalent anionic  $\text{H}_2\text{L}^{2-}$  acts as a pentadentate chelating ligand to two cadmium atoms. The remainder coordinating sites of  $\text{Cd}^{2+}$  are occupied by two O atoms from water molecules in *trans*-positions which complete the seven-coordinated pentagonal-bipyramid. Two deprotonated amide nitrogen atoms, two carbonyl O atoms, one pyridine N atom complete the equatorial plane and the mean deviation is 0.0064 Å indicating that the five atoms are ideally coplanar. Such planarity was observed in  $[\text{Cd}(\text{H}_2\text{daps})\text{Cl}_2](\text{CHCl}_3)(\text{CH}_3\text{OH})$  (less than 0.007 Å) ( $\text{H}_2\text{daps}$  = 2,6-diacetylpyridine bis(salicyloylhydrazone) (Pelizzi, *et al.*, 1987). The Cd—N distances are in the range of 2.287 (2) Å to 2.387 (3) Å; its average value of 2.320 (2) Å is shorter than those observed in  $[\text{Cd}(\text{L}')(\text{1.5H}_2\text{O})]_n$  ( $\text{L}' = \text{N,N'$ -bis(4-pyridylcarboxyl)-2,6-pyridine dicarbohydrazide) (Wang *et al.*, 2005) and  $[\text{Cd}(\text{H}_2\text{daps})\text{Cl}_2](\text{CHCl}_3)(\text{CH}_3\text{OH})$  (Pelizzi, *et al.*, 1987). Both, two Cd—O(carbonyl) bond lengths (2.4441 (19) Å) are comparable to those in other seven-coordinated pentagonal-bipyramidal cadmium complexes (Charles *et al.*, 1983). The Cd—O (water) distance is 2.341 (2) Å, being shorter than the mean lengths of Cd—O in the the equatorial plane of 2.444 (19) Å.

The crystal structure of the title complex is predominantly determined by N—H $\cdots$ O and O—H $\cdots$ O hydrogen bonds (Table 2 and Fig. 2) generating 2-D network.

**Experimental**

All chemicals were of reagent grade and were used without further purification. A solution of cadmium nitrate tetrahydrate (2 mmol, 0.457 g) dissolved in methanol (10 ml) was added dropwise to a DMF solution containing the ligand (2 mmol, 0.783 g). The mixture was stirred at room temperature for 6 h and then filtered. The filtrate was left to evaporate slowly at room temperature and yellow block-shaped crystals suitable for X-ray diffraction analysis were obtained after three weeks (m.p. >573 K). Elemental analysis calculated for (I): C: 36.88, H: 4.27, N: 14.34%; found: C: 36.11, H: 4.66, N: 14.02%. IR (KBr pellet,  $\text{cm}^{-1}$ ): 3467 (O—H), 3134 (N—H), 1709 (C=O) (acid carboxyl segment), 1647 (C=C).

## Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms, with pyridine C—H distances of 0.930 Å, hydrazide N—H distances of 0.860 Å, alkene C—H distances of 0.930 Å, methyl C—H distances of 0.960 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{O})$  and  $1.5 U_{\text{eq}}$  for methyl and hydroxy groups.

## Figures




Fig. 1. The molecular structure of the complex (I) showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes:  $-x + 1, y, -z + 3/2$ ]



Fig. 2. Part of the crystal structure of the complex, showing hydrogen bonds as dashed lines. [Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 1$ ].

## Diaqua[*N,N'*-bis(3-carboxyprop-2-enoyl)pyridine-2,6-dicarbohydrazidato(2-)]cadmium(II) *N,N*-dimethylformamide disolvate

### Crystal data

$[\text{Cd}(\text{C}_{15}\text{H}_{11}\text{N}_5\text{O}_8)(\text{H}_2\text{O})_2] \cdot 2\text{C}_3\text{H}_7\text{NO}$

$M_r = 683.91$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 18.6176 (2) \text{ \AA}$

$b = 12.6065 (8) \text{ \AA}$

$c = 12.0038 (6) \text{ \AA}$

$\beta = 99.51^\circ$

$V = 2778.6 (2) \text{ \AA}^3$

$Z = 4$

$F_{000} = 1392$

$D_x = 1.635 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3321 reflections

$\theta = 2.5\text{--}27.6^\circ$

$\mu = 0.86 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Block, yellow

$0.20 \times 0.18 \times 0.17 \text{ mm}$

### Data collection

Siemens SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298 \text{ K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

2448 independent reflections

2071 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\text{max}} = 25.0^\circ$

$\theta_{\text{min}} = 2.0^\circ$

$h = -22 \rightarrow 21$

(SADABS; Sheldrick, 1996)

$T_{\min} = 0.847$ ,  $T_{\max} = 0.868$

6846 measured reflections

$k = -14 \rightarrow 8$

$l = -14 \rightarrow 14$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.075$

$S = 1.00$

2448 reflections

189 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0426P)^2 + 1.9365P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.73 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.5000	0.32546 (2)	0.7500	0.03647 (12)
N1	0.5000	0.5148 (2)	0.7500	0.0315 (7)
N2	0.56574 (12)	0.39595 (18)	0.62292 (19)	0.0356 (5)
N3	0.59654 (12)	0.32818 (17)	0.55520 (19)	0.0346 (5)
H3A	0.6204	0.3519	0.5049	0.042*
N4	0.73608 (14)	0.4660 (2)	0.2710 (2)	0.0474 (6)
O1	0.60699 (11)	0.53782 (16)	0.53204 (17)	0.0457 (5)
O2	0.55458 (11)	0.18980 (15)	0.64485 (17)	0.0413 (5)
O3	0.55912 (13)	-0.00807 (17)	0.6368 (2)	0.0596 (6)
H3	0.5574	0.0566	0.6435	0.089*
O4	0.60364 (16)	-0.12393 (19)	0.5313 (2)	0.0733 (8)
O5	0.39670 (11)	0.29907 (15)	0.61261 (16)	0.0420 (5)
H5A	0.3988	0.2396	0.5800	0.050*
H5B	0.3967	0.3475	0.5634	0.050*
O6	0.68682 (14)	0.3634 (2)	0.3932 (2)	0.0656 (7)
C1	0.57294 (15)	0.4982 (2)	0.6033 (2)	0.0345 (6)
C2	0.53537 (14)	0.5671 (2)	0.6791 (2)	0.0340 (6)
C3	0.53679 (16)	0.6767 (2)	0.6772 (3)	0.0403 (7)
H3B	0.5619	0.7127	0.6280	0.048*
C4	0.5000	0.7315 (3)	0.7500	0.0414 (10)

## supplementary materials

---

H4	0.5000	0.8053	0.7500	0.050*
C5	0.58854 (14)	0.2254 (2)	0.5694 (2)	0.0336 (6)
C6	0.62026 (18)	0.1570 (2)	0.4921 (3)	0.0460 (8)
H6	0.6438	0.1921	0.4403	0.055*
C7	0.62029 (19)	0.0514 (2)	0.4854 (3)	0.0523 (8)
H7	0.6419	0.0254	0.4264	0.063*
C8	0.59292 (18)	-0.0317 (2)	0.5527 (3)	0.0483 (8)
C9	0.70087 (17)	0.4503 (3)	0.3561 (3)	0.0519 (8)
H9	0.6854	0.5100	0.3910	0.062*
C10	0.75008 (19)	0.5723 (3)	0.2334 (3)	0.0612 (10)
H10A	0.7260	0.6230	0.2741	0.092*
H10B	0.7320	0.5786	0.1540	0.092*
H10C	0.8016	0.5856	0.2472	0.092*
C11	0.7598 (2)	0.3769 (3)	0.2101 (3)	0.0672 (10)
H11A	0.7465	0.3119	0.2431	0.101*
H11B	0.8117	0.3796	0.2143	0.101*
H11C	0.7368	0.3801	0.1325	0.101*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cd1	0.0481 (2)	0.02519 (17)	0.03978 (19)	0.000	0.01812 (13)	0.000
N1	0.0379 (17)	0.0217 (16)	0.0355 (18)	0.000	0.0083 (14)	0.000
N2	0.0455 (13)	0.0251 (12)	0.0391 (14)	0.0013 (10)	0.0155 (11)	-0.0008 (10)
N3	0.0424 (13)	0.0293 (13)	0.0351 (13)	-0.0007 (10)	0.0150 (10)	0.0013 (10)
N4	0.0528 (15)	0.0427 (15)	0.0515 (16)	0.0070 (12)	0.0226 (13)	0.0104 (12)
O1	0.0627 (13)	0.0346 (12)	0.0450 (12)	0.0011 (10)	0.0240 (10)	0.0079 (9)
O2	0.0546 (12)	0.0284 (11)	0.0466 (12)	0.0010 (9)	0.0247 (10)	-0.0009 (9)
O3	0.0866 (17)	0.0307 (12)	0.0707 (16)	-0.0033 (11)	0.0394 (14)	-0.0011 (11)
O4	0.116 (2)	0.0311 (14)	0.0787 (19)	-0.0008 (13)	0.0324 (16)	-0.0118 (12)
O5	0.0580 (12)	0.0299 (10)	0.0391 (11)	-0.0008 (9)	0.0109 (9)	0.0010 (9)
O6	0.0778 (17)	0.0552 (15)	0.0728 (17)	-0.0020 (13)	0.0393 (14)	0.0140 (13)
C1	0.0398 (15)	0.0314 (16)	0.0321 (15)	-0.0003 (12)	0.0055 (12)	0.0047 (12)
C2	0.0387 (15)	0.0285 (15)	0.0346 (15)	0.0005 (12)	0.0053 (12)	0.0035 (12)
C3	0.0521 (17)	0.0254 (15)	0.0437 (17)	-0.0033 (13)	0.0093 (14)	0.0056 (12)
C4	0.058 (3)	0.020 (2)	0.047 (3)	0.000	0.008 (2)	0.000
C5	0.0384 (15)	0.0285 (16)	0.0358 (16)	0.0001 (12)	0.0119 (12)	0.0030 (12)
C6	0.0576 (19)	0.0354 (18)	0.0506 (19)	0.0015 (14)	0.0253 (15)	0.0005 (14)
C7	0.069 (2)	0.0397 (19)	0.054 (2)	0.0049 (16)	0.0299 (17)	-0.0071 (15)
C8	0.064 (2)	0.0311 (18)	0.0510 (19)	-0.0008 (14)	0.0121 (16)	-0.0051 (14)
C9	0.0539 (19)	0.050 (2)	0.056 (2)	0.0015 (16)	0.0210 (16)	0.0046 (16)
C10	0.062 (2)	0.054 (2)	0.072 (3)	0.0002 (17)	0.0235 (19)	0.0211 (19)
C11	0.079 (3)	0.060 (2)	0.069 (3)	0.012 (2)	0.032 (2)	0.004 (2)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cd1—N2 <sup>i</sup>	2.287 (2)	O5—H5A	0.8500
Cd1—N2	2.287 (2)	O5—H5B	0.8500

Cd1—O5 <sup>i</sup>	2.3412 (19)	O6—C9	1.227 (4)
Cd1—O5	2.3412 (19)	C1—C2	1.511 (4)
Cd1—N1	2.387 (3)	C2—C3	1.382 (4)
Cd1—O2 <sup>i</sup>	2.4441 (19)	C3—C4	1.382 (4)
Cd1—O2	2.4441 (19)	C3—H3B	0.9300
N1—C2 <sup>i</sup>	1.334 (3)	C4—C3 <sup>i</sup>	1.382 (4)
N1—C2	1.334 (3)	C4—H4	0.9300
N2—C1	1.321 (4)	C5—C6	1.460 (4)
N2—N3	1.369 (3)	C6—C7	1.335 (4)
N3—C5	1.319 (3)	C6—H6	0.9300
N3—H3A	0.8600	C7—C8	1.465 (4)
N4—C9	1.316 (4)	C7—H7	0.9300
N4—C11	1.448 (4)	C9—H9	0.9300
N4—C10	1.452 (4)	C10—H10A	0.9600
O1—C1	1.250 (3)	C10—H10B	0.9600
O2—C5	1.269 (3)	C10—H10C	0.9600
O3—C8	1.309 (4)	C11—H11A	0.9600
O3—H3	0.8200	C11—H11B	0.9600
O4—C8	1.214 (4)	C11—H11C	0.9600
N2 <sup>i</sup> —Cd1—N2	134.27 (11)	O1—C1—C2	121.3 (2)
N2 <sup>i</sup> —Cd1—O5 <sup>i</sup>	93.05 (8)	N2—C1—C2	112.5 (2)
N2—Cd1—O5 <sup>i</sup>	93.28 (8)	N1—C2—C3	121.2 (3)
N2 <sup>i</sup> —Cd1—O5	93.28 (8)	N1—C2—C1	115.2 (2)
N2—Cd1—O5	93.05 (8)	C3—C2—C1	123.6 (3)
O5 <sup>i</sup> —Cd1—O5	163.66 (9)	C4—C3—C2	118.5 (3)
N2 <sup>i</sup> —Cd1—N1	67.13 (6)	C4—C3—H3B	120.8
N2—Cd1—N1	67.13 (6)	C2—C3—H3B	120.8
O5 <sup>i</sup> —Cd1—N1	98.17 (5)	C3 <sup>i</sup> —C4—C3	120.0 (4)
O5—Cd1—N1	98.17 (5)	C3 <sup>i</sup> —C4—H4	120.0
N2 <sup>i</sup> —Cd1—O2 <sup>i</sup>	67.27 (7)	C3—C4—H4	120.0
N2—Cd1—O2 <sup>i</sup>	158.46 (8)	O2—C5—N3	121.3 (2)
O5 <sup>i</sup> —Cd1—O2 <sup>i</sup>	84.26 (7)	O2—C5—C6	123.1 (2)
O5—Cd1—O2 <sup>i</sup>	84.33 (7)	N3—C5—C6	115.6 (2)
N1—Cd1—O2 <sup>i</sup>	134.40 (4)	C7—C6—C5	129.2 (3)
N2 <sup>i</sup> —Cd1—O2	158.46 (8)	C7—C6—H6	115.4
N2—Cd1—O2	67.27 (7)	C5—C6—H6	115.4
O5 <sup>i</sup> —Cd1—O2	84.33 (7)	C6—C7—C8	132.6 (3)
O5—Cd1—O2	84.26 (7)	C6—C7—H7	113.7
N1—Cd1—O2	134.40 (4)	C8—C7—H7	113.7
O2 <sup>i</sup> —Cd1—O2	91.20 (9)	O4—C8—O3	119.9 (3)
C2 <sup>i</sup> —N1—C2	120.7 (3)	O4—C8—C7	118.9 (3)
C2 <sup>i</sup> —N1—Cd1	119.64 (16)	O3—C8—C7	121.2 (3)
C2—N1—Cd1	119.64 (16)	O6—C9—N4	125.4 (3)
C1—N2—N3	116.0 (2)	O6—C9—H9	117.3

## supplementary materials

C1—N2—Cd1	125.41 (19)	N4—C9—H9	117.3
N3—N2—Cd1	118.45 (16)	N4—C10—H10A	109.5
C5—N3—N2	118.0 (2)	N4—C10—H10B	109.5
C5—N3—H3A	121.0	H10A—C10—H10B	109.5
N2—N3—H3A	121.0	N4—C10—H10C	109.5
C9—N4—C11	120.5 (3)	H10A—C10—H10C	109.5
C9—N4—C10	121.2 (3)	H10B—C10—H10C	109.5
C11—N4—C10	118.3 (3)	N4—C11—H11A	109.5
C5—O2—Cd1	114.84 (16)	N4—C11—H11B	109.5
C8—O3—H3	109.5	H11A—C11—H11B	109.5
Cd1—O5—H5A	110.8	N4—C11—H11C	109.5
Cd1—O5—H5B	107.1	H11A—C11—H11C	109.5
H5A—O5—H5B	108.0	H11B—C11—H11C	109.5
O1—C1—N2	126.2 (3)		
N2 <sup>i</sup> —Cd1—N1—C2 <sup>i</sup>	0.87 (14)	O5—Cd1—O2—C5	92.38 (19)
N2—Cd1—N1—C2 <sup>i</sup>	-179.13 (14)	N1—Cd1—O2—C5	-3.4 (2)
O5 <sup>i</sup> —Cd1—N1—C2 <sup>i</sup>	-89.00 (14)	O2 <sup>i</sup> —Cd1—O2—C5	176.6 (2)
O5—Cd1—N1—C2 <sup>i</sup>	91.00 (14)	N3—N2—C1—O1	-2.6 (4)
O2 <sup>i</sup> —Cd1—N1—C2 <sup>i</sup>	0.95 (15)	Cd1—N2—C1—O1	-177.9 (2)
O2—Cd1—N1—C2 <sup>i</sup>	-179.05 (15)	N3—N2—C1—C2	178.2 (2)
N2 <sup>i</sup> —Cd1—N1—C2	-179.13 (14)	Cd1—N2—C1—C2	2.8 (3)
N2—Cd1—N1—C2	0.87 (14)	C2 <sup>i</sup> —N1—C2—C3	0.3 (2)
O5 <sup>i</sup> —Cd1—N1—C2	91.00 (14)	Cd1—N1—C2—C3	-179.7 (2)
O5—Cd1—N1—C2	-89.00 (14)	C2 <sup>i</sup> —N1—C2—C1	-179.9 (2)
O2 <sup>i</sup> —Cd1—N1—C2	-179.05 (15)	Cd1—N1—C2—C1	0.1 (2)
O2—Cd1—N1—C2	0.95 (15)	O1—C1—C2—N1	179.0 (2)
N2 <sup>i</sup> —Cd1—N2—C1	-2.1 (2)	N2—C1—C2—N1	-1.7 (3)
O5 <sup>i</sup> —Cd1—N2—C1	-99.6 (2)	O1—C1—C2—C3	-1.2 (4)
O5—Cd1—N2—C1	95.5 (2)	N2—C1—C2—C3	178.1 (3)
N1—Cd1—N2—C1	-2.1 (2)	N1—C2—C3—C4	-0.6 (4)
O2 <sup>i</sup> —Cd1—N2—C1	177.7 (2)	C1—C2—C3—C4	179.6 (2)
O2—Cd1—N2—C1	178.0 (3)	C2—C3—C4—C3 <sup>i</sup>	0.29 (19)
N2 <sup>i</sup> —Cd1—N2—N3	-177.3 (2)	Cd1—O2—C5—N3	3.8 (3)
O5 <sup>i</sup> —Cd1—N2—N3	85.19 (19)	Cd1—O2—C5—C6	-175.4 (2)
O5—Cd1—N2—N3	-79.74 (19)	N2—N3—C5—O2	-1.3 (4)
N1—Cd1—N2—N3	-177.3 (2)	N2—N3—C5—C6	177.9 (2)
O2 <sup>i</sup> —Cd1—N2—N3	2.5 (3)	O2—C5—C6—C7	-0.1 (5)
O2—Cd1—N2—N3	2.74 (17)	N3—C5—C6—C7	-179.3 (4)
C1—N2—N3—C5	-177.8 (2)	C5—C6—C7—C8	-3.2 (7)
Cd1—N2—N3—C5	-2.1 (3)	C6—C7—C8—O4	-176.1 (4)
N2 <sup>i</sup> —Cd1—O2—C5	176.8 (2)	C6—C7—C8—O3	2.0 (6)
N2—Cd1—O2—C5	-3.36 (18)	C11—N4—C9—O6	2.1 (5)
O5 <sup>i</sup> —Cd1—O2—C5	-99.33 (19)	C10—N4—C9—O6	-180.0 (3)

Symmetry codes: (i)  $-x+1, y, -z+3/2$ .

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O5—H5A···O4 <sup>ii</sup>	0.85	1.97	2.802 (3)	165
O5—H5B···O1 <sup>iii</sup>	0.85	1.84	2.685 (3)	174
N3—H3A···O6	0.86	1.97	2.808 (3)	163
O3—H3···O2	0.82	1.68	2.498 (3)	175

Symmetry codes: (ii)  $-x+1, -y, -z+1$ ; (iii)  $-x+1, -y+1, -z+1$ .

Fig. 1




Fig. 2

