

(5E)-Dimethyl 2-bromomethyl-5-cyclohexylimino-2-phenyl-2,5-dihydrofuran-3,4-dicarboxylate

Afsaneh Zonouzi,^a Mojtaba Biniaz,^a Hossein Rahmani^b
and Seik Weng Ng^{c*}

^aDepartment of Chemistry, College of Science, University of Tehran, PO Box 13145-143, Tehran, Iran, ^bInstitute of Chemical Industries, Iranian Research Organization for Science and Technology, PO Box 15815-358, Tehran, Iran, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 8 March 2009; accepted 9 March 2009

Key indicators: single-crystal X-ray study; $T = 123\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.053; wR factor = 0.156; data-to-parameter ratio = 19.1.

The molecule of the title compound, $\text{C}_{21}\text{H}_{24}\text{BrNO}_5$, has a planar furan ring [maximum deviation = 0.025 (3) Å]. The carboxymethyl group in the 3-position is nearly coplanar with this ring [dihedral angle = 7.9 (1)°], whereas that in the 4-position is nearly perpendicular to it [dihedral angle = 78.9 (1) Å].

Related literature

The iminolactone was synthesized by the one-pot, solvent-free reaction of dimethyl acetylenedicarboxylate, cyclohexyl isocyanide and α -bromoacetophenone under microwave irradiation; for other synthetic methods, see: Ma & Xie (2002, 2005); Nair *et al.* (2000); Villemain & Liao (2003).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{24}\text{BrNO}_5$
 $M_r = 450.32$
Monoclinic, $P2_1/c$
 $a = 16.8599 (3)\text{ \AA}$
 $b = 7.2871 (1)\text{ \AA}$
 $c = 17.4145 (3)\text{ \AA}$
 $\beta = 97.330 (1)^\circ$

$V = 2122.06 (6)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.97\text{ mm}^{-1}$
 $T = 123\text{ K}$
 $0.30 \times 0.15 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.590$, $T_{\max} = 0.828$

19142 measured reflections
4873 independent reflections
4034 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.156$
 $S = 1.08$
4873 reflections

255 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 2.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.67\text{ e \AA}^{-3}$

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the Research Council of Tehran University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2389).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ma, S. & Xie, H. (2002). *J. Org. Chem.* **67**, 6575–6578.
- Ma, S. & Xie, H. (2005). *Tetrahedron*, **61**, 251–258.
- Nair, V., Vinod, A. U., Nair, J. S., Sreekanth, A. R. & Rath, N. P. (2000). *Tetrahedron Lett.* **41**, 6675–6679.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Villemain, D. & Liao, L. (2003). *Synth. Commun.* **33**, 1575–1585.
- Westrip, S. P. (2009). publCIF. In preparation.

supporting information

Acta Cryst. (2009). E65, o751 [doi:10.1107/S1600536809008496]

(5E)-Dimethyl 2-bromomethyl-5-cyclohexylimino-2-phenyl-2,5-dihydro-furan-3,4-dicarboxylate

Afsaneh Zonouzi, Mojtaba Biniaz, Hossein Rahmani and Seik Weng Ng


S1. Experimental

To a mixture of 2-bromo-1-phenyl ethanone (α -bromo acetophenone, 0.398 g, 2 mmol) and dimethyl acetylenedicarboxylate (0.25 ml, 2 mmol), cyclohexyl isocyanide (0.25 ml, 2 mmol) was added. Irradiation of the mixture with microwave radiation (180 W) for 5 min produced the title iminolactone. The reaction was monitored by TLC (ethyl acetate *n*-hexane 4:1) until no α -bromoacetophenone was detectable. The product was recrystallized from methanol; yield 90%, m.p. 351 K.

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2 to 1.5 $U_{eq}(C)$.

The final difference Fourier map had a large peak深深 hole in the vicinity of the bromide atom. Attempts to model the bromide atom as being disordered over two positions did not lead to any improvement in the refinement.

**Figure 1**

Thermal ellisoid plot (Barbour, 2001) of $C_{21}H_{24}BrNO_5$; probability levels are set at 70% and H-atoms are drawn as spheres of arbitrary radius.

(5E)-Dimethyl 2-bromomethyl-5-cyclohexylimino-2-phenyl-2,5-dihydrofuran-3,4-dicarboxylate

Crystal data

$C_{21}H_{24}BrNO_5$

$M_r = 450.32$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.8599 (3)$ Å

$b = 7.2871 (1)$ Å

$c = 17.4145 (3)$ Å

$\beta = 97.330 (1)^\circ$

$V = 2122.06 (6)$ Å³

$Z = 4$

$F(000) = 928$

$D_x = 1.410 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3661 reflections

$\theta = 2.7\text{--}28.3^\circ$

$\mu = 1.97 \text{ mm}^{-1}$

$T = 123$ K

Prism, colorless

$0.30 \times 0.15 \times 0.10$ mm

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.590$, $T_{\max} = 0.828$

19142 measured reflections

4873 independent reflections

4034 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.2^\circ$
 $h = -21 \rightarrow 21$

$k = -9 \rightarrow 9$
 $l = -22 \rightarrow 22$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.156$
 $S = 1.08$
4873 reflections
255 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0837P)^2 + 3.6326P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 2.31 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.67 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.25611 (3)	1.02341 (5)	0.45239 (2)	0.03930 (16)
O1	0.29800 (12)	0.6088 (3)	0.49013 (12)	0.0216 (4)
O2	0.09230 (13)	0.3244 (3)	0.36320 (12)	0.0240 (5)
O3	0.11434 (15)	0.5554 (4)	0.28251 (13)	0.0297 (5)
O4	0.06942 (14)	0.8094 (3)	0.54175 (13)	0.0289 (5)
O5	0.02390 (13)	0.6629 (3)	0.43103 (13)	0.0259 (5)
N1	0.30562 (15)	0.4537 (4)	0.37308 (15)	0.0207 (5)
C1	0.23476 (17)	0.6816 (4)	0.53027 (16)	0.0186 (6)
C2	0.16009 (17)	0.6470 (4)	0.47364 (16)	0.0185 (6)
C3	0.17954 (17)	0.5531 (4)	0.41267 (16)	0.0183 (6)
C4	0.26637 (18)	0.5307 (4)	0.42102 (16)	0.0185 (6)
C5	0.25368 (19)	0.8840 (4)	0.54633 (17)	0.0240 (6)
H5A	0.3062	0.8943	0.5787	0.029*
H5B	0.2128	0.9369	0.5759	0.029*
C6	0.23270 (17)	0.5743 (4)	0.60559 (16)	0.0190 (6)
C7	0.25969 (18)	0.3940 (4)	0.61047 (17)	0.0219 (6)
H7	0.2790	0.3382	0.5672	0.026*
C8	0.25852 (19)	0.2952 (4)	0.67858 (19)	0.0251 (6)
H8	0.2771	0.1720	0.6815	0.030*
C9	0.2307 (2)	0.3741 (5)	0.74210 (19)	0.0281 (7)
H9	0.2305	0.3060	0.7886	0.034*
C10	0.2030 (2)	0.5530 (5)	0.73749 (19)	0.0308 (7)
H10	0.1836	0.6080	0.7809	0.037*
C11	0.20363 (19)	0.6531 (5)	0.66923 (18)	0.0252 (6)
H11	0.1841	0.7755	0.6662	0.030*
C12	0.08049 (18)	0.7155 (4)	0.48724 (17)	0.0212 (6)
C13	-0.05632 (18)	0.7330 (5)	0.4356 (2)	0.0291 (7)
H13A	-0.0942	0.6741	0.3957	0.044*
H13B	-0.0571	0.8660	0.4273	0.044*
H13C	-0.0717	0.7059	0.4868	0.044*
C14	0.12496 (18)	0.4812 (4)	0.34460 (17)	0.0194 (6)

C15	0.0330 (2)	0.2459 (5)	0.3044 (2)	0.0329 (8)
H15A	0.0136	0.1294	0.3232	0.049*
H15B	0.0572	0.2240	0.2570	0.049*
H15C	-0.0120	0.3312	0.2934	0.049*
C16	0.39323 (18)	0.4490 (5)	0.38899 (18)	0.0236 (6)
H16	0.4097	0.4544	0.4462	0.028*
C17	0.4273 (2)	0.6150 (5)	0.3507 (2)	0.0330 (8)
H17A	0.4051	0.6189	0.2953	0.040*
H17B	0.4109	0.7288	0.3755	0.040*
C18	0.5186 (2)	0.6070 (6)	0.3579 (2)	0.0382 (9)
H18A	0.5385	0.7119	0.3297	0.046*
H18B	0.5410	0.6176	0.4130	0.046*
C19	0.5467 (2)	0.4285 (7)	0.3250 (2)	0.0402 (9)
H19A	0.5280	0.4226	0.2688	0.048*
H19B	0.6059	0.4249	0.3321	0.048*
C20	0.5149 (2)	0.2658 (6)	0.3648 (2)	0.0405 (9)
H20A	0.5370	0.2663	0.4202	0.049*
H20B	0.5323	0.1512	0.3414	0.049*
C21	0.4232 (2)	0.2705 (5)	0.3573 (2)	0.0338 (8)
H21A	0.4010	0.2575	0.3022	0.041*
H21B	0.4040	0.1656	0.3860	0.041*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0740 (3)	0.0243 (2)	0.0196 (2)	-0.01265 (16)	0.00615 (16)	0.00297 (13)
O1	0.0240 (10)	0.0273 (12)	0.0134 (10)	-0.0023 (8)	0.0023 (8)	-0.0049 (8)
O2	0.0316 (11)	0.0228 (11)	0.0167 (10)	-0.0060 (9)	-0.0010 (8)	-0.0029 (8)
O3	0.0384 (13)	0.0344 (13)	0.0146 (11)	-0.0051 (10)	-0.0029 (9)	0.0058 (9)
O4	0.0321 (12)	0.0330 (13)	0.0214 (11)	0.0071 (10)	0.0022 (9)	-0.0057 (10)
O5	0.0228 (10)	0.0296 (12)	0.0245 (11)	0.0021 (9)	0.0007 (8)	-0.0072 (9)
N1	0.0246 (13)	0.0224 (13)	0.0157 (12)	-0.0007 (10)	0.0047 (9)	-0.0015 (10)
C1	0.0241 (14)	0.0189 (14)	0.0129 (13)	-0.0010 (10)	0.0033 (10)	-0.0019 (10)
C2	0.0261 (14)	0.0156 (13)	0.0132 (12)	-0.0013 (11)	0.0006 (10)	0.0021 (10)
C3	0.0244 (14)	0.0189 (14)	0.0115 (13)	-0.0008 (11)	0.0015 (10)	0.0005 (10)
C4	0.0260 (14)	0.0190 (14)	0.0100 (13)	-0.0037 (11)	0.0007 (10)	-0.0002 (10)
C5	0.0340 (16)	0.0218 (15)	0.0161 (14)	-0.0043 (12)	0.0027 (12)	-0.0008 (11)
C6	0.0209 (13)	0.0217 (14)	0.0139 (13)	-0.0006 (11)	0.0004 (10)	-0.0009 (11)
C7	0.0258 (14)	0.0210 (15)	0.0184 (14)	-0.0001 (11)	0.0007 (11)	-0.0031 (11)
C8	0.0304 (16)	0.0195 (14)	0.0241 (15)	0.0003 (12)	-0.0013 (12)	0.0023 (12)
C9	0.0359 (17)	0.0299 (17)	0.0178 (14)	-0.0010 (13)	0.0009 (12)	0.0061 (13)
C10	0.045 (2)	0.0323 (18)	0.0158 (15)	0.0074 (15)	0.0073 (13)	0.0037 (13)
C11	0.0366 (17)	0.0228 (15)	0.0167 (14)	0.0068 (13)	0.0046 (12)	0.0001 (12)
C12	0.0279 (15)	0.0191 (14)	0.0164 (13)	0.0015 (11)	0.0025 (11)	0.0027 (11)
C13	0.0236 (15)	0.0311 (17)	0.0322 (17)	0.0024 (13)	0.0016 (12)	0.0002 (14)
C14	0.0225 (14)	0.0216 (14)	0.0141 (13)	0.0012 (11)	0.0026 (10)	-0.0018 (11)
C15	0.0348 (18)	0.0362 (19)	0.0265 (17)	-0.0122 (15)	-0.0015 (13)	-0.0112 (14)
C16	0.0241 (15)	0.0324 (17)	0.0141 (14)	-0.0019 (12)	0.0021 (11)	-0.0023 (12)

C17	0.0313 (17)	0.036 (2)	0.0314 (18)	-0.0065 (14)	0.0037 (13)	0.0027 (15)
C18	0.0310 (18)	0.048 (2)	0.036 (2)	-0.0099 (16)	0.0038 (14)	0.0017 (17)
C19	0.0253 (16)	0.063 (3)	0.0332 (19)	-0.0053 (17)	0.0070 (14)	-0.0088 (19)
C20	0.0305 (18)	0.046 (2)	0.044 (2)	0.0075 (16)	0.0026 (15)	-0.0067 (18)
C21	0.0279 (17)	0.0329 (19)	0.041 (2)	0.0002 (14)	0.0073 (14)	-0.0066 (16)

Geometric parameters (\AA , $^\circ$)

Br1—C5	1.930 (3)	C9—H9	0.9500
O1—C4	1.376 (3)	C10—C11	1.396 (4)
O1—C1	1.448 (3)	C10—H10	0.9500
O2—C14	1.326 (4)	C11—H11	0.9500
O2—C15	1.454 (4)	C13—H13A	0.9800
O3—C14	1.202 (4)	C13—H13B	0.9800
O4—C12	1.204 (4)	C13—H13C	0.9800
O5—C12	1.333 (4)	C15—H15A	0.9800
O5—C13	1.457 (4)	C15—H15B	0.9800
N1—C4	1.261 (4)	C15—H15C	0.9800
N1—C16	1.468 (4)	C16—C21	1.524 (5)
C1—C2	1.518 (4)	C16—C17	1.528 (5)
C1—C5	1.528 (4)	C16—H16	1.0000
C1—C6	1.531 (4)	C17—C18	1.529 (5)
C2—C3	1.339 (4)	C17—H17A	0.9900
C2—C12	1.479 (4)	C17—H17B	0.9900
C3—C4	1.462 (4)	C18—C19	1.521 (6)
C3—C14	1.499 (4)	C18—H18A	0.9900
C5—H5A	0.9900	C18—H18B	0.9900
C5—H5B	0.9900	C19—C20	1.506 (6)
C6—C11	1.392 (4)	C19—H19A	0.9900
C6—C7	1.389 (4)	C19—H19B	0.9900
C7—C8	1.390 (4)	C20—C21	1.535 (5)
C7—H7	0.9500	C20—H20A	0.9900
C8—C9	1.381 (5)	C20—H20B	0.9900
C8—H8	0.9500	C21—H21A	0.9900
C9—C10	1.384 (5)	C21—H21B	0.9900
C4—O1—C1	110.3 (2)	O5—C13—H13C	109.5
C14—O2—C15	115.9 (3)	H13A—C13—H13C	109.5
C12—O5—C13	116.2 (2)	H13B—C13—H13C	109.5
C4—N1—C16	119.1 (3)	O3—C14—O2	126.0 (3)
O1—C1—C2	103.1 (2)	O3—C14—C3	124.1 (3)
O1—C1—C5	107.0 (2)	O2—C14—C3	109.9 (2)
C2—C1—C5	114.6 (2)	O2—C15—H15A	109.5
O1—C1—C6	109.0 (2)	O2—C15—H15B	109.5
C2—C1—C6	111.4 (2)	H15A—C15—H15B	109.5
C5—C1—C6	111.3 (2)	O2—C15—H15C	109.5
C3—C2—C12	128.2 (3)	H15A—C15—H15C	109.5
C3—C2—C1	109.5 (3)	H15B—C15—H15C	109.5

C12—C2—C1	122.3 (3)	N1—C16—C21	108.9 (3)
C2—C3—C4	108.8 (2)	N1—C16—C17	108.9 (3)
C2—C3—C14	128.2 (3)	C21—C16—C17	111.0 (3)
C4—C3—C14	123.1 (3)	N1—C16—H16	109.3
N1—C4—O1	125.8 (3)	C21—C16—H16	109.3
N1—C4—C3	126.0 (3)	C17—C16—H16	109.3
O1—C4—C3	108.2 (2)	C18—C17—C16	111.4 (3)
C1—C5—Br1	112.3 (2)	C18—C17—H17A	109.3
C1—C5—H5A	109.1	C16—C17—H17A	109.3
Br1—C5—H5A	109.1	C18—C17—H17B	109.3
C1—C5—H5B	109.1	C16—C17—H17B	109.3
Br1—C5—H5B	109.1	H17A—C17—H17B	108.0
H5A—C5—H5B	107.9	C19—C18—C17	111.2 (3)
C11—C6—C7	119.2 (3)	C19—C18—H18A	109.4
C11—C6—C1	121.3 (3)	C17—C18—H18A	109.4
C7—C6—C1	119.5 (3)	C19—C18—H18B	109.4
C6—C7—C8	120.1 (3)	C17—C18—H18B	109.4
C6—C7—H7	119.9	H18A—C18—H18B	108.0
C8—C7—H7	119.9	C20—C19—C18	110.8 (3)
C9—C8—C7	120.7 (3)	C20—C19—H19A	109.5
C9—C8—H8	119.6	C18—C19—H19A	109.5
C7—C8—H8	119.6	C20—C19—H19B	109.5
C8—C9—C10	119.5 (3)	C18—C19—H19B	109.5
C8—C9—H9	120.3	H19A—C19—H19B	108.1
C10—C9—H9	120.3	C19—C20—C21	110.9 (3)
C9—C10—C11	120.2 (3)	C19—C20—H20A	109.5
C9—C10—H10	119.9	C21—C20—H20A	109.5
C11—C10—H10	119.9	C19—C20—H20B	109.5
C6—C11—C10	120.3 (3)	C21—C20—H20B	109.5
C6—C11—H11	119.9	H20A—C20—H20B	108.0
C10—C11—H11	119.9	C16—C21—C20	111.4 (3)
O4—C12—O5	125.0 (3)	C16—C21—H21A	109.3
O4—C12—C2	123.6 (3)	C20—C21—H21A	109.3
O5—C12—C2	111.4 (3)	C16—C21—H21B	109.3
O5—C13—H13A	109.5	C20—C21—H21B	109.3
O5—C13—H13B	109.5	H21A—C21—H21B	108.0
H13A—C13—H13B	109.5		
C4—O1—C1—C2	-3.2 (3)	C11—C6—C7—C8	-0.8 (4)
C4—O1—C1—C5	-124.4 (2)	C1—C6—C7—C8	179.7 (3)
C4—O1—C1—C6	115.1 (2)	C6—C7—C8—C9	0.0 (5)
O1—C1—C2—C3	4.5 (3)	C7—C8—C9—C10	0.6 (5)
C5—C1—C2—C3	120.4 (3)	C8—C9—C10—C11	-0.3 (5)
C6—C1—C2—C3	-112.2 (3)	C7—C6—C11—C10	1.1 (5)
O1—C1—C2—C12	-174.1 (2)	C1—C6—C11—C10	-179.4 (3)
C5—C1—C2—C12	-58.2 (4)	C9—C10—C11—C6	-0.6 (5)
C6—C1—C2—C12	69.2 (3)	C13—O5—C12—O4	2.8 (5)
C12—C2—C3—C4	174.5 (3)	C13—O5—C12—C2	-176.0 (3)

C1—C2—C3—C4	−4.0 (3)	C3—C2—C12—O4	−175.0 (3)
C12—C2—C3—C14	−4.6 (5)	C1—C2—C12—O4	3.3 (5)
C1—C2—C3—C14	176.9 (3)	C3—C2—C12—O5	3.8 (4)
C16—N1—C4—O1	−2.1 (5)	C1—C2—C12—O5	−177.9 (3)
C16—N1—C4—C3	177.6 (3)	C15—O2—C14—O3	−5.8 (4)
C1—O1—C4—N1	−179.2 (3)	C15—O2—C14—C3	175.4 (3)
C1—O1—C4—C3	1.1 (3)	C2—C3—C14—O3	100.1 (4)
C2—C3—C4—N1	−177.8 (3)	C4—C3—C14—O3	−78.9 (4)
C14—C3—C4—N1	1.4 (5)	C2—C3—C14—O2	−81.1 (4)
C2—C3—C4—O1	2.0 (3)	C4—C3—C14—O2	99.9 (3)
C14—C3—C4—O1	−178.9 (3)	C4—N1—C16—C21	146.3 (3)
O1—C1—C5—Br1	61.4 (3)	C4—N1—C16—C17	−92.6 (3)
C2—C1—C5—Br1	−52.2 (3)	N1—C16—C17—C18	−173.9 (3)
C6—C1—C5—Br1	−179.67 (19)	C21—C16—C17—C18	−54.1 (4)
O1—C1—C6—C11	153.4 (3)	C16—C17—C18—C19	55.4 (4)
C2—C1—C6—C11	−93.6 (3)	C17—C18—C19—C20	−57.0 (4)
C5—C1—C6—C11	35.6 (4)	C18—C19—C20—C21	57.2 (4)
O1—C1—C6—C7	−27.2 (3)	N1—C16—C21—C20	174.2 (3)
C2—C1—C6—C7	85.9 (3)	C17—C16—C21—C20	54.4 (4)
C5—C1—C6—C7	−145.0 (3)	C19—C20—C21—C16	−56.3 (4)