

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(2-Ethoxy-2-methyl-2H-chromen-3-yl)-ethanone

Afsaneh Zonouzi,^a Mojtaba Biniaz,^a Hossein Rahmani^b and Seik Weng Ng^{c*}

^aDepartment of Chemistry, College of Science, University of Tehran, PO Box 13145-143, Tehran, Iran, ^bInstitute of Chemical Industries, Iranian Research Organization for Science and Technology, PO Box 15815-358, Tehran, Iran, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 8 March 2009; accepted 9 March 2009

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.047; wR factor = 0.140; data-to-parameter ratio = 16.9.

The C_{sp^3} atom of the chromenyl fused-ring system in the title compound, $C_{14}H_{16}O_3$, deviates by 0.407 (2) Å from the plane of the other atoms (r.m.s. deviation = 0.041 Å). The ethoxy substituent occupies a pseudo-axial position.

Related literature

For the synthesis, see: Zonouzi *et al.* (2008b). For related crystal structures, see: Bardajee *et al.* (2007); Zhan & Lin (2006); Zonouzi *et al.* (2008a,b).



Experimental

Crystal data

$C_{14}H_{16}O_3$
 $M_r = 232.27$
Monoclinic, $C2/c$
 $a = 20.0084$ (5) Å
 $b = 7.2637$ (2) Å
 $c = 19.1056$ (5) Å
 $\beta = 123.294$ (1)°
 $V = 2320.96$ (11) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 123$ K
 $0.25 \times 0.20 \times 0.15$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: none
7793 measured reflections
2655 independent reflections
2123 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.140$
 $S = 1.03$
2655 reflections
157 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.36$ e Å⁻³
 $\Delta\rho_{min} = -0.43$ e Å⁻³

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Research Council of Tehran University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2390).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bardajee, G. R., Winnik, M. A. & Lough, A. J. (2007). *Acta Cryst.* **E63**, o1513–o1514.
Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2009). *publCIF*. In preparation.
Zhan, Y. & Lin, S.-K. (2006). *Acta Cryst.* **E62**, o2097–o2098.
Zonouzi, A., Rahmani, H. & Jahangiri, A. (2008a). *Z. Kristallogr. New Cryst. Struct.* **223**, 155–156.
Zonouzi, A., Sadeghi Googheri, M. & Miraliaaghi, P. S. (2008b). *Org. Prep. Proced. Int.* **40**, 383–389.

supplementary materials

Acta Cryst. (2009). E65, o752 [doi:10.1107/S1600536809008502]

1-(2-Ethoxy-2-methyl-2*H*-chromen-3-yl)ethanone

A. Zonouzi, M. Biniiaz, H. Rahmani and S. W. Ng

Comment

(type here to add)

Experimental

The compound was synthesized by using a reported method (Zonouzi *et al.*, 2008*b*). Crystals were obtained by recrystallization from ethanol.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2 to 1.5 $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{14}\text{H}_{16}\text{O}_3$; probability levels are set at 70% and H-atoms are drawn as spheres of arbitrary radius.

1-(2-Ethoxy-2-methyl-2*H*-chromen-3-yl)ethanone

Crystal data

$\text{C}_{14}\text{H}_{16}\text{O}_3$

$M_r = 232.27$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 20.0084$ (5) Å

$b = 7.2637$ (2) Å

$c = 19.1056$ (5) Å

$\beta = 123.294$ (1)°

$V = 2320.96$ (11) Å³

$Z = 8$

$F_{000} = 992$

$D_x = 1.329$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2233 reflections

$\theta = 2.4\text{--}28.1^\circ$

$\mu = 0.09$ mm⁻¹

$T = 123$ K

Irregular block, colorless

$0.25 \times 0.20 \times 0.15$ mm

supplementary materials

Data collection

Bruker SMART APEX diffractometer	2123 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.035$
Monochromator: graphite	$\theta_{\text{max}} = 27.5^\circ$
$T = 123$ K	$\theta_{\text{min}} = 2.4^\circ$
ω scans	$h = -21 \rightarrow 25$
Absorption correction: None	$k = -9 \rightarrow 9$
7793 measured reflections	$l = -24 \rightarrow 23$
2655 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.047$	H-atom parameters constrained
$wR(F^2) = 0.140$	$w = 1/[\sigma^2(F_o^2) + (0.0739P)^2 + 1.4605P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
2655 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
157 parameters	$\Delta\rho_{\text{max}} = 0.36 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.43 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.13572 (6)	0.55707 (15)	0.32959 (6)	0.0208 (3)
O2	0.17621 (6)	0.25748 (14)	0.32719 (6)	0.0211 (3)
O3	0.36134 (7)	0.35871 (19)	0.38669 (7)	0.0350 (3)
C1	0.19962 (9)	0.4438 (2)	0.33800 (9)	0.0193 (3)
C2	0.27375 (9)	0.4496 (2)	0.42713 (9)	0.0189 (3)
C3	0.26440 (9)	0.4851 (2)	0.49036 (9)	0.0198 (3)
H3	0.3098	0.4792	0.5463	0.024*
C4	0.18788 (9)	0.5318 (2)	0.47624 (9)	0.0190 (3)
C5	0.17449 (9)	0.5519 (2)	0.54044 (9)	0.0223 (3)
H5	0.2169	0.5303	0.5969	0.027*
C6	0.10015 (10)	0.6026 (2)	0.52253 (10)	0.0244 (3)
H6	0.0917	0.6164	0.5666	0.029*
C7	0.03757 (10)	0.6336 (2)	0.43999 (10)	0.0239 (3)
H7	-0.0135	0.6686	0.4280	0.029*
C8	0.04917 (9)	0.6137 (2)	0.37524 (9)	0.0222 (3)
H8	0.0063	0.6340	0.3189	0.027*
C9	0.12413 (9)	0.5638 (2)	0.39363 (9)	0.0185 (3)
C10	0.20936 (10)	0.5212 (2)	0.27015 (9)	0.0250 (4)

H10A	0.1566	0.5384	0.2185	0.037*
H10B	0.2407	0.4352	0.2597	0.037*
H10C	0.2371	0.6399	0.2885	0.037*
C11	0.11063 (10)	0.2021 (2)	0.24546 (9)	0.0282 (4)
H11A	0.1250	0.2167	0.2037	0.034*
H11B	0.0628	0.2781	0.2278	0.034*
C12	0.09445 (12)	0.0053 (3)	0.25244 (12)	0.0409 (5)
H12A	0.0504	-0.0390	0.1978	0.061*
H12B	0.0798	-0.0068	0.2935	0.061*
H12C	0.1425	-0.0678	0.2706	0.061*
C13	0.35266 (9)	0.3964 (2)	0.44326 (9)	0.0218 (3)
C14	0.42348 (9)	0.3868 (2)	0.53284 (9)	0.0256 (4)
H14A	0.4721	0.3623	0.5337	0.038*
H14B	0.4152	0.2876	0.5621	0.038*
H14C	0.4291	0.5043	0.5608	0.038*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0232 (5)	0.0224 (6)	0.0196 (5)	0.0066 (4)	0.0136 (4)	0.0041 (4)
O2	0.0220 (6)	0.0178 (6)	0.0183 (5)	0.0008 (4)	0.0077 (4)	0.0006 (4)
O3	0.0311 (7)	0.0482 (8)	0.0313 (6)	0.0106 (6)	0.0207 (5)	0.0023 (5)
C1	0.0206 (7)	0.0186 (7)	0.0207 (7)	0.0030 (6)	0.0126 (6)	0.0013 (5)
C2	0.0208 (7)	0.0146 (7)	0.0216 (7)	0.0000 (6)	0.0119 (6)	0.0015 (5)
C3	0.0203 (7)	0.0171 (7)	0.0202 (7)	-0.0003 (6)	0.0101 (6)	0.0003 (5)
C4	0.0222 (7)	0.0155 (7)	0.0212 (7)	-0.0011 (6)	0.0131 (6)	-0.0001 (5)
C5	0.0258 (8)	0.0206 (8)	0.0211 (7)	0.0000 (6)	0.0132 (6)	0.0001 (6)
C6	0.0323 (9)	0.0222 (8)	0.0270 (7)	0.0007 (7)	0.0215 (7)	-0.0003 (6)
C7	0.0238 (8)	0.0213 (8)	0.0321 (8)	0.0023 (6)	0.0189 (7)	0.0005 (6)
C8	0.0224 (8)	0.0199 (7)	0.0236 (7)	0.0030 (6)	0.0122 (6)	0.0017 (6)
C9	0.0237 (7)	0.0142 (7)	0.0200 (7)	0.0002 (6)	0.0135 (6)	0.0006 (5)
C10	0.0318 (9)	0.0237 (8)	0.0238 (7)	0.0049 (7)	0.0180 (7)	0.0043 (6)
C11	0.0257 (8)	0.0265 (9)	0.0207 (7)	-0.0003 (7)	0.0054 (6)	-0.0016 (6)
C12	0.0394 (11)	0.0386 (11)	0.0360 (9)	-0.0071 (9)	0.0151 (8)	-0.0031 (8)
C13	0.0229 (8)	0.0176 (7)	0.0264 (7)	0.0003 (6)	0.0145 (6)	0.0007 (6)
C14	0.0201 (7)	0.0249 (8)	0.0299 (8)	-0.0002 (6)	0.0125 (6)	-0.0008 (6)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.3654 (17)	C7—C8	1.386 (2)
O1—C1	1.4531 (18)	C7—H7	0.9500
O2—C1	1.4101 (18)	C8—C9	1.388 (2)
O2—C11	1.4397 (17)	C8—H8	0.9500
O3—C13	1.2164 (18)	C10—H10A	0.9800
C1—C10	1.520 (2)	C10—H10B	0.9800
C1—C2	1.5259 (19)	C10—H10C	0.9800
C2—C3	1.345 (2)	C11—C12	1.488 (3)
C2—C13	1.484 (2)	C11—H11A	0.9900
C3—C4	1.442 (2)	C11—H11B	0.9900

supplementary materials

C3—H3	0.9500	C12—H12A	0.9800
C4—C5	1.400 (2)	C12—H12B	0.9800
C4—C9	1.4006 (19)	C12—H12C	0.9800
C5—C6	1.381 (2)	C13—C14	1.510 (2)
C5—H5	0.9500	C14—H14A	0.9800
C6—C7	1.392 (2)	C14—H14B	0.9800
C6—H6	0.9500	C14—H14C	0.9800
C9—O1—C1	119.55 (11)	O1—C9—C4	120.41 (13)
C1—O2—C11	117.31 (11)	C8—C9—C4	121.18 (13)
O2—C1—O1	109.01 (12)	C1—C10—H10A	109.5
O2—C1—C10	114.69 (12)	C1—C10—H10B	109.5
O1—C1—C10	102.32 (11)	H10A—C10—H10B	109.5
O2—C1—C2	103.36 (11)	C1—C10—H10C	109.5
O1—C1—C2	111.41 (11)	H10A—C10—H10C	109.5
C10—C1—C2	116.12 (13)	H10B—C10—H10C	109.5
C3—C2—C13	121.33 (13)	O2—C11—C12	106.55 (13)
C3—C2—C1	118.58 (13)	O2—C11—H11A	110.4
C13—C2—C1	119.74 (13)	C12—C11—H11A	110.4
C2—C3—C4	122.28 (13)	O2—C11—H11B	110.4
C2—C3—H3	118.9	C12—C11—H11B	110.4
C4—C3—H3	118.9	H11A—C11—H11B	108.6
C5—C4—C9	118.45 (14)	C11—C12—H12A	109.5
C5—C4—C3	123.72 (13)	C11—C12—H12B	109.5
C9—C4—C3	117.81 (13)	H12A—C12—H12B	109.5
C6—C5—C4	120.60 (14)	C11—C12—H12C	109.5
C6—C5—H5	119.7	H12A—C12—H12C	109.5
C4—C5—H5	119.7	H12B—C12—H12C	109.5
C5—C6—C7	120.05 (14)	O3—C13—C2	121.91 (13)
C5—C6—H6	120.0	O3—C13—C14	119.59 (14)
C7—C6—H6	120.0	C2—C13—C14	118.49 (13)
C8—C7—C6	120.47 (14)	C13—C14—H14A	109.5
C8—C7—H7	119.8	C13—C14—H14B	109.5
C6—C7—H7	119.8	H14A—C14—H14B	109.5
C7—C8—C9	119.25 (14)	C13—C14—H14C	109.5
C7—C8—H8	120.4	H14A—C14—H14C	109.5
C9—C8—H8	120.4	H14B—C14—H14C	109.5
O1—C9—C8	118.29 (13)		
C11—O2—C1—O1	66.57 (15)	C3—C4—C5—C6	177.92 (15)
C11—O2—C1—C10	-47.43 (18)	C4—C5—C6—C7	0.3 (2)
C11—O2—C1—C2	-174.83 (12)	C5—C6—C7—C8	0.1 (2)
C9—O1—C1—O2	75.85 (15)	C6—C7—C8—C9	-0.5 (2)
C9—O1—C1—C10	-162.31 (12)	C1—O1—C9—C8	-157.01 (13)
C9—O1—C1—C2	-37.58 (17)	C1—O1—C9—C4	26.77 (19)
O2—C1—C2—C3	-90.42 (15)	C7—C8—C9—O1	-175.65 (13)
O1—C1—C2—C3	26.51 (19)	C7—C8—C9—C4	0.5 (2)
C10—C1—C2—C3	143.09 (15)	C5—C4—C9—O1	175.92 (13)
O2—C1—C2—C13	82.91 (16)	C3—C4—C9—O1	-2.3 (2)
O1—C1—C2—C13	-160.17 (13)	C5—C4—C9—C8	-0.2 (2)

supplementary materials

C10—C1—C2—C13	-43.59 (19)	C3—C4—C9—C8	-178.45 (14)
C13—C2—C3—C4	-178.17 (14)	C1—O2—C11—C12	-176.06 (14)
C1—C2—C3—C4	-5.0 (2)	C3—C2—C13—O3	177.78 (15)
C2—C3—C4—C5	173.50 (14)	C1—C2—C13—O3	4.6 (2)
C2—C3—C4—C9	-8.3 (2)	C3—C2—C13—C14	-1.4 (2)
C9—C4—C5—C6	-0.2 (2)	C1—C2—C13—C14	-174.50 (13)

Fig. 1

