

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

 Ethyl 3-amino-1-phenyl-1*H*-benzo[*f*]-chromene-2-carboxylate

 Yuan-Hong Jiao,<sup>a</sup> Qian Zhang,<sup>a\*</sup> Fa-Yan Meng,<sup>a</sup> Lei Teng,<sup>a</sup> Jia Yuan<sup>a</sup> and Seik Weng Ng<sup>b</sup>
<sup>a</sup>School of Chemistry and Material Engineering, Huangshi Institute of Technology, Huangshi 435003, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: jiaoyuanhong1995@yahoo.com.cn


Received 5 March 2009; accepted 11 March 2009

 Key indicators: single-crystal X-ray study;  $T = 153$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.030;  $wR$  factor = 0.078; data-to-parameter ratio = 8.3.

The pyranyl ring of the title compound,  $\text{C}_{22}\text{H}_{19}\text{NO}_3$ , adopts a flattened-boat conformation. The dihedral angle between naphthalene and phenyl rings is  $78.3(1)^\circ$ . The molecule also features an intramolecular  $\text{N}-\text{H}\cdots\text{O}_{\text{carbonyl}}$  hydrogen bond. Adjacent molecules are linked by an intermolecular  $\text{N}-\text{H}\cdots\text{O}_{\text{carbonyl}}$  hydrogen bond, forming a zigzag chain that runs along the  $c$  axis.

## Related literature

For the crystal structures of other ethyl 3-amino-1-aryl-1*H*-benzo[*f*]chromene-2-carboxylate derivatives, see: Klokol *et al.* (1987); Shi *et al.* (2003*a,b*); Wang *et al.* (2003); Zhuang *et al.* (2003*a,b*).



## Experimental

## Crystal data

$\text{C}_{22}\text{H}_{19}\text{NO}_3$	$V = 1766.05(9)$ Å <sup>3</sup>
$M_r = 345.38$	$Z = 4$
Monoclinic, $Cc$	Mo $K\alpha$ radiation
$a = 13.7835(5)$ Å	$\mu = 0.09$ mm <sup>-1</sup>
$b = 14.6460(4)$ Å	$T = 153$ K
$c = 8.8713(2)$ Å	$0.36 \times 0.25 \times 0.14$ mm
$\beta = 99.551(1)^\circ$	

## Data collection

Rigaku R-Axis RAPID diffractometer  
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)  
 $T_{\text{min}} = 0.864$ ,  $T_{\text{max}} = 0.988$

8563 measured reflections  
2029 independent reflections  
1958 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.016$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.078$   
 $S = 1.09$   
2029 reflections  
244 parameters  
4 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.19$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.27$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H11}\cdots\text{O3}^i$	0.88 (1)	2.05 (1)	2.907 (2)	166 (2)
$\text{N1}-\text{H12}\cdots\text{O3}$	0.88 (1)	2.07 (2)	2.707 (2)	129 (2)

 Symmetry code: (i)  $x, -y + 2, z + \frac{1}{2}$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Huangshi Institute of Technology (grant No. 08yjz23B) and the University of Malaya for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2186).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
Klokol, G. V., Sharanina, L. G., Nesterov, V. N., Shklover, V. E., Sharanin, Yu. A. & Struchkov, Yu. T. (1987). *Russ. J. Org. Chem.* **23**, 412–421.  
Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.  
Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Shi, D.-Q., Wang, J., Wang, X., Zhuang, Q. & Yu, K. (2003*a*). *Acta Cryst.* **E59**, o1733–o1734.  
Shi, D., Wu, H., Wang, X., Zhuang, Q. & Hu, H. (2003*b*). *Acta Cryst.* **E59**, o1263–o1264.  
Wang, X.-S., Shi, D.-Q. & Tu, S.-J. (2003). *Chin. J. Chem.* **21**, 1114–1117.  
Westrip, S. P. (2009). *publCIF*. In preparation.  
Zhuang, Q.-Y., Rong, L.-C., Wang, J.-X., Tu, S.-J., Wang, X.-S. & Shi, D.-Q. (2003*a*). *Chin. J. Org. Chem.* **23**, 671–673.  
Zhuang, Q.-Y., Zhang, S., Shi, D.-Q., Tu, S.-J. & Wang, X.-S. (2003*b*). *J. Xuzhou Normal Univ. (Natur. Sci.)*, **21**, 49–52.

**supplementary materials**

*Acta Cryst.* (2009). E65, o775 [ doi:10.1107/S1600536809008885 ]

## Ethyl 3-amino-1-phenyl-1*H*-benzo[*f*]chromene-2-carboxylate

Y.-H. Jiao, Q. Zhang, F.-Y. Meng, L. Teng, J. Yuan and S. W. Ng

### Comment

2-Naphthol, substituted benzaldehydes and ethyl 2-cyanoacetate react in the presence of a catalyst to form the ethyl 3-amino-1-aryl-1*H*-benzo[*f*]chromene-2-carboxylates; the compounds possess a primary amine group. The crystal structures of several 1-aryl derivatives have been reported (Klokol *et al.*, 1987; Shi *et al.*, 2003*a*; Shi *et al.*, 2003*b*; Wang *et al.*, 2003; Zhuang *et al.*, 2003*a*; Zhuang *et al.*, 2003*b*). Interestingly, the crystal structure of the unsubstituted 1-phenyl compound was not known, and its structure is reported here. The title compound (Scheme I, Fig. 1) exhibits a pyranyl ring in a flattened boat conformation. The molecule also features an intramolecular *N*—*H*⋯*O*<sub>carbonyl</sub> bond. Adjacent molecules are linked by an intermolecular *N*—*H*⋯*O*<sub>carbonyl</sub> hydrogen bond to furnish a zigzag chain that runs along the *c*-axis.

### Experimental

2-Naphthol (1.4 g, 10 mmol), benzaldehyde (1.1 g, 10 mmol), ethyl 2-cyanoacetate (1.0 g, 10 mmol) and piperidine (1 ml) were dissolved in ethanol (30 ml). The solution was heated for 5 h. The solvent was removed under reduced pressure and the residue recrystallized from dichloromethane/methanol (1:1/v:v) to give yellow crystals in 80% yield; m.p. 436–437 K.

### Refinement

In the absence of significant anomalous scattering Friedel pairs were merged prior to refinement and the absolute configuration of the molecule was not refined. Carbon-bound H atoms were placed in calculated positions [C—H 0.95–1.00 Å and  $U_{\text{iso}}(\text{H})$  1.2–1.5 $U_{\text{eq}}(\text{C})$ ], and were included in the refinement in the riding-model approximation. The amino H-atoms were located in a difference Fourier map, and were refined with an N—H distance restraint of 0.88 (1) Å; their isotropic temperature factors were refined.

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{22}\text{H}_{19}\text{NO}_3$ ; displacement ellipsoids are drawn at the 70% probability level, and H atoms as spheres of arbitrary radius.




Fig. 2. Thermal ellipsoid plot (Barbour, 2001) showing the intra and intermolecular hydrogen bonding interactions (dashed lines); displacement ellipsoids are drawn at the 70% probability level, and H atoms as spheres of arbitrary radius.

## Ethyl 3-amino-1-phenyl-1H-benzo[f]chromene-2-carboxylate

### Crystal data

$C_{22}H_{19}NO_3$	$F_{000} = 728$
$M_r = 345.38$	$D_x = 1.299 \text{ Mg m}^{-3}$
Monoclinic, $Cc$	Mo $K\alpha$ radiation
Hall symbol: C -2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 13.7835 (5) \text{ \AA}$	Cell parameters from 8058 reflections
$b = 14.6460 (4) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$c = 8.8713 (2) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 99.551 (1)^\circ$	$T = 153 \text{ K}$
$V = 1766.05 (9) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.36 \times 0.25 \times 0.14 \text{ mm}$

### Data collection

Rigaku R-Axis RAPID diffractometer	2029 independent reflections
Radiation source: fine-focus sealed tube	1958 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.016$
Detector resolution: $10.000 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.5^\circ$
$T = 153 \text{ K}$	$\theta_{\text{min}} = 3.0^\circ$
$\omega$ scans	$h = -17 \rightarrow 17$
Absorption correction: Multi-scan (ABSCOR; Higashi, 1995)	$k = -18 \rightarrow 18$
$T_{\text{min}} = 0.864$ , $T_{\text{max}} = 0.988$	$l = -10 \rightarrow 11$
8563 measured reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.078$	$w = 1/[\sigma^2(F_o^2) + (0.0522P)^2 + 0.3553P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
2029 reflections	$(\Delta/\sigma)_{\max} = 0.001$
244 parameters	$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
4 restraints	$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.50000 (10)	0.89966 (8)	0.50000 (13)	0.0251 (3)
O2	0.57192 (9)	0.73346 (8)	0.07916 (12)	0.0221 (3)
O3	0.57203 (10)	0.88667 (8)	0.05998 (13)	0.0243 (3)
N1	0.54563 (13)	0.98073 (10)	0.31277 (17)	0.0289 (3)
H11	0.5480 (18)	1.0271 (12)	0.376 (2)	0.038 (7)*
H12	0.5641 (18)	0.9858 (18)	0.2234 (17)	0.040 (7)*
C1	0.53026 (11)	0.72524 (11)	0.36834 (17)	0.0176 (3)
H1	0.4968	0.6810	0.2908	0.021*
C2	0.63064 (12)	0.68559 (10)	0.43558 (18)	0.0199 (3)
C3	0.66322 (13)	0.60409 (11)	0.3813 (2)	0.0254 (4)
H3	0.6228	0.5728	0.3004	0.031*
C4	0.75447 (14)	0.56763 (13)	0.4441 (3)	0.0335 (4)
H4	0.7759	0.5118	0.4060	0.040*
C5	0.81378 (14)	0.61264 (14)	0.5619 (3)	0.0364 (5)
H5	0.8756	0.5874	0.6056	0.044*
C6	0.78287 (14)	0.69460 (15)	0.6159 (2)	0.0357 (5)
H6	0.8239	0.7260	0.6959	0.043*
C7	0.69180 (13)	0.73093 (13)	0.5532 (2)	0.0276 (4)
H7	0.6711	0.7871	0.5908	0.033*
C8	0.46549 (12)	0.73777 (11)	0.48933 (17)	0.0191 (3)
C9	0.41664 (12)	0.66266 (12)	0.54835 (17)	0.0204 (3)
C10	0.43046 (13)	0.57101 (12)	0.50594 (18)	0.0247 (3)
H10	0.4732	0.5580	0.4349	0.030*
C11	0.38294 (15)	0.50035 (13)	0.5660 (2)	0.0305 (4)
H11A	0.3941	0.4392	0.5378	0.037*
C12	0.31800 (14)	0.51858 (14)	0.6691 (2)	0.0322 (4)
H12A	0.2840	0.4698	0.7079	0.039*
C13	0.30372 (14)	0.60584 (14)	0.7134 (2)	0.0297 (4)
H13	0.2601	0.6172	0.7837	0.036*
C14	0.35287 (12)	0.68005 (13)	0.65636 (19)	0.0240 (4)
C15	0.34044 (13)	0.77096 (13)	0.7054 (2)	0.0278 (4)
H15	0.2974	0.7827	0.7764	0.033*
C16	0.38932 (14)	0.84127 (13)	0.65223 (19)	0.0262 (4)
H16	0.3816	0.9018	0.6867	0.031*
C17	0.45205 (12)	0.82298 (12)	0.54462 (18)	0.0217 (3)
C18	0.53012 (12)	0.89655 (11)	0.36102 (17)	0.0213 (3)

## supplementary materials

---

C19	0.53928 (12)	0.81562 (10)	0.28807 (18)	0.0186 (3)
C20	0.56246 (12)	0.81763 (10)	0.13523 (18)	0.0185 (3)
C21	0.59893 (13)	0.72831 (11)	-0.07209 (18)	0.0239 (3)
H21A	0.6644	0.7561	-0.0717	0.029*
H21B	0.5502	0.7611	-0.1475	0.029*
C22	0.60081 (17)	0.62865 (13)	-0.1116 (2)	0.0336 (4)
H22A	0.5345	0.6031	-0.1183	0.050*
H22B	0.6459	0.5964	-0.0321	0.050*
H22C	0.6232	0.6214	-0.2101	0.050*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0364 (7)	0.0209 (6)	0.0193 (5)	-0.0010 (5)	0.0086 (5)	-0.0031 (4)
O2	0.0328 (6)	0.0182 (5)	0.0168 (5)	0.0025 (5)	0.0084 (4)	0.0014 (4)
O3	0.0334 (7)	0.0200 (6)	0.0199 (5)	0.0007 (5)	0.0059 (5)	0.0027 (4)
N1	0.0473 (9)	0.0188 (7)	0.0216 (7)	-0.0023 (6)	0.0088 (6)	-0.0033 (6)
C1	0.0189 (7)	0.0185 (7)	0.0156 (6)	0.0004 (5)	0.0036 (5)	0.0004 (5)
C2	0.0194 (7)	0.0223 (7)	0.0190 (7)	-0.0004 (6)	0.0061 (5)	0.0069 (6)
C3	0.0223 (8)	0.0188 (7)	0.0357 (10)	-0.0010 (6)	0.0060 (7)	0.0053 (6)
C4	0.0243 (8)	0.0223 (8)	0.0553 (12)	0.0029 (7)	0.0108 (8)	0.0109 (8)
C5	0.0202 (8)	0.0387 (11)	0.0489 (12)	0.0022 (7)	0.0012 (8)	0.0183 (9)
C6	0.0245 (9)	0.0507 (12)	0.0298 (9)	-0.0019 (8)	-0.0014 (7)	0.0060 (8)
C7	0.0236 (8)	0.0359 (9)	0.0230 (8)	-0.0006 (7)	0.0026 (6)	0.0006 (7)
C8	0.0187 (7)	0.0215 (7)	0.0168 (7)	0.0006 (6)	0.0024 (6)	0.0015 (6)
C9	0.0187 (7)	0.0267 (8)	0.0155 (6)	-0.0001 (6)	0.0017 (5)	0.0015 (6)
C10	0.0279 (8)	0.0250 (8)	0.0221 (8)	-0.0044 (6)	0.0067 (6)	0.0003 (6)
C11	0.0365 (10)	0.0277 (8)	0.0272 (9)	-0.0071 (8)	0.0053 (7)	0.0014 (7)
C12	0.0309 (9)	0.0365 (10)	0.0296 (9)	-0.0103 (7)	0.0059 (7)	0.0065 (8)
C13	0.0225 (8)	0.0440 (10)	0.0239 (8)	-0.0020 (7)	0.0078 (6)	0.0057 (7)
C14	0.0190 (7)	0.0320 (9)	0.0208 (8)	0.0011 (6)	0.0030 (6)	0.0024 (6)
C15	0.0241 (8)	0.0376 (10)	0.0233 (8)	0.0063 (7)	0.0085 (6)	-0.0007 (7)
C16	0.0277 (8)	0.0276 (9)	0.0240 (8)	0.0056 (7)	0.0063 (7)	-0.0033 (7)
C17	0.0228 (8)	0.0251 (8)	0.0173 (7)	0.0017 (6)	0.0031 (6)	0.0007 (6)
C18	0.0254 (8)	0.0226 (8)	0.0153 (7)	0.0002 (6)	0.0020 (6)	0.0005 (6)
C19	0.0206 (7)	0.0177 (7)	0.0169 (7)	0.0001 (5)	0.0016 (6)	0.0018 (5)
C20	0.0199 (7)	0.0173 (7)	0.0180 (7)	0.0018 (5)	0.0021 (5)	0.0012 (5)
C21	0.0330 (9)	0.0233 (8)	0.0173 (8)	0.0011 (7)	0.0092 (6)	-0.0003 (6)
C22	0.0519 (11)	0.0238 (8)	0.0280 (9)	0.0062 (8)	0.0155 (8)	-0.0017 (7)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—C18	1.3654 (19)	C8—C9	1.433 (2)
O1—C17	1.393 (2)	C9—C10	1.415 (2)
O2—C20	1.3437 (19)	C9—C14	1.427 (2)
O2—C21	1.4527 (18)	C10—C11	1.378 (2)
O3—C20	1.231 (2)	C10—H10	0.9500
N1—C18	1.334 (2)	C11—C12	1.407 (3)
N1—H11	0.876 (10)	C11—H11A	0.9500

N1—H12	0.875 (10)	C12—C13	1.361 (3)
C1—C8	1.517 (2)	C12—H12A	0.9500
C1—C19	1.518 (2)	C13—C14	1.418 (3)
C1—C2	1.528 (2)	C13—H13	0.9500
C1—H1	1.0000	C14—C15	1.420 (3)
C2—C3	1.389 (2)	C15—C16	1.357 (3)
C2—C7	1.396 (2)	C15—H15	0.9500
C3—C4	1.395 (2)	C16—C17	1.416 (2)
C3—H3	0.9500	C16—H16	0.9500
C4—C5	1.382 (3)	C18—C19	1.366 (2)
C4—H4	0.9500	C19—C20	1.444 (2)
C5—C6	1.386 (3)	C21—C22	1.502 (2)
C5—H5	0.9500	C21—H21A	0.9900
C6—C7	1.392 (3)	C21—H21B	0.9900
C6—H6	0.9500	C22—H22A	0.9800
C7—H7	0.9500	C22—H22B	0.9800
C8—C17	1.365 (2)	C22—H22C	0.9800
C18—O1—C17	117.34 (13)	C12—C11—H11A	119.9
C20—O2—C21	116.42 (12)	C13—C12—C11	120.25 (17)
C18—N1—H11	120.2 (17)	C13—C12—H12A	119.9
C18—N1—H12	117.1 (17)	C11—C12—H12A	119.9
H11—N1—H12	122 (2)	C12—C13—C14	121.08 (17)
C8—C1—C19	109.27 (13)	C12—C13—H13	119.5
C8—C1—C2	111.78 (13)	C14—C13—H13	119.5
C19—C1—C2	112.04 (12)	C13—C14—C15	121.35 (16)
C8—C1—H1	107.9	C13—C14—C9	119.15 (16)
C19—C1—H1	107.9	C15—C14—C9	119.50 (15)
C2—C1—H1	107.9	C16—C15—C14	120.85 (16)
C3—C2—C7	118.56 (15)	C16—C15—H15	119.6
C3—C2—C1	121.00 (14)	C14—C15—H15	119.6
C7—C2—C1	120.44 (15)	C15—C16—C17	118.99 (16)
C2—C3—C4	120.82 (17)	C15—C16—H16	120.5
C2—C3—H3	119.6	C17—C16—H16	120.5
C4—C3—H3	119.6	C8—C17—O1	122.40 (14)
C5—C4—C3	120.03 (18)	C8—C17—C16	123.36 (16)
C5—C4—H4	120.0	O1—C17—C16	114.24 (14)
C3—C4—H4	120.0	N1—C18—O1	110.36 (14)
C4—C5—C6	119.83 (17)	N1—C18—C19	128.13 (15)
C4—C5—H5	120.1	O1—C18—C19	121.50 (14)
C6—C5—H5	120.1	C18—C19—C20	118.64 (14)
C5—C6—C7	120.10 (18)	C18—C19—C1	120.87 (14)
C5—C6—H6	120.0	C20—C19—C1	120.42 (13)
C7—C6—H6	120.0	O3—C20—O2	121.82 (15)
C6—C7—C2	120.65 (18)	O3—C20—C19	125.89 (14)
C6—C7—H7	119.7	O2—C20—C19	112.28 (13)
C2—C7—H7	119.7	O2—C21—C22	106.43 (13)
C17—C8—C9	118.06 (14)	O2—C21—H21A	110.4
C17—C8—C1	119.73 (14)	C22—C21—H21A	110.4
C9—C8—C1	122.20 (14)	O2—C21—H21B	110.4

## supplementary materials

C10—C9—C14	118.23 (15)	C22—C21—H21B	110.4
C10—C9—C8	122.59 (15)	H21A—C21—H21B	108.6
C14—C9—C8	119.17 (15)	C21—C22—H22A	109.5
C11—C10—C9	121.04 (16)	C21—C22—H22B	109.5
C11—C10—H10	119.5	H22A—C22—H22B	109.5
C9—C10—H10	119.5	C21—C22—H22C	109.5
C10—C11—C12	120.21 (18)	H22A—C22—H22C	109.5
C10—C11—H11A	119.9	H22B—C22—H22C	109.5
C8—C1—C2—C3	121.86 (16)	C8—C9—C14—C15	-1.2 (2)
C19—C1—C2—C3	-115.10 (16)	C13—C14—C15—C16	178.73 (17)
C8—C1—C2—C7	-58.27 (19)	C9—C14—C15—C16	-0.8 (3)
C19—C1—C2—C7	64.76 (19)	C14—C15—C16—C17	1.0 (3)
C7—C2—C3—C4	0.8 (2)	C9—C8—C17—O1	176.85 (14)
C1—C2—C3—C4	-179.31 (16)	C1—C8—C17—O1	-3.5 (2)
C2—C3—C4—C5	0.0 (3)	C9—C8—C17—C16	-2.7 (2)
C3—C4—C5—C6	-0.8 (3)	C1—C8—C17—C16	176.99 (16)
C4—C5—C6—C7	0.8 (3)	C18—O1—C17—C8	24.8 (2)
C5—C6—C7—C2	0.0 (3)	C18—O1—C17—C16	-155.62 (15)
C3—C2—C7—C6	-0.8 (3)	C15—C16—C17—C8	0.8 (3)
C1—C2—C7—C6	179.31 (16)	C15—C16—C17—O1	-178.81 (15)
C19—C1—C8—C17	-20.86 (19)	C17—O1—C18—N1	160.53 (15)
C2—C1—C8—C17	103.73 (16)	C17—O1—C18—C19	-18.3 (2)
C19—C1—C8—C9	158.81 (13)	N1—C18—C19—C20	-4.4 (3)
C2—C1—C8—C9	-76.60 (18)	O1—C18—C19—C20	174.24 (15)
C17—C8—C9—C10	-175.95 (15)	N1—C18—C19—C1	172.70 (16)
C1—C8—C9—C10	4.4 (2)	O1—C18—C19—C1	-8.7 (2)
C17—C8—C9—C14	2.8 (2)	C8—C1—C19—C18	27.1 (2)
C1—C8—C9—C14	-176.84 (15)	C2—C1—C19—C18	-97.31 (17)
C14—C9—C10—C11	0.6 (2)	C8—C1—C19—C20	-155.85 (14)
C8—C9—C10—C11	179.41 (15)	C2—C1—C19—C20	79.71 (17)
C9—C10—C11—C12	1.2 (3)	C21—O2—C20—O3	3.0 (2)
C10—C11—C12—C13	-1.8 (3)	C21—O2—C20—C19	-177.75 (13)
C11—C12—C13—C14	0.5 (3)	C18—C19—C20—O3	-2.5 (2)
C12—C13—C14—C15	-178.19 (18)	C1—C19—C20—O3	-179.59 (16)
C12—C13—C14—C9	1.3 (3)	C18—C19—C20—O2	178.28 (15)
C10—C9—C14—C13	-1.9 (2)	C1—C19—C20—O2	1.20 (19)
C8—C9—C14—C13	179.31 (15)	C20—O2—C21—C22	-177.74 (14)
C10—C9—C14—C15	177.66 (15)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H11 $\cdots$ O3 <sup>i</sup>	0.88 (1)	2.05 (1)	2.907 (2)	166 (2)
N1—H12 $\cdots$ O3	0.88 (1)	2.07 (2)	2.707 (2)	129 (2)

Symmetry codes: (i)  $x, -y+2, z+1/2$ .

Fig. 1




Fig. 2

