

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

 (3*R*,6*S*,7*aS*)-3-Phenyl-6-(phenylsulfanyl)-perhydropyrrolo[1,2-*c*]oxazol-5-one

Graeme J. Gainsford,* Andreas Luxenburger and Anthony D. Woolhouse

Industrial Research Limited, PO Box 31-310, Lower Hutt, New Zealand

Correspondence e-mail: g.gainsford@irl.cri.nz


Received 2 March 2009; accepted 29 March 2009

 Key indicators: single-crystal X-ray study; $T = 121$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.033; wR factor = 0.087; data-to-parameter ratio = 29.9.

Molecules of the title compound [systematic name: (2*R*,5*S*,7*S*)-2-phenyl-7-phenylsulfanyl-1-aza-3-oxabicyclo[3.3.0]octan-8-one], $\text{C}_{18}\text{H}_{17}\text{NO}_2\text{S}$, form high quality crystals even though they are only packed using $\text{C}-\text{H}\cdots\text{O}$ (carbonyl) and weak $\text{C}-\text{H}\cdots\text{S}$ interactions. The dihedral angle between the aromatic rings is $85.53(5)^\circ$. The fused rings adopt envelope and twist conformations.

Related literature

For related structures, see Nagasaka & Imai (1995); Anwar *et al.* (2003); Bailey *et al.* (2000); McCarthy *et al.* (1999). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

 $\text{C}_{18}\text{H}_{17}\text{NO}_2\text{S}$
 $M_r = 311.39$

 Orthorhombic, $P2_12_12_1$
 $a = 5.3884(3)$ Å

 $b = 11.2227(7)$ Å

 $c = 25.3308(16)$ Å

 $V = 1531.81(16)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.22$ mm⁻¹
 $T = 121$ K

 $0.75 \times 0.39 \times 0.30$ mm

Data collection

Bruker–Nonius APEXII CCD area-detector diffractometer

Absorption correction: multi-scan (Blessing, 1995)

 $T_{\min} = 0.829$, $T_{\max} = 0.937$

45370 measured reflections

5948 independent reflections

 5711 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.087$
 $S = 1.11$

5948 reflections

199 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Absolute structure: Flack (1983),

2508 Friedel pairs

Flack parameter: 0.01 (4)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C7A-H7A\cdots O5^i$	1.00	2.55	3.4305 (13)	147
$C7-H72\cdots O5^{ii}$	0.99	2.62	3.3493 (16)	131
$C1-H1B\cdots O5^{ii}$	0.99	2.71	3.1513 (13)	108
$C1-H1A\cdots S1^{ii}$	0.99	3.00	3.9512 (14)	162
$C4-H4\cdots S1^{iii}$	0.95	2.98	3.7531 (13)	139

 Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT and SADABS (Bruker, 2006); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997), Mercury (Macrae *et al.*, 2006) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97 and PLATON.

We thank Dr J. Wikaira of the University of Canterbury for her assistance in data collection, as well as the New Zealand Foundation for Science and Technology and New Zealand Pharmaceuticals Ltd for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ER2064).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Anwar, M., Bailey, J. H., Dickinson, L. C., Edwards, H. J., Goswami, R. & Moloney, M. G. (2003). *Org. Biomol. Chem.* **1**, 2364–2376.
- Bailey, J. H., Byfield, A. T. J., Davis, P. J., Foster, A. C., Leech, M., Moloney, M. G., Muller, M. & Prout, C. K. (2000). *J. Chem. Soc. Perkin Trans. 1*, pp. 1977–1982.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Bruker (2006). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- McCarthy, D. G., Collins, C. C., O'Driscoll, J. P. & Lawrence, S. E. (1999). *J. Chem. Soc. Perkin Trans. 1*, pp. 3667–3675.
- Nagasaka, T. & Imai, T. (1995). *Chem. Pharm. Bull.* **43**, 1081–1088.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o943 [doi:10.1107/S1600536809011623]

(3*R*,6*S*,7*aS*)-3-Phenyl-6-(phenylsulfanyl)perhydropyrrolo[1,2-*c*]oxazol-5-one

G. J. Gainsford, A. Luxenburger and A. D. Woolhouse

Comment

During the course of our efforts to delineate, in detail, the conversion of the bicyclic lactam 1 into its highly prized, chiral α,β -unsaturated derivative 5 employing *sulfur chemistry* (Fig. 3), we were able to isolate and separate the three anticipated products (2,3 & 4) from the rapid quenching of the lithium enolate of 1 with the electrophile, PhSSPh: the two 6-(phenylthio)-mono-adducts 2 and 3 were found to be predominant in the mixture with up to 10% only of the 6,6-bis(phenylthio)-adduct. In order to be able to unequivocally assign structures to each of the mono-adducts, we carried out an X-ray crystallographic study of the title *cis(endo-)* adduct 2 (Scheme) so that structural inferences made on the basis of nOe experiments could be corroborated (see bottom right entry in fig. 3).

The 2-(*p*-methoxyphenyl) derivative is described (molecule 21*b*) by Nagasaka & Imai (1995). Related compounds in the Cambridge Structural Database [C.S.D. Version 5.30 with November 2008 updates (Allen, 2002)] have been reported by Anwar *et al.* (2003) (IJAGUV) and Bailey *et al.* (2000) (QINMEF & QINMEJ).

The asymmetric unit is shown in Figure 1. The 5-membered fused rings (N4,C5,C6,C7,C7a) and (C1,O2,C3,N4,C7a) are best described as having envelope (on C6) and twist (on C1—O2) conformations (Spek, 2009). The 3-phenyl (C2,C4,C9—C12) and 6-phenylthio (C13—C18) phenyl rings subtend angles of 85.53 (5)° to each other and 71.74 (5) and 39.74 (5)° respectively to the mean plane through the fused ring (octan-8-one) atoms. The corresponding three angles for the 6-benzyl closest structural relative (QINMEF) are 80.14 (7), 71.18 (6) and 36.82 (7)° while the interplanar angles found for the 3-phenyl rings in IJAGUV & QINMIJ structures are 74.1 (4) and 76.73 (7)%, respectively.

The molecular packing is provided by mainly weak C—H \cdots O interactions (entries 1–3, Table 1, involving a bifurcated O) as partly shown in Figure 2. The weak, just significant, C—H \cdots S interactions have been observed before contributing to dimer formation in CUQXUH with a C—H \cdots S angles of 140° (McCarthy *et al.*, 1999).

These crystals were of superb quality, confirmed by the final agreement indices and difference density maps, which raises the issue of how many of the weak intermolecular interactions may just be fortuitous, with packing largely determined by the overall molecular shape and van der Waal's forces.

Experimental

To a solution of the lactam 1 (19.7 g, 97.0 mmol) in dry THF (180 ml) was added dropwise LiHMDS (1 *M* solution in hexanes; 116 ml, 116 mmol) at -78 °C and the solution was stirred for further 60 min before a concentrated solution of diphenylsulfide (25.4 g, 116 mmol) in dry THF (25 ml) was added rapidly in one single portion at the same temperature. After being stirred for 3 h the reaction mixture was quenched at -78 °C with saturated aqueous ammonium chloride solution (50 ml) and allowed to warm to ambient temperature before being diluted with ethyl acetate (400 ml). The separated organic phase was washed with water and brine, dried over MgSO₄ and concentrated to give a crude oil, the composition of which was determined by HPLC analysis. Based on this analysis the two diastereomeric phenylthioethers 2 and 3 were formed in

supplementary materials

40% and 49% yield, respectively. In addition, the reaction provided the bis(phenylthio)-adduct 4 in 9% as the by-product and 2% of the starting material remained unreacted. The crude product was finally fractionated by flash column chromatography (silica gel, 10% and 20% ethyl acetate/petroleum ether) to yield the three adducts 2, 3 and 4.

(2): (3*R*,6*S*,7*aS*)-3-Phenyl-6-(phenylthio)tetrahydropyrrolo [1,2-*c*]oxazol-5(1*H*)-one was recrystallized from ethyl acetate/petroleum ether to give colourless crystals, $R_f = 0.44$ (30% ethyl acetate/petroleum ether); m.p. 97–98 °C, $[\alpha]_D^{21} = +215.6$ (c 0.545, CHCl₃); FTIR (neat) 1692 (C=O) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.57–7.53 (m, 2H), 7.46–7.42 (m, 2H), 7.38–7.27 (m, 6H), 6.29 (s, 1H, H3), 4.26 (t, $J = 9.7$ Hz, 1H, H6), 4.17 (dd, $J = 8.2, 6.2$ Hz, 1H, H1 α), 4.06–3.99 (m, 1H, H7 $\alpha\alpha$), 3.21 (t, $J = 8.1$ Hz, 1H, H1 β), 2.81 (ddd, $J = 13.5, 9.2, 7.3$ Hz, 1H, H7 α), 1.93 (ddd, $J = 13.5, 10.2, 6.4$ Hz, 1H, H7 β); ¹³C NMR (125 MHz, CDCl₃) δ 173.54, 138.18, 132.91, 132.85, 128.94, 128.55, 128.29, 127.89, 125.82, 87.18, 71.88, 55.73, 51.06, 32.03; HRMS (ES+) m/z calcd for C₁₈H₁₇NO₂SNa⁺ 334.0878, obsd 334.0872; Anal. calcd for C₁₈H₁₇NO₂S: C, 69.43; H, 5.50; N, 4.50. Found: C, 69.32; H, 5.63; N, 4.63.

(3): (3*R*,6*R*,7*aS*)-3-Phenyl-6-(phenylthio)tetrahydropyrrolo [1,2-*c*]oxazol-5(1*H*)-one, colourless crystals, $R_f = 0.30$ (30% ethyl acetate/petroleum ether); m.p. 80 °C; $[\alpha]_D^{20} = +102.3$ (c 1.05, CHCl₃); FTIR (neat) 1702 (C=O) cm⁻¹; ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.52–7.48 (m, 2H); 7.41–7.30 (m, 6H); 7.27–7.23 (m, 2H), 6.05 (s, 1H, H3); 4.22 (dd, $J = 9.4, 3.3$ Hz, 1H, H6); 4.13 (dd, $J = 8.0, 6.2$ Hz, 1H, H1 α); 3.90–3.83 (m, 1H, H7 $\alpha\alpha$); 3.47 (t, $J = 8.4$ Hz, 1H, H1 β); 2.56 (ddd, $J = 14.5, 9.2, 5.3$ Hz, 1H, H7 β); 2.28 (ddd, $J = 14.2, 7.5, 3.3$ Hz, 1H, H7 α); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 174.58, 138.57, 132.55, 131.99, 129.02, 128.49, 128.25, 127.76, 125.93, 86.47, 70.97, 57.11, 49.25, 30.66; HRMS (ES+) m/z calcd for C₁₈H₁₇NO₂SNa⁺ 334.0878, obsd 334.0879; Anal. calcd for C₁₈H₁₇NO₂S: C, 69.43; H, 5.50; N, 4.50. Found C, 69.39; H, 5.59; N, 4.55.

(4): (3*R*,7*aS*)-3-Phenyl-6,6- bis(phenylthio)tetrahydropyrrolo[1,2-*c*]oxazol-5(1*H*)-one, colourless crystals, $R_f = 0.54$ (30% ethyl acetate/petroleum ether); m.p. 108 °C; $[\alpha]_D^{20} = +157.1$ (c 0.62, CDCl₃); FTIR (neat) 1698 (C=O) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.71–7.68 (m, 2H), 7.59–7.55 (m, 2H), 7.41–7.29 (m, 9H), 7.25–7.20 (m, 2H), 6.19 (s, 1H), 4.03 (dd, $J = 8.1, 6.3$ Hz, 1H), 3.69–3.63 (m, 1H), 3.09 (t, $J = 8.1$ Hz, 1H), 2.54 (dd, $J = 14.3, 7.0$ Hz, 1H), 2.35 (dd, $J = 14.3, 6.5$ Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 171.71, 137.95, 136.65, 136.27, 131.18, 130.44, 129.96, 129.57, 129.01, 128.87, 128.67, 128.35, 126.02, 87.09, 71.78, 68.68, 55.00, 39.79; HRMS (ES+) m/z calcd for C₂₄H₂₁NO₂S₂Na⁺ 442.0911, obsd 442.0191; Anal. calcd for C₂₄H₂₁NO₂S₂: C, 68.70; H, 5.04; N, 3.34. Found C, 68.88; H, 5.29; N, 3.37.

Refinement

A total of 9 reflections within 2 θ 68° were omitted from refinement as either outliers or partially screened by the backstop. All H atoms bound to carbon were constrained to their expected geometries (C—H 0.95, 0.99 & 1.00 Å). All H atoms were refined with U_{iso} 1.2 times that of the U_{eq} of their parent atom.

Figures




Fig. 1. Molecular structure of the asymmetric unit (Farrugia, 1997); displacement ellipsoids are shown at the 50% probability level.




Fig. 2. Packing diagram (Mercury; Macrae *et al.*, 2006) of the unit cell. Contact atoms are shown as balls; not all interactions and labels are shown for clarity (see Table 1).



Fig. 3. Preparation of the title compound.

(2*R*,5*S*,7*S*)-2-phenyl-7-phenylsulfonyl-1-aza-3-oxabicyclo[3.3.0]octan-8-one

Crystal data

$C_{18}H_{17}NO_2S$

$M_r = 311.39$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 5.3884$ (3) Å

$b = 11.2227$ (7) Å

$c = 25.3308$ (16) Å

$V = 1531.81$ (16) Å³

$Z = 4$

$F(000) = 656$

$D_x = 1.350$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9806 reflections

$\theta = 3.0$ – 34.4°

$\mu = 0.22$ mm⁻¹

$T = 121$ K

Block, colourless

$0.75 \times 0.39 \times 0.30$ mm

Data collection

Bruker–Nonius APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

Detector resolution: 8.333 pixels mm⁻¹

ϕ and ω scans

Absorption correction: multi-scan (Blessing, 1995)

$T_{\min} = 0.829$, $T_{\max} = 0.937$

45370 measured reflections

5948 independent reflections

5711 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 34.0^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -8 \rightarrow 8$

$k = -17 \rightarrow 16$

$l = -38 \rightarrow 38$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.033$	H-atom parameters constrained
$wR(F^2) = 0.087$	$w = 1/[\sigma^2(F_o^2) + (0.0528P)^2 + 0.1721P]$
$S = 1.11$	where $P = (F_o^2 + 2F_c^2)/3$
5948 reflections	$(\Delta/\sigma)_{\max} = 0.001$
199 parameters	$\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 2508 Friedel pairs Flack parameter: 0.01 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.53402 (5)	0.52168 (2)	0.147539 (9)	0.02284 (6)
O2	0.63842 (15)	0.59460 (6)	0.36457 (3)	0.02259 (14)
O5	0.76332 (18)	0.36178 (6)	0.23809 (3)	0.02532 (15)
N4	0.80755 (16)	0.53105 (7)	0.28760 (3)	0.01873 (14)
C1	0.5697 (2)	0.67901 (8)	0.32437 (4)	0.02362 (18)
H1A	0.5735	0.7616	0.3381	0.028*
H1B	0.4020	0.6619	0.3103	0.028*
C2	0.88109 (18)	0.41716 (8)	0.37229 (3)	0.01718 (15)
C3	0.70664 (17)	0.48760 (8)	0.33825 (3)	0.01756 (15)
H3	0.5543	0.4392	0.3313	0.021*
C4	0.8234 (2)	0.29955 (8)	0.38433 (4)	0.02221 (17)
H4	0.6793	0.2640	0.3695	0.027*
C5	0.77799 (18)	0.46990 (8)	0.24169 (3)	0.01866 (15)
C6	0.7641 (2)	0.56099 (8)	0.19659 (4)	0.02006 (16)
H6	0.9305	0.5691	0.1795	0.024*
C7	0.6940 (3)	0.67833 (8)	0.22432 (4)	0.0274 (2)
H71	0.7857	0.7462	0.2087	0.033*

H72	0.5138	0.6938	0.2212	0.033*
C7A	0.7678 (2)	0.66023 (8)	0.28236 (4)	0.02122 (17)
H7A	0.9223	0.7055	0.2911	0.025*
C9	0.9747 (2)	0.23345 (9)	0.41785 (5)	0.02726 (19)
H9	0.9343	0.1531	0.4259	0.033*
C10	1.1843 (2)	0.28528 (10)	0.43943 (4)	0.0288 (2)
H10	1.2878	0.2406	0.4624	0.035*
C11	1.2439 (2)	0.40310 (10)	0.42745 (5)	0.0274 (2)
H11	1.3877	0.4386	0.4424	0.033*
C12	1.09340 (18)	0.46886 (9)	0.39371 (4)	0.02237 (17)
H12	1.1353	0.5489	0.3853	0.027*
C13	0.71008 (18)	0.45302 (7)	0.09728 (4)	0.01856 (15)
C14	0.9116 (2)	0.37876 (9)	0.10770 (4)	0.02280 (18)
H14	0.9565	0.3611	0.1431	0.027*
C15	1.0467 (2)	0.33070 (10)	0.06609 (4)	0.02566 (19)
H15	1.1859	0.2813	0.0732	0.031*
C16	0.9801 (2)	0.35426 (9)	0.01414 (4)	0.02609 (19)
H16	1.0746	0.3221	-0.0142	0.031*
C17	0.7743 (2)	0.42516 (9)	0.00390 (4)	0.02471 (18)
H17	0.7254	0.4397	-0.0316	0.030*
C18	0.63983 (19)	0.47488 (9)	0.04504 (4)	0.02167 (16)
H18	0.5001	0.5238	0.0377	0.026*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.02433 (10)	0.02516 (10)	0.01902 (10)	0.00568 (8)	-0.00096 (8)	-0.00069 (8)
O2	0.0304 (3)	0.0205 (3)	0.0169 (3)	0.0075 (3)	0.0009 (3)	-0.0014 (2)
O5	0.0404 (4)	0.0151 (3)	0.0205 (3)	0.0020 (3)	-0.0005 (3)	-0.0027 (2)
N4	0.0270 (4)	0.0144 (3)	0.0148 (3)	0.0010 (3)	0.0016 (3)	-0.0007 (2)
C1	0.0323 (5)	0.0203 (4)	0.0183 (4)	0.0070 (3)	-0.0006 (3)	-0.0002 (3)
C2	0.0203 (4)	0.0172 (3)	0.0141 (3)	0.0012 (3)	0.0009 (3)	-0.0011 (2)
C3	0.0218 (4)	0.0166 (3)	0.0143 (3)	0.0006 (3)	0.0007 (3)	-0.0006 (2)
C4	0.0250 (4)	0.0179 (3)	0.0236 (4)	-0.0004 (3)	-0.0009 (3)	-0.0001 (3)
C5	0.0243 (4)	0.0167 (3)	0.0150 (3)	0.0018 (3)	0.0006 (3)	-0.0009 (3)
C6	0.0282 (4)	0.0173 (3)	0.0147 (4)	0.0012 (3)	0.0007 (3)	0.0002 (3)
C7	0.0481 (6)	0.0161 (3)	0.0180 (4)	0.0049 (4)	0.0023 (4)	0.0002 (3)
C7A	0.0305 (5)	0.0140 (3)	0.0191 (4)	-0.0005 (3)	0.0014 (3)	-0.0016 (3)
C9	0.0311 (5)	0.0220 (4)	0.0287 (5)	0.0036 (4)	0.0001 (4)	0.0047 (3)
C10	0.0285 (5)	0.0328 (5)	0.0251 (5)	0.0093 (4)	-0.0033 (4)	0.0021 (4)
C11	0.0213 (4)	0.0334 (5)	0.0276 (5)	0.0029 (4)	-0.0047 (4)	-0.0034 (4)
C12	0.0197 (4)	0.0229 (4)	0.0245 (4)	-0.0016 (3)	0.0002 (3)	-0.0016 (3)
C13	0.0228 (4)	0.0169 (3)	0.0160 (3)	-0.0008 (3)	-0.0019 (3)	0.0000 (3)
C14	0.0280 (5)	0.0238 (4)	0.0167 (4)	0.0053 (3)	-0.0025 (3)	0.0008 (3)
C15	0.0268 (4)	0.0281 (4)	0.0222 (4)	0.0062 (4)	-0.0019 (4)	-0.0027 (3)
C16	0.0301 (5)	0.0293 (4)	0.0189 (4)	0.0002 (4)	0.0015 (4)	-0.0029 (3)
C17	0.0311 (5)	0.0272 (4)	0.0158 (4)	-0.0028 (4)	-0.0047 (4)	0.0004 (3)
C18	0.0251 (4)	0.0216 (4)	0.0183 (4)	-0.0007 (3)	-0.0053 (3)	0.0022 (3)

supplementary materials

Geometric parameters (Å, °)

S1—C13	1.7648 (10)	C7—H71	0.9900
S1—C6	1.8098 (10)	C7—H72	0.9900
O2—C3	1.4217 (11)	C7A—H7A	1.0000
O2—C1	1.4393 (12)	C9—C10	1.3831 (17)
O5—C5	1.2194 (11)	C9—H9	0.9500
N4—C5	1.3598 (11)	C10—C11	1.3940 (17)
N4—C7A	1.4716 (11)	C10—H10	0.9500
N4—C3	1.4762 (11)	C11—C12	1.3901 (15)
C1—C7A	1.5218 (15)	C11—H11	0.9500
C1—H1A	0.9900	C12—H12	0.9500
C1—H1B	0.9900	C13—C14	1.3942 (13)
C2—C4	1.3900 (12)	C13—C18	1.3981 (12)
C2—C12	1.3927 (13)	C14—C15	1.3899 (14)
C2—C3	1.5007 (13)	C14—H14	0.9500
C3—H3	1.0000	C15—C16	1.3896 (14)
C4—C9	1.3913 (15)	C15—H15	0.9500
C4—H4	0.9500	C16—C17	1.3892 (16)
C5—C6	1.5348 (13)	C16—H16	0.9500
C6—C7	1.5395 (13)	C17—C18	1.3866 (15)
C6—H6	1.0000	C17—H17	0.9500
C7—C7A	1.5363 (14)	C18—H18	0.9500
C13—S1—C6	103.50 (5)	H71—C7—H72	108.8
C3—O2—C1	106.90 (7)	N4—C7A—C1	100.10 (8)
C5—N4—C7A	113.75 (8)	N4—C7A—C7	104.74 (7)
C5—N4—C3	122.24 (8)	C1—C7A—C7	118.00 (10)
C7A—N4—C3	110.50 (7)	N4—C7A—H7A	111.1
O2—C1—C7A	102.90 (8)	C1—C7A—H7A	111.1
O2—C1—H1A	111.2	C7—C7A—H7A	111.1
C7A—C1—H1A	111.2	C10—C9—C4	119.71 (10)
O2—C1—H1B	111.2	C10—C9—H9	120.1
C7A—C1—H1B	111.2	C4—C9—H9	120.1
H1A—C1—H1B	109.1	C9—C10—C11	120.05 (10)
C4—C2—C12	119.60 (9)	C9—C10—H10	120.0
C4—C2—C3	119.09 (9)	C11—C10—H10	120.0
C12—C2—C3	121.26 (8)	C12—C11—C10	120.22 (10)
O2—C3—N4	102.93 (7)	C12—C11—H11	119.9
O2—C3—C2	109.72 (7)	C10—C11—H11	119.9
N4—C3—C2	116.27 (8)	C11—C12—C2	119.82 (9)
O2—C3—H3	109.2	C11—C12—H12	120.1
N4—C3—H3	109.2	C2—C12—H12	120.1
C2—C3—H3	109.2	C14—C13—C18	119.67 (9)
C2—C4—C9	120.59 (10)	C14—C13—S1	122.90 (7)
C2—C4—H4	119.7	C18—C13—S1	117.43 (7)
C9—C4—H4	119.7	C15—C14—C13	119.76 (9)
O5—C5—N4	125.01 (9)	C15—C14—H14	120.1
O5—C5—C6	127.15 (8)	C13—C14—H14	120.1

N4—C5—C6	107.84 (7)	C16—C15—C14	120.60 (10)
C5—C6—C7	104.01 (7)	C16—C15—H15	119.7
C5—C6—S1	112.46 (6)	C14—C15—H15	119.7
C7—C6—S1	110.72 (8)	C17—C16—C15	119.47 (10)
C5—C6—H6	109.8	C17—C16—H16	120.3
C7—C6—H6	109.8	C15—C16—H16	120.3
S1—C6—H6	109.8	C18—C17—C16	120.49 (9)
C7A—C7—C6	105.07 (8)	C18—C17—H17	119.8
C7A—C7—H71	110.7	C16—C17—H17	119.8
C6—C7—H71	110.7	C17—C18—C13	119.95 (9)
C7A—C7—H72	110.7	C17—C18—H18	120.0
C6—C7—H72	110.7	C13—C18—H18	120.0
C3—O2—C1—C7A	-42.19 (10)	C5—N4—C7A—C1	124.87 (9)
C1—O2—C3—N4	30.62 (10)	C3—N4—C7A—C1	-17.02 (10)
C1—O2—C3—C2	155.00 (8)	C5—N4—C7A—C7	2.19 (12)
C5—N4—C3—O2	-145.28 (9)	C3—N4—C7A—C7	-139.70 (9)
C7A—N4—C3—O2	-7.18 (10)	O2—C1—C7A—N4	34.70 (9)
C5—N4—C3—C2	94.76 (11)	O2—C1—C7A—C7	147.49 (8)
C7A—N4—C3—C2	-127.14 (9)	C6—C7—C7A—N4	-14.67 (12)
C4—C2—C3—O2	125.46 (9)	C6—C7—C7A—C1	-124.87 (9)
C12—C2—C3—O2	-51.84 (11)	C2—C4—C9—C10	0.09 (16)
C4—C2—C3—N4	-118.31 (9)	C4—C9—C10—C11	-0.26 (17)
C12—C2—C3—N4	64.39 (11)	C9—C10—C11—C12	-0.13 (17)
C12—C2—C4—C9	0.47 (15)	C10—C11—C12—C2	0.70 (16)
C3—C2—C4—C9	-176.88 (9)	C4—C2—C12—C11	-0.86 (14)
C7A—N4—C5—O5	-167.92 (11)	C3—C2—C12—C11	176.43 (9)
C3—N4—C5—O5	-31.03 (16)	C6—S1—C13—C14	37.85 (9)
C7A—N4—C5—C6	11.47 (12)	C6—S1—C13—C18	-142.53 (7)
C3—N4—C5—C6	148.36 (8)	C18—C13—C14—C15	2.42 (15)
O5—C5—C6—C7	159.26 (12)	S1—C13—C14—C15	-177.97 (8)
N4—C5—C6—C7	-20.11 (11)	C13—C14—C15—C16	-1.15 (17)
O5—C5—C6—S1	39.41 (14)	C14—C15—C16—C17	-0.94 (17)
N4—C5—C6—S1	-139.96 (7)	C15—C16—C17—C18	1.75 (16)
C13—S1—C6—C5	-98.36 (7)	C16—C17—C18—C13	-0.48 (15)
C13—S1—C6—C7	145.76 (7)	C14—C13—C18—C17	-1.62 (14)
C5—C6—C7—C7A	20.79 (12)	S1—C13—C18—C17	178.75 (8)
S1—C6—C7—C7A	141.81 (8)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C7A—H7A \cdots O5 ⁱ	1.00	2.55	3.4305 (13)	147
C7—H72 \cdots O5 ⁱⁱ	0.99	2.62	3.3493 (16)	131
C1—H1B \cdots O5 ⁱⁱ	0.99	2.71	3.1513 (13)	108
C1—H1A \cdots S1 ⁱⁱ	0.99	3.00	3.9512 (14)	162
C4—H4 \cdots S1 ⁱⁱⁱ	0.95	2.98	3.7531 (13)	139

Symmetry codes: (i) $-x+2, y+1/2, -z+1/2$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $-x+1, y-1/2, -z+1/2$.

Fig. 1




Fig. 2




Fig. 3

