

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[[pyridinecopper(II)]-(μ -2-oxidonaphthalene-1-carbaldehyde picolinoylhydrazonato)-[pyridinecopper(II)]- μ -sulfato] diethyl ether hemisolvate]

Quanchang Wang, Dacheng Li* and Daqi Wang

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: lidacheng62@lcu.edu.cn


Received 18 February 2009; accepted 3 April 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.045; wR factor = 0.130; data-to-parameter ratio = 13.2.

The title compound, $[\text{Cu}_2(\text{C}_{17}\text{H}_{11}\text{N}_3\text{O})(\text{SO}_4)(\text{C}_5\text{H}_5\text{N})_2] \cdot 0.5\text{C}_4\text{H}_{10}\text{O}]_n$, was synthesized by the reaction of 2-hydroxy-1-naphthylaldehyde-2-pyridinecarboxylhydrazone with copper sulfonate. A one-dimensional polymer was obtained *via* self-assembly. Each Cu ion is located in a distorted square-pyramidal coordination environment, with one Cu ion coordinated by two N and three O atoms, while the other links to two O and three N atoms. In the crystal, weak intermolecular $\text{C}-\text{H} \cdots \text{O}$ interactions connect the chains into a two-dimensional network.

Related literature

For the biological activity of aroylhydrazones, see Armstrong *et al.* (2003). For the crystal structure of a copper complex with a related picolinoylhydrazone derivative, see: Bai *et al.* (2006).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_{17}\text{H}_{11}\text{N}_3\text{O})(\text{SO}_4)(\text{C}_5\text{H}_5\text{N})_2] \cdot 0.5\text{C}_4\text{H}_{10}\text{O}$
 $M_r = 707.69$
 Monoclinic, $C2/c$

$a = 26.484$ (2) Å
 $b = 14.0374$ (15) Å
 $c = 16.8083$ (17) Å
 $\beta = 108.404$ (2)°

$V = 5929.2$ (10) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 1.56$ mm⁻¹
 $T = 298$ K
 $0.38 \times 0.32 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.589$, $T_{\max} = 0.789$

14633 measured reflections
 5215 independent reflections
 3268 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.130$
 $S = 1.01$
 5215 reflections

394 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.85$ e Å⁻³
 $\Delta\rho_{\min} = -0.49$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—O2	1.888 (3)	Cu2—O3	2.011 (3)
Cu1—N2	1.957 (4)	Cu2—N3	2.015 (4)
Cu1—O1	1.964 (3)	Cu2—N1	2.016 (4)
Cu1—N4	2.008 (4)	Cu2—O4	2.036 (3)
Cu1—O5 ⁱ	2.389 (3)	Cu2—N5	2.189 (4)

Symmetry code: (i) $x, -y + 2, z + \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{C}26-\text{H}26 \cdots \text{O}2^{\text{i}}$	0.93	2.46	3.386 (7)	174
$\text{C}5-\text{H}5 \cdots \text{O}3^{\text{ii}}$	0.93	2.43	3.340 (7)	165
$\text{C}19-\text{H}19 \cdots \text{O}7^{\text{iii}}$	0.93	2.46	3.266 (7)	145

Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, y, -z - \frac{1}{2}$; (iii) $-x + 1, y, -z + \frac{1}{2}$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL

We acknowledge the financial support of the National Natural Science Foundation of China (grant No. 20671048).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZZ166).

References

- Armstrong, C. M., Bernhardt, P. V., Chin, P. & Richardson, D. R. (2003). *Eur. J. Inorg. Chem.* pp. 1145–1156.
 Bai, Y., Dang, D. B., Cao, X., Duan, C. Y. & Meng, Q. J. (2006). *Inorg. Chem. Commun.* pp. 86–89.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Siemens (1996). SMART and SAINT. Siemens Analytical X-Ray Systems, Inc., Madison, Wisconsin, USA.

supplementary materials

Acta Cryst. (2009). E65, m520 [doi:10.1107/S1600536809012653]

***catena*-Poly[[[pyridinecopper(II)]-(μ -2-oxidonaphthalene-1-carbaldehyde picolinoylhydrazonato)-[pyridinecopper(II)]- μ -sulfato] diethyl ether hemisolvate]**

Q. Wang, D. Li and D. Wang

Comment

Hydrazone complexes play an important role in the fields of photoelectric materials and medicines due to their biological and pharmacological activities (Armstrong *et al.*, 2003). The molecular structure of the related salicylaldehyde-2-pyridine-carboxyl-hydrazone has been reported (Bai *et al.*, 2006). To throw further light on the coordination characteristics of 2-pyridinecarboxyl-hydrazone and to explore the properties of their complexes, we report the structure of the title complex (I).

The structure of repeating unit of complex I is shown in Fig. 1 and the one-dimensional polymeric chain structure of the complex is shown in Fig. 2. In the complex, each Cu ion is located in a distorted square pyramidal coordination environment. Cu1 is coordinated to two N and three O atoms, while Cu2 links to two O and three N atoms. The Cu—O (2-naphthol) distance [1.888 (3) Å] is slightly shorter than the previously reported Cu—O (phenol) distance [1.954 (3) Å] (Bai *et al.*, 2006), whereas the Cu—O (carbozone) distance [1.964 (3) Å] is longer than the related Cu—O (carbozone) distance of 1.942 (3) Å in the related complex (Bai *et al.*, 2006). In the crystal, weak intermolecular C—H \cdots O interactions connect the chains into a two-dimensional net structure.

Experimental

The title compound was synthesized by mixing 2-hydroxy-1-naphthylaldehyde-2-pyridinecarboxyl-hydrazone (0.0291 g, 0.1 mmol) and copper sulfonate (0.0319 g, 0.2 mmol) and stirring in 10 ml of pyridine for 6 h. The product was filtered and then layered with ether. 2 weeks later brown single crystals were obtained. Anal. Calcd (%) for 2(C₂₇ H₂₁ Cu₂ N₅ O₆ S), C₄ H₁₀ O (Mr = 1415.44): C:49.22; H:3.70; N:9.90; Found (%): C: 49.30; H: 3.52; N: 9.76

Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms, with C—H 0.97 (methylene), C—H 0.93 (pyridine) C—H 0.93 (naphthalene) Å [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$], and with C—H 0.96 Å (methyl) [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$].

Figures




Fig. 1. The molecular structure of the compound, showing 30% probability displacement ellipsoids. Unlabelled atoms are related to the labelled ones by symmetry operation (+x, 2 - y, -1/2 + z). H atoms have been omitted for clarity.




Fig. 2. The one-dimensional polymeric structure of the title complex.

catena-Poly[[[pyridinecopper(II)]-(μ-2-oxidonaphthalene-1-carbaldehyde picolinoylhydrazonato)-[pyridinecopper(II)]-μ-sulfonato] diethyl ether hemisolvate]

Crystal data

[Cu₂(C₁₇H₁₁N₃O)(SO₄)(C₅H₅N)₂]₂·0.5C₄H₁₀O

M_r = 707.69

Monoclinic, *C2/c*

Hall symbol: -C 2yc

a = 26.484 (2) Å

b = 14.0374 (15) Å

c = 16.8083 (17) Å

β = 108.404 (2)°

V = 5929.2 (10) Å³

Z = 8

*F*₀₀₀ = 2888

D_x = 1.586 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 2858 reflections

θ = 2.3–25.3°

μ = 1.56 mm⁻¹

T = 298 K

Block, brown

0.38 × 0.32 × 0.16 mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298 K

phi and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

*T*_{min} = 0.589, *T*_{max} = 0.789

14633 measured reflections

5215 independent reflections

3268 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.055

θ_{max} = 25.0°

θ_{min} = 1.6°

h = -31→31

k = -16→16

l = -12→19

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H-atom parameters constrained
$wR(F^2) = 0.130$	$w = 1/[\sigma^2(F_o^2) + (0.052P)^2 + 19.1978P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
5215 reflections	$(\Delta/\sigma)_{\max} = 0.001$
394 parameters	$\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.19607 (2)	0.78202 (4)	0.14330 (4)	0.0327 (2)
Cu2	0.14395 (2)	1.00409 (4)	-0.08707 (4)	0.03123 (19)
N1	0.16100 (16)	0.9077 (3)	0.0066 (2)	0.0299 (10)
N2	0.20801 (16)	0.8671 (3)	0.0587 (3)	0.0297 (10)
N3	0.06961 (16)	0.9731 (3)	-0.0874 (3)	0.0331 (11)
N4	0.17578 (17)	0.6701 (3)	0.2012 (3)	0.0347 (11)
N5	0.16690 (18)	1.1384 (3)	-0.0190 (3)	0.0368 (11)
O1	0.12165 (13)	0.8167 (3)	0.0856 (2)	0.0366 (9)
O2	0.26870 (13)	0.7475 (3)	0.1794 (2)	0.0374 (9)
O3	0.12067 (13)	1.0678 (2)	-0.2000 (2)	0.0355 (9)
O4	0.20303 (13)	0.9942 (2)	-0.1402 (2)	0.0352 (9)
O5	0.19783 (14)	1.1264 (3)	-0.2360 (2)	0.0406 (9)
O6	0.15828 (16)	0.9758 (3)	-0.2900 (2)	0.0449 (10)
O7	1.0000	0.6854 (6)	0.2500	0.092 (3)
S1	0.17080 (5)	1.04166 (9)	-0.22080 (8)	0.0293 (3)
C1	0.1195 (2)	0.8746 (4)	0.0264 (3)	0.0307 (12)
C2	0.0669 (2)	0.9078 (4)	-0.0292 (3)	0.0326 (12)
C3	0.0197 (2)	0.8731 (4)	-0.0246 (4)	0.0456 (15)

supplementary materials

H3	0.0191	0.8286	0.0161	0.055*
C4	-0.0273 (2)	0.9056 (5)	-0.0820 (4)	0.0550 (17)
H4	-0.0600	0.8828	-0.0808	0.066*
C5	-0.0247 (2)	0.9719 (5)	-0.1405 (4)	0.0532 (17)
H5	-0.0557	0.9948	-0.1794	0.064*
C6	0.0243 (2)	1.0040 (4)	-0.1408 (4)	0.0442 (15)
H6	0.0257	1.0494	-0.1803	0.053*
C7	0.2514 (2)	0.8798 (3)	0.0399 (3)	0.0313 (12)
H7	0.2495	0.9184	-0.0059	0.038*
C8	0.3020 (2)	0.8388 (3)	0.0846 (3)	0.0297 (12)
C9	0.3072 (2)	0.7729 (4)	0.1518 (3)	0.0321 (12)
C10	0.3584 (2)	0.7317 (4)	0.1922 (4)	0.0411 (14)
H10	0.3616	0.6859	0.2335	0.049*
C11	0.4018 (2)	0.7567 (4)	0.1727 (4)	0.0439 (15)
H11	0.4345	0.7299	0.2024	0.053*
C12	0.3994 (2)	0.8238 (4)	0.1071 (4)	0.0396 (14)
C13	0.3491 (2)	0.8645 (4)	0.0621 (3)	0.0332 (13)
C14	0.3484 (2)	0.9312 (4)	-0.0009 (4)	0.0464 (15)
H14	0.3162	0.9585	-0.0319	0.056*
C15	0.3939 (3)	0.9568 (5)	-0.0177 (4)	0.0624 (19)
H15	0.3920	1.0009	-0.0598	0.075*
C16	0.4427 (3)	0.9177 (5)	0.0272 (5)	0.067 (2)
H16	0.4734	0.9358	0.0156	0.080*
C17	0.4451 (2)	0.8525 (5)	0.0886 (4)	0.0556 (18)
H17	0.4779	0.8265	0.1188	0.067*
C18	0.1284 (2)	0.6663 (4)	0.2132 (4)	0.0499 (16)
H18	0.1068	0.7200	0.2012	0.060*
C19	0.1099 (3)	0.5857 (5)	0.2430 (4)	0.0609 (19)
H19	0.0768	0.5859	0.2512	0.073*
C20	0.1414 (3)	0.5058 (5)	0.2601 (4)	0.0580 (18)
H20	0.1294	0.4502	0.2785	0.070*
C21	0.1905 (3)	0.5093 (4)	0.2498 (4)	0.0531 (17)
H21	0.2129	0.4567	0.2628	0.064*
C22	0.2066 (2)	0.5922 (4)	0.2198 (3)	0.0404 (14)
H22	0.2399	0.5938	0.2124	0.048*
C23	0.1707 (2)	1.2197 (4)	-0.0595 (4)	0.0447 (15)
H23	0.1622	1.2175	-0.1176	0.054*
C24	0.1862 (3)	1.3053 (4)	-0.0203 (4)	0.0517 (17)
H24	0.1877	1.3598	-0.0510	0.062*
C25	0.1995 (3)	1.3085 (5)	0.0656 (4)	0.0590 (18)
H25	0.2109	1.3654	0.0939	0.071*
C26	0.1959 (3)	1.2281 (5)	0.1093 (4)	0.0578 (18)
H26	0.2042	1.2295	0.1673	0.069*
C27	0.1797 (2)	1.1443 (4)	0.0646 (4)	0.0473 (16)
H27	0.1776	1.0894	0.0943	0.057*
C28	0.9808 (4)	0.7425 (7)	0.1773 (8)	0.109 (4)
H28A	1.0083	0.7862	0.1733	0.131*
H28B	0.9504	0.7795	0.1798	0.131*
C29	0.9644 (4)	0.6756 (9)	0.1002 (7)	0.136 (4)

H29A	0.9954	0.6439	0.0952	0.204*
H29B	0.9481	0.7122	0.0505	0.204*
H29C	0.9396	0.6291	0.1072	0.204*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0341 (4)	0.0334 (4)	0.0307 (4)	0.0030 (3)	0.0104 (3)	0.0091 (3)
Cu2	0.0336 (4)	0.0347 (4)	0.0247 (3)	0.0023 (3)	0.0083 (3)	0.0051 (3)
N1	0.028 (2)	0.034 (2)	0.024 (2)	0.005 (2)	0.002 (2)	0.007 (2)
N2	0.032 (2)	0.028 (2)	0.026 (2)	0.002 (2)	0.004 (2)	0.0031 (19)
N3	0.029 (2)	0.038 (3)	0.029 (2)	0.004 (2)	0.004 (2)	0.001 (2)
N4	0.036 (3)	0.036 (3)	0.033 (3)	0.008 (2)	0.012 (2)	0.005 (2)
N5	0.049 (3)	0.037 (3)	0.025 (2)	-0.004 (2)	0.012 (2)	-0.002 (2)
O1	0.033 (2)	0.044 (2)	0.033 (2)	0.0032 (17)	0.0119 (18)	0.0116 (18)
O2	0.036 (2)	0.043 (2)	0.034 (2)	0.0059 (18)	0.0127 (18)	0.0159 (18)
O3	0.034 (2)	0.044 (2)	0.0280 (19)	0.0051 (17)	0.0087 (17)	0.0052 (17)
O4	0.034 (2)	0.045 (2)	0.0258 (19)	0.0071 (17)	0.0084 (17)	0.0092 (17)
O5	0.050 (2)	0.039 (2)	0.033 (2)	-0.0119 (19)	0.0138 (19)	0.0061 (18)
O6	0.061 (3)	0.044 (2)	0.034 (2)	-0.006 (2)	0.020 (2)	-0.0134 (18)
O7	0.071 (5)	0.065 (5)	0.150 (8)	0.000	0.047 (6)	0.000
S1	0.0337 (7)	0.0319 (7)	0.0224 (7)	-0.0006 (6)	0.0089 (6)	0.0004 (6)
C1	0.033 (3)	0.033 (3)	0.026 (3)	0.000 (2)	0.008 (2)	-0.002 (2)
C2	0.030 (3)	0.037 (3)	0.030 (3)	0.000 (2)	0.009 (2)	0.002 (2)
C3	0.039 (3)	0.053 (4)	0.046 (4)	0.004 (3)	0.016 (3)	0.011 (3)
C4	0.030 (3)	0.071 (5)	0.064 (4)	-0.004 (3)	0.016 (3)	0.006 (4)
C5	0.033 (3)	0.079 (5)	0.044 (4)	0.009 (3)	0.007 (3)	0.010 (4)
C6	0.040 (4)	0.053 (4)	0.036 (3)	0.008 (3)	0.007 (3)	0.010 (3)
C7	0.038 (3)	0.028 (3)	0.029 (3)	-0.003 (2)	0.011 (3)	0.000 (2)
C8	0.032 (3)	0.027 (3)	0.029 (3)	0.000 (2)	0.009 (2)	-0.002 (2)
C9	0.036 (3)	0.029 (3)	0.032 (3)	0.003 (2)	0.012 (3)	-0.002 (2)
C10	0.034 (3)	0.040 (3)	0.045 (3)	0.006 (3)	0.006 (3)	0.008 (3)
C11	0.031 (3)	0.041 (3)	0.055 (4)	0.008 (3)	0.006 (3)	0.004 (3)
C12	0.032 (3)	0.035 (3)	0.054 (4)	0.003 (3)	0.016 (3)	-0.007 (3)
C13	0.031 (3)	0.034 (3)	0.037 (3)	-0.004 (2)	0.015 (3)	-0.007 (3)
C14	0.043 (4)	0.053 (4)	0.047 (4)	-0.005 (3)	0.019 (3)	0.006 (3)
C15	0.065 (5)	0.069 (5)	0.063 (5)	-0.011 (4)	0.034 (4)	0.010 (4)
C16	0.054 (5)	0.071 (5)	0.085 (6)	-0.009 (4)	0.037 (4)	0.003 (4)
C17	0.043 (4)	0.057 (4)	0.070 (5)	0.003 (3)	0.022 (4)	-0.002 (4)
C18	0.050 (4)	0.047 (4)	0.060 (4)	0.021 (3)	0.028 (3)	0.021 (3)
C19	0.052 (4)	0.064 (5)	0.080 (5)	0.008 (4)	0.040 (4)	0.027 (4)
C20	0.059 (4)	0.055 (4)	0.067 (5)	0.004 (4)	0.028 (4)	0.032 (4)
C21	0.057 (4)	0.041 (4)	0.066 (4)	0.011 (3)	0.024 (4)	0.023 (3)
C22	0.043 (3)	0.038 (3)	0.044 (4)	0.007 (3)	0.018 (3)	0.008 (3)
C23	0.060 (4)	0.044 (4)	0.027 (3)	-0.003 (3)	0.010 (3)	0.000 (3)
C24	0.066 (4)	0.036 (3)	0.048 (4)	-0.004 (3)	0.011 (3)	-0.002 (3)
C25	0.068 (5)	0.047 (4)	0.055 (4)	-0.004 (3)	0.010 (4)	-0.016 (3)
C26	0.072 (5)	0.066 (5)	0.030 (3)	-0.006 (4)	0.009 (3)	-0.016 (3)

supplementary materials

C27	0.064 (4)	0.047 (4)	0.033 (3)	-0.002 (3)	0.017 (3)	0.007 (3)
C28	0.076 (6)	0.081 (7)	0.184 (12)	0.020 (5)	0.058 (7)	0.039 (7)
C29	0.115 (9)	0.170 (12)	0.139 (10)	0.002 (8)	0.063 (8)	0.014 (10)

Geometric parameters (Å, °)

Cu1—O2	1.888 (3)	C8—C13	1.459 (7)
Cu1—N2	1.957 (4)	C9—C10	1.432 (7)
Cu1—O1	1.964 (3)	C10—C11	1.339 (7)
Cu1—N4	2.008 (4)	C10—H10	0.9300
Cu1—O5 ⁱ	2.389 (3)	C11—C12	1.436 (8)
Cu2—O3	2.011 (3)	C11—H11	0.9300
Cu2—N3	2.015 (4)	C12—C17	1.402 (8)
Cu2—N1	2.016 (4)	C12—C13	1.426 (7)
Cu2—O4	2.036 (3)	C13—C14	1.409 (8)
Cu2—N5	2.189 (4)	C14—C15	1.368 (8)
N1—C1	1.328 (6)	C14—H14	0.9300
N1—N2	1.398 (5)	C15—C16	1.388 (9)
N2—C7	1.296 (6)	C15—H15	0.9300
N3—C6	1.325 (6)	C16—C17	1.365 (9)
N3—C2	1.359 (6)	C16—H16	0.9300
N4—C18	1.333 (7)	C17—H17	0.9300
N4—C22	1.341 (6)	C18—C19	1.387 (8)
N5—C27	1.340 (7)	C18—H18	0.9300
N5—C23	1.349 (7)	C19—C20	1.372 (8)
O1—C1	1.273 (6)	C19—H19	0.9300
O2—C9	1.297 (6)	C20—C21	1.367 (8)
O3—S1	1.520 (4)	C20—H20	0.9300
O4—S1	1.510 (3)	C21—C22	1.387 (7)
O5—S1	1.452 (4)	C21—H21	0.9300
O5—Cu1 ⁱⁱ	2.389 (3)	C22—H22	0.9300
O6—S1	1.440 (4)	C23—C24	1.371 (8)
O7—C28 ⁱⁱⁱ	1.416 (10)	C23—H23	0.9300
O7—C28	1.416 (10)	C24—C25	1.374 (9)
C1—C2	1.486 (7)	C24—H24	0.9300
C2—C3	1.366 (7)	C25—C26	1.366 (9)
C3—C4	1.391 (8)	C25—H25	0.9300
C3—H3	0.9300	C26—C27	1.389 (8)
C4—C5	1.371 (8)	C26—H26	0.9300
C4—H4	0.9300	C27—H27	0.9300
C5—C6	1.376 (8)	C28—C29	1.547 (13)
C5—H5	0.9300	C28—H28A	0.9700
C6—H6	0.9300	C28—H28B	0.9700
C7—C8	1.434 (7)	C29—H29A	0.9600
C7—H7	0.9300	C29—H29B	0.9600
C8—C9	1.433 (7)	C29—H29C	0.9600
O2—Cu1—N2	90.44 (16)	C7—C8—C13	120.0 (5)
O2—Cu1—O1	169.76 (15)	O2—C9—C10	117.0 (5)

N2—Cu1—O1	81.79 (16)	O2—C9—C8	124.6 (5)
O2—Cu1—N4	92.59 (16)	C10—C9—C8	118.5 (5)
N2—Cu1—N4	163.41 (17)	C11—C10—C9	122.4 (5)
O1—Cu1—N4	93.03 (16)	C11—C10—H10	118.8
O2—Cu1—O5 ⁱ	96.92 (14)	C9—C10—H10	118.8
N2—Cu1—O5 ⁱ	108.92 (15)	C10—C11—C12	121.8 (5)
O1—Cu1—O5 ⁱ	91.92 (14)	C10—C11—H11	119.1
N4—Cu1—O5 ⁱ	86.91 (16)	C12—C11—H11	119.1
O3—Cu2—N3	94.91 (16)	C17—C12—C13	119.7 (6)
O3—Cu2—N1	164.13 (16)	C17—C12—C11	121.7 (5)
N3—Cu2—N1	80.49 (17)	C13—C12—C11	118.6 (5)
O3—Cu2—O4	70.46 (13)	C14—C13—C12	117.0 (5)
N3—Cu2—O4	150.44 (16)	C14—C13—C8	123.3 (5)
N1—Cu2—O4	106.43 (15)	C12—C13—C8	119.7 (5)
O3—Cu2—N5	93.57 (15)	C15—C14—C13	121.7 (6)
N3—Cu2—N5	107.27 (17)	C15—C14—H14	119.1
N1—Cu2—N5	102.30 (16)	C13—C14—H14	119.1
O4—Cu2—N5	99.37 (16)	C14—C15—C16	120.9 (6)
C1—N1—N2	110.1 (4)	C14—C15—H15	119.6
C1—N1—Cu2	115.5 (3)	C16—C15—H15	119.6
N2—N1—Cu2	134.3 (3)	C17—C16—C15	119.3 (6)
C7—N2—N1	118.2 (4)	C17—C16—H16	120.3
C7—N2—Cu1	128.3 (4)	C15—C16—H16	120.3
N1—N2—Cu1	112.8 (3)	C16—C17—C12	121.4 (6)
C6—N3—C2	117.8 (5)	C16—C17—H17	119.3
C6—N3—Cu2	127.3 (4)	C12—C17—H17	119.3
C2—N3—Cu2	114.7 (3)	N4—C18—C19	123.0 (5)
C18—N4—C22	117.6 (5)	N4—C18—H18	118.5
C18—N4—Cu1	120.9 (4)	C19—C18—H18	118.5
C22—N4—Cu1	121.0 (4)	C20—C19—C18	118.8 (6)
C27—N5—C23	116.2 (5)	C20—C19—H19	120.6
C27—N5—Cu2	122.6 (4)	C18—C19—H19	120.6
C23—N5—Cu2	121.1 (3)	C21—C20—C19	118.8 (6)
C1—O1—Cu1	109.5 (3)	C21—C20—H20	120.6
C9—O2—Cu1	130.7 (3)	C19—C20—H20	120.6
S1—O3—Cu2	94.59 (17)	C20—C21—C22	119.4 (6)
S1—O4—Cu2	93.93 (17)	C20—C21—H21	120.3
S1—O5—Cu1 ⁱⁱ	135.4 (2)	C22—C21—H21	120.3
C28 ⁱⁱⁱ —O7—C28	110.9 (11)	N4—C22—C21	122.3 (5)
O6—S1—O5	112.6 (2)	N4—C22—H22	118.8
O6—S1—O4	111.3 (2)	C21—C22—H22	118.8
O5—S1—O4	110.4 (2)	N5—C23—C24	123.9 (5)
O6—S1—O3	110.2 (2)	N5—C23—H23	118.0
O5—S1—O3	111.0 (2)	C24—C23—H23	118.0
O4—S1—O3	100.8 (2)	C23—C24—C25	118.2 (6)
O1—C1—N1	125.7 (5)	C23—C24—H24	120.9
O1—C1—C2	119.5 (5)	C25—C24—H24	120.9
N1—C1—C2	114.8 (5)	C26—C25—C24	120.0 (6)

supplementary materials

N3—C2—C3	122.6 (5)	C26—C25—H25	120.0
N3—C2—C1	114.3 (4)	C24—C25—H25	120.0
C3—C2—C1	123.1 (5)	C25—C26—C27	118.1 (6)
C2—C3—C4	118.6 (6)	C25—C26—H26	121.0
C2—C3—H3	120.7	C27—C26—H26	121.0
C4—C3—H3	120.7	N5—C27—C26	123.6 (6)
C5—C4—C3	118.9 (6)	N5—C27—H27	118.2
C5—C4—H4	120.5	C26—C27—H27	118.2
C3—C4—H4	120.5	O7—C28—C29	107.9 (8)
C4—C5—C6	119.1 (6)	O7—C28—H28A	110.1
C4—C5—H5	120.5	C29—C28—H28A	110.1
C6—C5—H5	120.5	O7—C28—H28B	110.1
N3—C6—C5	123.0 (6)	C29—C28—H28B	110.1
N3—C6—H6	118.5	H28A—C28—H28B	108.4
C5—C6—H6	118.5	C28—C29—H29A	109.5
N2—C7—C8	124.8 (5)	C28—C29—H29B	109.5
N2—C7—H7	117.6	H29A—C29—H29B	109.5
C8—C7—H7	117.6	C28—C29—H29C	109.5
C9—C8—C7	120.9 (5)	H29A—C29—H29C	109.5
C9—C8—C13	119.1 (5)	H29B—C29—H29C	109.5

Symmetry codes: (i) $x, -y+2, z+1/2$; (ii) $x, -y+2, z-1/2$; (iii) $-x+2, y, -z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C26—H26 \cdots O2 ^{iv}	0.93	2.46	3.386 (7)	174
C5—H5 \cdots O3 ^v	0.93	2.43	3.340 (7)	165
C19—H19 \cdots O7 ^{vi}	0.93	2.46	3.266 (7)	145

Symmetry codes: (iv) $-x+1/2, y+1/2, -z+1/2$; (v) $-x, y, -z-1/2$; (vi) $-x+1, y, -z+1/2$.

Fig. 1




Fig. 2

