

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Amino-2-methyl-4-oxo-3,4-dihydroquinazolin-1-ium *p*-toluenesulfonate monohydrate

 Mohammad Arfan,^a M. Nawaz Tahir,^{b*} Rasool Khan^a and Mohammad S. Iqbal^c
^aUniversity of Peshawar, Institute of Chemical Sciences, Peshawar 25120, Pakistan,

^bUniversity of Sargodha, Department of Physics, Sargodha, Pakistan, and ^cGovernment College University, Department of Chemistry, Lahore, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 13 April 2009; accepted 13 April 2009

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.034; wR factor = 0.091; data-to-parameter ratio = 18.1.

In the title hydrated molecular salt, $\text{C}_9\text{H}_{10}\text{N}_3\text{O}^{+}\cdot\text{C}_7\text{H}_7\text{O}_3\text{S}^{-}\cdot\text{H}_2\text{O}$, the cation is protonated at a quinazolinone N atom and an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond occurs. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\pi$ and weak aromatic $\pi-\pi$ stacking interactions [centroid-centroid separations = 3.8648 (12) and 3.9306 (13) Å] help to establish the packing; a short $\text{S}=\text{O}\cdots\pi$ contact is also seen.

Related literature

For a related structure, see: Atkinson & Meades (2000). For background on the properties of cyclic amidines and quinazolinones, see: Glaser & Traber (1984); Haverá (1979); Hori *et al.* (1990); Liverton *et al.* (1998). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_9\text{H}_{10}\text{N}_3\text{O}^{+}\cdot\text{C}_7\text{H}_7\text{O}_3\text{S}^{-}\cdot\text{H}_2\text{O}$
 $M_r = 365.40$

 Monoclinic, Cc
 $a = 20.838$ (1) Å

 $b = 6.2769$ (3) Å

 $c = 14.7897$ (7) Å

 $\beta = 116.676$ (1)°

 $V = 1728.56$ (14) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.22$ mm⁻¹
 $T = 296$ K

 $0.28 \times 0.24 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

 $T_{\min} = 0.935$, $T_{\max} = 0.958$

9546 measured reflections

4390 independent reflections

 3952 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.091$
 $S = 1.00$

4390 reflections

243 parameters

2 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Absolute structure: Flack (1983),

2109 Friedel pairs

Flack parameter: 0.04 (5)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O4}^i$	0.89 (3)	1.89 (3)	2.734 (3)	158 (3)
$\text{N3}-\text{H3A}\cdots\text{O1}$	0.83 (3)	2.31 (4)	2.693 (3)	109 (3)
$\text{N3}-\text{H3B}\cdots\text{O3}^{ii}$	0.86 (3)	2.22 (3)	2.963 (3)	146 (3)
$\text{O5}-\text{H5A}\cdots\text{O3}^i$	0.73 (5)	2.16 (5)	2.871 (3)	166 (7)
$\text{O5}-\text{H5B}\cdots\text{O2}$	0.82 (6)	2.07 (6)	2.862 (4)	162 (5)
$\text{C4}-\text{H4}\cdots\text{O5}^{iii}$	0.93	2.58	3.423 (5)	152
$\text{C9}-\text{H9C}\cdots\text{O4}^i$	0.96	2.43	3.251 (3)	144
$\text{C2}-\text{H2}\cdots\text{CgC}^c$	0.93	2.84	3.533 (2)	132
$\text{S1}-\text{O2}\cdots\text{CgB}$	1.44 (1)	3.17 (1)	3.8430 (9)	107 (1)

 Symmetry codes: (i) $x, y+1, z$; (ii) $x, -y+1, z+\frac{1}{2}$; (iii) $x-\frac{1}{2}, y-\frac{1}{2}, z$. CgB and CgC are the centroids of the C1/C6/C7/N2/C8/N1 and C10-C15 rings.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

The authors acknowledge the Higher Education Commission, Islamabad, Pakistan, for funding the purchase of the diffractometer at GCU, Lahore.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2950).

References

- Atkinson, R. S. & Meades, C. K. (2000). *Tetrahedron Lett.* **41**, 7769–7772.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Glaser, T. & Traber, J. (1984). *Inflamm. Res.* **15**, 341–348.
- Haverá, H. J. (1979). *J. Med. Chem.* **22**, 1548–1550.
- Hori, M., Iemura, R., Hara, H., Ozaki, A., Sukamoto, T. & Ohtaka, H. (1990). *Chem. Pharm. Bull.* **38**, 681–687.
- Liverton, N. J., Armstrong, D. J., Claremon, D. A., Remy, D. C., Bardwin, J. J., Lynch, R. J., Zhang, G. & Gould, R. J. (1998). *Bioorg. Med. Chem. Lett.* **8**, 483–486.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o1051 [doi:10.1107/S1600536809013841]

3-Amino-2-methyl-4-oxo-3,4-dihydroquinazolin-1-ium *p*-toluenesulfonate monohydrate

M. Arfan, M. N. Tahir, R. Khan and M. S. Iqbal

Comment

Cyclic amidines and quinazolinones, are known to possess diverse pharmacological activities as phosphodiesterase inhibitors (Glaser & Traber, 1984), anticonvulsants (Hori *et al.*, 1990), antihypertensives (Glaser & Traber, 1984), vasodilators (Havera, 1979) and fibrinogen receptor antagonists (Liverton *et al.*, 1998).

We now report synthesis and structure of the title compound (I), (Fig. 1), through a reaction of 2-aminobenzoic acid and hydrazine in presence of *p*-toluenesulfonic acid. A one pot, three component, *p*-toluenesulfonic acid catalyzed heterocyclization has yielded colourless prisms of (I) in the form of a *p*-toluenesulfonate salt with one water molecule of crystallization.

The crystal structure of (II) 3-(6-Azabicyclo(3.1.0)hex-2-en-6-yl)-2-((S)-1-hydroxy-2,2-dimethylpropyl)quinazolin-4(3*H*)-one (Atkinson & Meades, 2000) has been published. The title compound has also quinazoline with a chemically different attachments.

In the title compound the two fused rings A (C1—C6) and B(C1/C6/C7/N2/C8/N1) are essentially planar and the ring C (C10—C15) of the *p*-toluenesulfonate anion is of course planar. The title compound is stabilized due to intra- and intermolecular H-bonding as well as C—H \cdots π and S1=O2 \cdots CgB interactions (Table 1). There also exist interactions between the centroids CgA—CgCⁱ [symmetry code: $i = x, -y + 1, z + 1/2$] and CgB—CgCⁱ at a distance of 3.8648 (12) and 3.9306 (13) Å, respectively. The water molecule connects the *p*-toluenesulfonate ions only. In the title compound there exist $R_1^1(5)$ and $R_2^1(6)$ ring motifs (Bernstein *et al.*, 1995), (Fig 2).

Experimental

A mixture of anthranilic acid (0.14 g, 1 mmol), triethylorthoacetate (0.23 ml, 1.2 mmol), hydrazine hydrate (0.1 ml) and *p*-toluenesulfonic acid (1 g, 5 mmol) was stirred at room temperature for 1 h. After completion of the reaction as indicated by TLC, the reaction mixture was poured into water and allowed to settle, the product precipitated as colourless prisms of (I). The product was filtered, washed with water and dried. m.p. 585–594 K. yield: 72%.

Refinement

The coordinates of H-atoms connected with water molecule and NH₂ group were refined. C-bound H-atoms were positioned geometrically (C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C, N, O})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. The molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level. H-atoms are shown by small spheres of arbitrary radius and hydrogen bonds by dotted lines.




Fig. 2. A fragment of the packing of (I) showing $R_1^1(5)$ and $R_2^1(6)$ ring motifs.

(I)

Crystal data

$C_9H_{10}N_3O^+ \cdot C_7H_7O_3S^- \cdot H_2O$

$M_r = 365.40$

Monoclinic, Cc

Hall symbol: $C -2yc$

$a = 20.838$ (1) Å

$b = 6.2769$ (3) Å

$c = 14.7897$ (7) Å

$\beta = 116.676$ (1)°

$V = 1728.56$ (14) Å³

$Z = 4$

$F_{000} = 768$

$D_x = 1.404$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 4390 reflections

$\theta = 2.9$ – 28.9 °

$\mu = 0.22$ mm⁻¹

$T = 296$ K

Prism, colourless

$0.28 \times 0.24 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 7.30 pixels mm⁻¹

$T = 296$ K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2005)

$T_{\min} = 0.935$, $T_{\max} = 0.958$

9546 measured reflections

4390 independent reflections

3952 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\text{max}} = 28.9$ °

$\theta_{\text{min}} = 2.9$ °

$h = -28 \rightarrow 28$

$k = -8 \rightarrow 6$

$l = -19 \rightarrow 20$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.034$	$w = 1/[\sigma^2(F_o^2) + (0.0508P)^2 + 0.3201P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.091$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.00$	$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
4390 reflections	$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$
243 parameters	Extinction correction: none
2 restraints	Absolute structure: Flack (1983), 2109 Friedal pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.04 (5)
Secondary atom site location: difference Fourier map	

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.12729 (10)	0.3093 (3)	0.49282 (14)	0.0676 (6)
N1	0.15599 (8)	0.8870 (3)	0.40550 (11)	0.0404 (4)
N2	0.19969 (9)	0.5856 (3)	0.49997 (11)	0.0442 (5)
N3	0.26093 (12)	0.4718 (4)	0.56765 (18)	0.0611 (7)
C1	0.08550 (9)	0.8146 (3)	0.36387 (13)	0.0421 (5)
C2	0.02990 (11)	0.9399 (4)	0.29579 (15)	0.0534 (6)
C3	-0.03899 (12)	0.8605 (5)	0.25562 (17)	0.0676 (8)
C4	-0.05286 (12)	0.6617 (5)	0.28300 (18)	0.0723 (9)
C5	0.00197 (13)	0.5380 (4)	0.34994 (17)	0.0607 (8)
C6	0.07242 (10)	0.6139 (3)	0.39165 (14)	0.0458 (6)
C7	0.13149 (12)	0.4861 (3)	0.46325 (15)	0.0483 (7)
C8	0.21066 (10)	0.7761 (3)	0.46961 (13)	0.0407 (5)
C9	0.28411 (11)	0.8657 (4)	0.50903 (18)	0.0573 (6)
S1	0.19585 (2)	0.35554 (7)	0.26006 (3)	0.0413 (1)
O2	0.19879 (10)	0.5841 (2)	0.27106 (14)	0.0654 (5)
O3	0.24598 (7)	0.2742 (3)	0.22604 (11)	0.0595 (5)

supplementary materials

O4	0.20275 (8)	0.2499 (3)	0.35157 (10)	0.0551 (4)
C10	0.10901 (9)	0.2884 (3)	0.16552 (13)	0.0390 (5)
C11	0.09715 (10)	0.0902 (3)	0.12006 (14)	0.0443 (5)
C12	0.02813 (11)	0.0315 (3)	0.05242 (15)	0.0504 (6)
C13	-0.02945 (11)	0.1669 (4)	0.02991 (15)	0.0528 (6)
C14	-0.01601 (12)	0.3654 (4)	0.07515 (18)	0.0593 (7)
C15	0.05282 (11)	0.4272 (3)	0.14287 (15)	0.0497 (6)
C16	-0.10500 (13)	0.1024 (6)	-0.0422 (2)	0.0780 (9)
O5	0.30414 (16)	0.8511 (4)	0.2540 (3)	0.1157 (13)
H1	0.1621 (12)	1.020 (4)	0.3909 (17)	0.0485*
H2	0.03897	1.07432	0.27772	0.0641*
H3	-0.07674	0.94197	0.20938	0.0809*
H3A	0.2495 (16)	0.346 (5)	0.551 (2)	0.0734*
H3B	0.2630 (16)	0.493 (5)	0.626 (2)	0.0734*
H4	-0.09983	0.61162	0.25573	0.0867*
H5	-0.00766	0.40379	0.36759	0.0728*
H9A	0.31436	0.76770	0.49599	0.0688*
H9B	0.30317	0.88850	0.58061	0.0688*
H9C	0.28237	0.99873	0.47601	0.0688*
H11	0.13528	-0.00298	0.13486	0.0532*
H12	0.02028	-0.10145	0.02150	0.0605*
H14	-0.05397	0.45945	0.05983	0.0712*
H15	0.06092	0.56133	0.17269	0.0596*
H16A	-0.13244	0.08167	-0.00532	0.0934*
H16B	-0.10387	-0.02800	-0.07543	0.0934*
H16C	-0.12681	0.21226	-0.09184	0.0934*
H5A	0.284 (3)	0.949 (8)	0.248 (4)	0.1388*
H5B	0.271 (3)	0.770 (8)	0.245 (4)	0.1388*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0927 (12)	0.0463 (8)	0.0836 (11)	-0.0072 (9)	0.0571 (10)	0.0044 (8)
N1	0.0439 (7)	0.0344 (7)	0.0429 (7)	-0.0061 (6)	0.0195 (6)	-0.0032 (6)
N2	0.0556 (9)	0.0390 (8)	0.0417 (8)	0.0027 (7)	0.0252 (7)	-0.0019 (6)
N3	0.0707 (12)	0.0523 (12)	0.0608 (12)	0.0167 (10)	0.0299 (10)	0.0081 (9)
C1	0.0436 (8)	0.0496 (10)	0.0373 (8)	-0.0099 (7)	0.0218 (7)	-0.0078 (7)
C2	0.0488 (10)	0.0620 (13)	0.0474 (10)	-0.0027 (9)	0.0197 (8)	0.0049 (9)
C3	0.0440 (10)	0.103 (2)	0.0505 (11)	-0.0070 (11)	0.0166 (9)	0.0046 (12)
C4	0.0484 (11)	0.114 (2)	0.0559 (12)	-0.0311 (13)	0.0247 (10)	-0.0118 (13)
C5	0.0637 (12)	0.0704 (15)	0.0578 (12)	-0.0289 (11)	0.0361 (10)	-0.0144 (10)
C6	0.0545 (10)	0.0497 (11)	0.0427 (9)	-0.0131 (8)	0.0304 (8)	-0.0092 (8)
C7	0.0670 (13)	0.0425 (11)	0.0494 (11)	-0.0065 (9)	0.0387 (10)	-0.0048 (8)
C8	0.0475 (9)	0.0374 (9)	0.0388 (8)	-0.0016 (7)	0.0207 (7)	-0.0063 (7)
C9	0.0430 (9)	0.0518 (12)	0.0664 (12)	-0.0025 (9)	0.0150 (9)	-0.0011 (10)
S1	0.0448 (2)	0.0404 (2)	0.0389 (2)	-0.0109 (2)	0.0191 (2)	-0.0002 (2)
O2	0.0721 (9)	0.0420 (7)	0.0751 (10)	-0.0179 (8)	0.0267 (8)	-0.0047 (8)
O3	0.0470 (7)	0.0789 (11)	0.0566 (8)	-0.0051 (7)	0.0269 (7)	-0.0023 (7)

O4	0.0604 (8)	0.0616 (9)	0.0386 (6)	-0.0227 (7)	0.0180 (6)	0.0002 (6)
C10	0.0436 (8)	0.0402 (9)	0.0353 (8)	-0.0043 (7)	0.0197 (7)	0.0000 (7)
C11	0.0449 (9)	0.0429 (10)	0.0463 (9)	-0.0019 (7)	0.0215 (7)	-0.0053 (7)
C12	0.0517 (10)	0.0508 (11)	0.0472 (10)	-0.0104 (9)	0.0209 (8)	-0.0081 (8)
C13	0.0438 (9)	0.0718 (14)	0.0432 (10)	-0.0085 (9)	0.0198 (8)	0.0012 (9)
C14	0.0447 (10)	0.0708 (15)	0.0591 (13)	0.0161 (10)	0.0204 (9)	0.0053 (10)
C15	0.0572 (11)	0.0434 (10)	0.0492 (10)	0.0058 (9)	0.0246 (9)	-0.0023 (8)
C16	0.0473 (12)	0.114 (2)	0.0647 (16)	-0.0166 (13)	0.0180 (11)	-0.0023 (15)
O5	0.1089 (19)	0.0754 (15)	0.205 (3)	-0.0269 (14)	0.108 (2)	-0.0230 (18)

Geometric parameters (Å, °)

S1—O3	1.4412 (17)	C8—C9	1.482 (3)
S1—O4	1.4551 (16)	C2—H2	0.9300
S1—O2	1.4420 (13)	C3—H3	0.9300
S1—C10	1.7707 (19)	C4—H4	0.9300
O1—C7	1.210 (3)	C5—H5	0.9300
O5—H5B	0.82 (6)	C9—H9C	0.9600
O5—H5A	0.73 (5)	C9—H9B	0.9600
N1—C1	1.390 (3)	C9—H9A	0.9600
N1—C8	1.308 (3)	C10—C11	1.383 (3)
N2—N3	1.412 (3)	C10—C15	1.375 (3)
N2—C7	1.418 (3)	C11—C12	1.383 (3)
N2—C8	1.333 (3)	C12—C13	1.384 (3)
N1—H1	0.89 (3)	C13—C14	1.382 (3)
N3—H3A	0.83 (3)	C13—C16	1.507 (4)
N3—H3B	0.86 (3)	C14—C15	1.386 (3)
C1—C6	1.390 (3)	C11—H11	0.9300
C1—C2	1.388 (3)	C12—H12	0.9300
C2—C3	1.377 (4)	C14—H14	0.9300
C3—C4	1.382 (4)	C15—H15	0.9300
C4—C5	1.369 (4)	C16—H16C	0.9600
C5—C6	1.396 (4)	C16—H16A	0.9600
C6—C7	1.453 (3)	C16—H16B	0.9600
O2—S1—C10	107.61 (11)	C2—C3—H3	119.00
O2—S1—O3	113.02 (12)	C5—C4—H4	120.00
O2—S1—O4	111.53 (11)	C3—C4—H4	120.00
O4—S1—C10	105.38 (9)	C4—C5—H5	120.00
O3—S1—O4	112.26 (10)	C6—C5—H5	120.00
O3—S1—C10	106.50 (9)	C8—C9—H9A	109.00
H5A—O5—H5B	96 (7)	C8—C9—H9B	109.00
C1—N1—C8	123.39 (18)	C8—C9—H9C	109.00
N3—N2—C7	118.69 (19)	H9A—C9—H9B	109.00
N3—N2—C8	117.1 (2)	H9A—C9—H9C	109.00
C7—N2—C8	124.06 (17)	H9B—C9—H9C	110.00
C1—N1—H1	116.1 (17)	S1—C10—C11	119.63 (15)
C8—N1—H1	120.3 (17)	S1—C10—C15	119.88 (15)
N2—N3—H3A	103 (2)	C11—C10—C15	120.33 (18)
H3A—N3—H3B	109 (3)	C10—C11—C12	119.49 (19)

supplementary materials

N2—N3—H3B	105 (2)	C11—C12—C13	121.21 (19)
C2—C1—C6	121.2 (2)	C12—C13—C14	118.2 (2)
N1—C1—C2	120.53 (19)	C14—C13—C16	120.3 (2)
N1—C1—C6	118.30 (17)	C12—C13—C16	121.5 (2)
C1—C2—C3	118.5 (2)	C13—C14—C15	121.4 (2)
C2—C3—C4	121.0 (2)	C10—C15—C14	119.38 (19)
C3—C4—C5	120.5 (3)	C12—C11—H11	120.00
C4—C5—C6	119.8 (2)	C10—C11—H11	120.00
C5—C6—C7	120.83 (19)	C11—C12—H12	119.00
C1—C6—C7	120.12 (19)	C13—C12—H12	119.00
C1—C6—C5	119.05 (19)	C15—C14—H14	119.00
N2—C7—C6	114.30 (17)	C13—C14—H14	119.00
O1—C7—N2	119.2 (2)	C10—C15—H15	120.00
O1—C7—C6	126.5 (2)	C14—C15—H15	120.00
N2—C8—C9	120.72 (19)	C13—C16—H16B	109.00
N1—C8—N2	119.8 (2)	C13—C16—H16C	109.00
N1—C8—C9	119.51 (19)	C13—C16—H16A	109.00
C1—C2—H2	121.00	H16A—C16—H16C	109.00
C3—C2—H2	121.00	H16B—C16—H16C	109.00
C4—C3—H3	120.00	H16A—C16—H16B	109.00
O4—S1—C10—C11	-78.48 (18)	N1—C1—C6—C7	0.6 (3)
O4—S1—C10—C15	96.94 (18)	C2—C1—C6—C5	0.1 (3)
O2—S1—C10—C11	162.40 (17)	C1—C2—C3—C4	0.7 (4)
O2—S1—C10—C15	-22.2 (2)	C2—C3—C4—C5	-0.8 (4)
O3—S1—C10—C11	40.94 (19)	C3—C4—C5—C6	0.6 (4)
O3—S1—C10—C15	-143.64 (17)	C4—C5—C6—C7	179.7 (2)
C1—N1—C8—C9	179.25 (18)	C4—C5—C6—C1	-0.2 (3)
C8—N1—C1—C6	-0.1 (3)	C5—C6—C7—N2	-179.2 (2)
C1—N1—C8—N2	-1.8 (3)	C1—C6—C7—O1	-179.4 (2)
C8—N1—C1—C2	-179.68 (19)	C1—C6—C7—N2	0.7 (3)
N3—N2—C7—O1	2.0 (3)	C5—C6—C7—O1	0.7 (4)
C7—N2—C8—C9	-177.77 (19)	S1—C10—C11—C12	174.65 (16)
N3—N2—C7—C6	-178.09 (19)	C15—C10—C11—C12	-0.8 (3)
C8—N2—C7—O1	177.4 (2)	S1—C10—C15—C14	-174.44 (17)
C8—N2—C7—C6	-2.7 (3)	C11—C10—C15—C14	0.9 (3)
N3—N2—C8—N1	178.78 (19)	C10—C11—C12—C13	-0.5 (3)
N3—N2—C8—C9	-2.3 (3)	C11—C12—C13—C14	1.5 (3)
C7—N2—C8—N1	3.3 (3)	C11—C12—C13—C16	-178.8 (2)
C2—C1—C6—C7	-179.83 (19)	C12—C13—C14—C15	-1.3 (4)
N1—C1—C6—C5	-179.51 (19)	C16—C13—C14—C15	179.0 (2)
N1—C1—C2—C3	179.3 (2)	C13—C14—C15—C10	0.1 (3)
C6—C1—C2—C3	-0.3 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O4 ⁱ	0.89 (3)	1.89 (3)	2.734 (3)	158 (3)
N3—H3A \cdots O1	0.83 (3)	2.31 (4)	2.693 (3)	109 (3)

N3—H3B···O3 ⁱⁱ	0.86 (3)	2.22 (3)	2.963 (3)	146 (3)
O5—H5A···O3 ⁱ	0.73 (5)	2.16 (5)	2.871 (3)	166 (7)
O5—H5B···O2	0.82 (6)	2.07 (6)	2.862 (4)	162 (5)
C4—H4···O5 ⁱⁱⁱ	0.93	2.58	3.423 (5)	152
C9—H9C···O4 ⁱ	0.96	2.43	3.251 (3)	144
C2—H2···CgC ⁱ	0.93	2.84	3.533 (2)	132
S1—O2···CgB	1.4420 (13)	3.169 (2)	3.8430 (9)	106.86 (10)

Symmetry codes: (i) $x, y+1, z$; (ii) $x, -y+1, z+1/2$; (iii) $x-1/2, y-1/2, z$.

Fig. 1




Fig. 2

