

# Bis(2,6-diamino-3,5-dibromopyridinium) hexabromidostannate(IV)

 Rawhi H. Al-Far,<sup>a</sup> Salim F. Haddad<sup>b</sup> and Basem Fares Ali<sup>c\*</sup>
<sup>a</sup>Faculty of Information Technology and Science, Al-Balqa'a Applied University, Salt, Jordan, <sup>b</sup>Department of Chemistry, University of Jordan, Amman, Jordan, and <sup>c</sup>Department of Chemistry, Al al-Bayt University, Mafraq 25113, Jordan

Correspondence e-mail: bfali@aabu.edu.jo


Received 16 April 2009; accepted 23 April 2009

 Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.019$  Å;  $R$  factor = 0.059;  $wR$  factor = 0.146; data-to-parameter ratio = 17.4.

The asymmetric unit of the title compound,  $(\text{C}_5\text{H}_6\text{Br}_2\text{N}_3)_2[\text{SnBr}_6]$ , contains one cation and one half-anion in which the Sn atom is located on a crystallographic centre of inversion and is in a quasi-octahedral geometry. The crystal structure is assembled *via* hydrogen-bonding interactions of two kinds,  $\text{N}(\text{pyridine/amine})-\text{H}\cdots\text{Br}-\text{Sn}$ , along with  $\text{C}-\text{Br}\cdots\text{Br}-\text{Sn}$  interactions [3.4925 (19) Å]. The cations are involved in  $\pi-\pi$  stacking, which adds an extra supramolecularity as it presents a strong case of offset-face-to-face motifs [centroid-centroid distance = 3.577 (3) Å]. The intermolecular hydrogen bonds, short  $\text{Br}\cdots\text{Br}$  interactions and  $\pi-\pi$  stacking result in the formation of a three-dimensional supramolecular architecture.

## Related literature

For general background to hybrid organic-inorganic compounds, see: Aruta *et al.* (2005); Hill (1998); Kagan *et al.* (1999); Knutson *et al.* (2005); Raptopoulou *et al.* (2002). For related structures, see: Al-Far & Ali (2007); Al-Far, Ali & Al-Sou'od (2007); Ali & Al-Far (2007); Ali *et al.* (2008); Ali, Al-Far & Ng (2007); Awwadi *et al.* (2007); Tudela & Khan (1991); Willey *et al.* (1998). For bond-length data, see: Allen *et al.* (1987).



## Experimental

### Crystal data

$(\text{C}_5\text{H}_6\text{Br}_2\text{N}_3)_2[\text{SnBr}_6]$   
 $M_r = 1133.97$   
 Monoclinic,  $P2_1/c$   
 $a = 8.3696$  (14) Å  
 $b = 16.720$  (2) Å  
 $c = 9.5814$  (15) Å  
 $\beta = 112.556$  (12)°

$V = 1238.3$  (3) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 17.18$  mm<sup>-1</sup>  
 $T = 295$  K  
 0.30 × 0.30 × 0.20 mm

### Data collection

Bruker P4 diffractometer  
 Absorption correction:  $\psi$  scan  
 (XSCANS; Bruker, 1996)  
 $T_{\min} = 0.007$ ,  $T_{\max} = 0.035$   
 2825 measured reflections  
 2162 independent reflections

1437 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.078$   
 3 standard reflections  
 every 97 reflections  
 intensity decay: 0.01 %

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$   
 $wR(F^2) = 0.146$   
 $S = 1.00$   
 2162 reflections

124 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.97$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -1.63$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

|         |             |         |             |
|---------|-------------|---------|-------------|
| Sn1—Br1 | 2.6002 (13) | Sn1—Br3 | 2.6131 (14) |
| Sn1—Br2 | 2.5768 (14) |         |             |

 Symmetry code: (i)  $-x + 2, -y + 1, -z + 1$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$                      | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|------------------------------------|-------|-------------|-------------|---------------|
| N1—H1 $\cdots$ Br3 <sup>ii</sup>   | 0.86  | 2.54        | 3.354 (9)   | 159           |
| N2—H2B $\cdots$ Br3 <sup>ii</sup>  | 0.86  | 2.88        | 3.612 (12)  | 144           |
| N3—H3A $\cdots$ Br2 <sup>ii</sup>  | 0.86  | 2.79        | 3.608 (10)  | 160           |
| N3—H3B $\cdots$ Br1 <sup>iii</sup> | 0.86  | 2.82        | 3.604 (10)  | 153           |

 Symmetry codes: (ii)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $-x + 1, -y + 1, -z + 1$ .

Data collection: XSCANS (Bruker, 1996); cell refinement: XSCANS; data reduction: SHELXTL (Sheldrick, 2008); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The University of Jordan, Al-Balqa'a Applied University and Al al-Bayt University are thanked for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2669).

## References

- Al-Far, R. & Ali, B. F. (2007). *Acta Cryst.* **C63**, m137–m139.  
 Ali, B. F. & Al-Far, R. (2007). *Acta Cryst.* **E63**, m892–m894.  
 Ali, B. F., Al-Far, R. & Al-Sou'od, K. (2007). *J. Chem. Crystallogr.* **37**, 265–273.  
 Ali, B. F., Al-Far, R. H. & Haddad, S. F. (2008). *Acta Cryst.* **E64**, m749–m750.  
 Ali, B. F., Al-Far, R. & Ng, S. W. (2007). *Acta Cryst.* **E63**, m2102–m2103.

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Aruta, C., Licci, F., Zappettini, A., Bolzoni, F., Rastelli, F., Ferro, P. & Besagni, T. (2005). *Appl. Phys. A*, **81**, 963–968.
- Awwadi, F. F., Willett, R. D., Peterson, K. A. & Twamley, B. (2007). *J. Phys. Chem. A*, **111**, 2319–2328.
- Bruker (1996). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hill, C. L. (1998). *Chem. Rev.* **98**, 1–2.
- Kagan, C. R., Mitzi, D. B. & Dimitrakopoulos, C. D. (1999). *Science*, **286**, 945–947.
- Knutson, J. L., Martin, J. D. & Mitzi, D. B. (2005). *Inorg. Chem.* **44**, 4699–4705.
- Raptopoulou, C. P., Terzis, A., Mousdis, G. A. & Papavassiliou, G. C. (2002). *Z. Naturforsch. Teil B*, **57**, 645–650.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tudela, D. & Khan, M. A. (1991). *J. Chem. Soc. Dalton Trans.* pp. 1003–1006.
- Wiley, G. R., Woodman, T. J., Somasundaram, U., Aris, D. R. & Errington, W. (1998). *J. Chem. Soc. Dalton Trans.* pp. 2573–2576.

**supplementary materials**

*Acta Cryst.* (2009). E65, m583-m584 [ doi:10.1107/S1600536809015189 ]

## Bis(2,6-diamino-3,5-dibromopyridinium) hexabromidostannate(IV)

R. H. Al-Far, S. F. Haddad and B. F. Ali

### Comment

Non-covalent interactions play an important role in organizing structural units in both natural and artificial systems. Hybrid organic-inorganic compounds are of great interest owing to their ionic, electrical, magnetic and optical properties (Hill, 1998; Kagan *et al.*, 1999; Raptopoulou *et al.*, 2002). Tin metal-halo based hybrids are of particular interest as being materials with interesting optical and magnetic properties (Aruta *et al.*, 2005; Knutson *et al.*, 2005; Kagan *et al.*, 1999). We are currently carrying out studies about lattice including different types of intermolecular interactions (aryl $\cdots$ aryl,  $X\cdots X$ ,  $X\cdots$ aryl and  $X\cdots H$ ). Within our research of hybrid compounds containing tin metal (Al-Far & Ali 2007; Al-Far, Ali & Al-Sou'od, 2007; Ali & Al-Far, 2007; Ali, Al-Far & Ng, 2007), we report herein the crystal structure of the title compound.

The asymmetric unit of the title compound contains one cation and one-half anion, in which the Sn atom is located on a crystallographic centre of inversion and is in a quasi-octahedral geometry (Fig. 1 and Table 1). The Sn-Br bonds are in accordance with the corresponding values in similar compounds (Willey *et al.*, 1998; Tudela & Khan 1991; Ali *et al.*, 2008; Al-Far & Ali 2007). The pyridine ring of the starting cation have undergone bromination during the synthesis process (Al-Far & Ali, 2007). In the cation, the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges.

In the crystal structure, weak intermolecular N-H $\cdots$ Br interactions (Table 2) link the molecules into alternating layers of cations and stacks of anions (Fig. 2), in which they may be effective in the stabilization of the structure. The anion stacks are interacting with the cation layers in an extensive hydrogen bonding and Br $\cdots$ Br halogen bonding interactions. Each anion is surrounded by six cations *via* three H—N—H $\cdots$ Br, one N—H $\cdots$ Br interactions and the symmetry related ones along with one Br $\cdots$ Br interaction [ $Br2\cdots Br4^i = 3.4925(19)$  Å, symmetry code (i): 2 - x, 1 - y, -z] and the symmetry related one. On the other hand, each cation is associated with three anions, through six (N<sub>pyridinic</sub>, N<sub>aminic</sub>)—H $\cdots$ Br—Sn hydrogen bonding interactions, and by one C—Br $\cdots$ Br—Sn interaction. It is noteworthy that structural and theoretical results (Awwadi *et al.* 2007; and references therein), show the significance of linear C—Br $\cdots$ Br synthons in influencing structures of crystalline materials and in use as potential building blocks in crystal engineering *via* supramolecular synthesis.

Moreover, interactions between cations perpendicular to [101] represent a case of strong offset face-to-face  $\pi$ - $\pi$  motif. This is evident by the centroids separation distance of 5.059 (3) Å, with the perpendicular distance between planes being 3.577 (3) Å (the sliding angle between planes is 45.0 (3)°).

### Experimental

For the preparation of the title compound, the warm solution of SnCl<sub>2</sub> metal (1.0 mmol) dissolved in absolute ethanol (15 ml) was mixed with a stirred hot solution of 2,6-diaminopyridine (98%; 2 mmol) dissolved in ethanol (20 ml). The mixture was acidified with HBr (48%, 2-3 ml), and then treated with liquid Br<sub>2</sub> (2-3 ml). The resulting mixture was stirred for 3 h, and then allowed to evaporate at room temperature. The salt crystallized over 2 d, as nice parallelepiped yellow crystals (yield; 82%).

## Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH and NH<sub>2</sub>) and C-H = 0.93 Å for aromatic H, respectively, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ .

## Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level [symmetry code (A): 2 - x, 1 - y, 1 - z].




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds and Br...Br interactions are shown as dashed lines.

## Bis(2,6-diamino-3,5-dibromopyridinium) hexabromidostannate(IV)

### Crystal data

(C<sub>5</sub>H<sub>6</sub>Br<sub>2</sub>N<sub>3</sub>)<sub>2</sub>[SnBr<sub>6</sub>]

$M_r = 1133.97$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.3696$  (14) Å

$b = 16.720$  (2) Å

$c = 9.5814$  (15) Å

$\beta = 112.556$  (12)°

$V = 1238.3$  (3) Å<sup>3</sup>

$Z = 2$

$F_{000} = 1028$

$D_x = 3.042$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 29 reflections

$\theta = 5.7\text{--}12.5^\circ$

$\mu = 17.18$  mm<sup>-1</sup>

$T = 295$  K

Parallelepiped, yellow

$0.30 \times 0.30 \times 0.20$  mm

### Data collection

Bruker P4  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$  K

$\omega$  scans

Absorption correction:  $\psi$  scan  
(*XSCANS*; Bruker, 1996)

$T_{\text{min}} = 0.008$ ,  $T_{\text{max}} = 0.035$

2825 measured reflections

$R_{\text{int}} = 0.078$

$\theta_{\text{max}} = 25.0^\circ$

$\theta_{\text{min}} = 2.4^\circ$

$h = -9 \rightarrow 1$

$k = -19 \rightarrow 1$

$l = -10 \rightarrow 11$

3 standard reflections

every 97 reflections

2162 independent reflections  
 1437 reflections with  $I > 2\sigma(I)$

intensity decay: 0.01%

### Refinement

|  |  |
|--|--|
| Refinement on $F^2$  | Secondary atom site location: difference Fourier map     |
| Least-squares matrix: full                                     | Hydrogen site location: inferred from neighbouring sites |
| $R[F^2 > 2\sigma(F^2)] = 0.059$                                | H-atom parameters constrained                            |
| $wR(F^2) = 0.146$  | $w = 1/[\sigma^2(F_o^2) + (0.0766P)^2]$                  |
| $S = 1.00$   | where $P = (F_o^2 + 2F_c^2)/3$                           |
| 2162 reflections   | $(\Delta/\sigma)_{\max} < 0.001$                         |
| 124 parameters   | $\Delta\rho_{\max} = 0.97 \text{ e } \text{\AA}^{-3}$    |
| Primary atom site location: structure-invariant direct methods | $\Delta\rho_{\min} = -1.62 \text{ e } \text{\AA}^{-3}$   |
|  | Extinction correction: none                              |

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

|     | $x$          | $y$         | $z$           | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|-------------|---------------|----------------------------------|
| Sn1 | 1.0000       | 0.5000      | 0.5000        | 0.0284 (3)                       |
| Br1 | 1.06913 (19) | 0.61962 (8) | 0.68737 (17)  | 0.0445 (4)                       |
| Br2 | 1.03622 (19) | 0.59582 (8) | 0.30353 (17)  | 0.0447 (4)                       |
| Br3 | 0.66959 (17) | 0.53289 (8) | 0.39554 (18)  | 0.0423 (4)                       |
| Br4 | 0.7063 (2)   | 0.32003 (9) | -0.12875 (19) | 0.0510 (5)                       |
| Br5 | 0.2397 (2)   | 0.45315 (8) | 0.10813 (18)  | 0.0483 (4)                       |
| N1  | 0.3496 (13)  | 0.2261 (5)  | 0.0181 (12)   | 0.032 (3)                        |
| H1  | 0.3148       | 0.1789      | 0.0281        | 0.039*                           |
| N2  | 0.5374 (16)  | 0.1644 (6)  | -0.0743 (13)  | 0.047 (3)                        |
| H2A | 0.6200       | 0.1663      | -0.1066       | 0.056*                           |
| H2B | 0.4940       | 0.1190      | -0.0650       | 0.056*                           |
| N3  | 0.1518 (14)  | 0.2743 (6)  | 0.1119 (13)   | 0.044 (3)                        |
| H3A | 0.1211       | 0.2259      | 0.1191        | 0.053*                           |
| H3B | 0.1029       | 0.3134      | 0.1386        | 0.053*                           |
| C1  | 0.4759 (17)  | 0.2325 (7)  | -0.0381 (15)  | 0.034 (3)                        |
| C2  | 0.5364 (17)  | 0.3061 (7)  | -0.0492 (17)  | 0.036 (3)                        |

## supplementary materials

---

|    |             |            |              |           |
|----|-------------|------------|--------------|-----------|
| C3 | 0.4653 (17) | 0.3720 (8) | -0.0099 (15) | 0.040 (4) |
| H3 | 0.5047      | 0.4227     | -0.0212      | 0.048*    |
| C4 | 0.3354 (18) | 0.3649 (7) | 0.0466 (15)  | 0.034 (3) |
| C5 | 0.2745 (17) | 0.2888 (8) | 0.0595 (15)  | 0.039 (3) |

### Atomic displacement parameters ( $\text{\AA}^2$ )

|     | $U^{11}$   | $U^{22}$   | $U^{33}$    | $U^{12}$    | $U^{13}$   | $U^{23}$    |
|-----|------------|------------|-------------|-------------|------------|-------------|
| Sn1 | 0.0293 (7) | 0.0234 (6) | 0.0365 (8)  | 0.0000 (5)  | 0.0169 (6) | -0.0002 (5) |
| Br1 | 0.0485 (9) | 0.0374 (7) | 0.0537 (10) | -0.0074 (7) | 0.0264 (8) | -0.0159 (7) |
| Br2 | 0.0494 (9) | 0.0430 (8) | 0.0499 (10) | -0.0016 (7) | 0.0279 (8) | 0.0093 (7)  |
| Br3 | 0.0271 (7) | 0.0319 (7) | 0.0677 (10) | 0.0009 (6)  | 0.0180 (7) | -0.0014 (7) |
| Br4 | 0.0472 (9) | 0.0493 (9) | 0.0706 (11) | -0.0030 (8) | 0.0381 (9) | -0.0045 (8) |
| Br5 | 0.0480 (9) | 0.0333 (7) | 0.0680 (11) | 0.0059 (7)  | 0.0272 (9) | -0.0091 (7) |
| N1  | 0.037 (6)  | 0.018 (5)  | 0.055 (8)   | -0.003 (5)  | 0.032 (6)  | -0.002 (5)  |
| N2  | 0.061 (9)  | 0.029 (6)  | 0.059 (8)   | 0.003 (6)   | 0.032 (7)  | -0.011 (6)  |
| N3  | 0.042 (7)  | 0.035 (6)  | 0.069 (9)   | -0.014 (6)  | 0.037 (7)  | -0.017 (6)  |
| C1  | 0.035 (8)  | 0.024 (6)  | 0.042 (9)   | 0.007 (6)   | 0.012 (7)  | -0.002 (6)  |
| C2  | 0.037 (8)  | 0.016 (6)  | 0.063 (9)   | 0.004 (6)   | 0.028 (7)  | 0.001 (6)   |
| C3  | 0.038 (9)  | 0.032 (7)  | 0.048 (9)   | -0.018 (7)  | 0.015 (8)  | 0.007 (6)   |
| C4  | 0.049 (9)  | 0.017 (6)  | 0.034 (8)   | -0.002 (6)  | 0.013 (7)  | -0.004 (5)  |
| C5  | 0.032 (8)  | 0.046 (8)  | 0.030 (8)   | -0.003 (7)  | 0.004 (6)  | -0.012 (6)  |

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

|  |             |            |            |
|--|-------------|------------|------------|
| Sn1—Br1                                | 2.6002 (13) | N3—H3A     | 0.8600     |
| Sn1—Br1 <sup>i</sup>                   | 2.6002 (13) | N3—H3B     | 0.8600     |
| Sn1—Br2                                | 2.5768 (14) | C1—N1      | 1.362 (16) |
| Sn1—Br2 <sup>i</sup>                   | 2.5768 (14) | C1—N2      | 1.349 (15) |
| Sn1—Br3                                | 2.6131 (14) | C1—C2      | 1.350 (17) |
| Sn1—Br3 <sup>i</sup>                   | 2.6131 (14) | C2—Br4     | 1.868 (12) |
| N1—C5                                  | 1.357 (15)  | C2—C3      | 1.372 (18) |
| N1—H1                                  | 0.8600      | C3—C4      | 1.392 (18) |
| N2—H2A                                 | 0.8600      | C3—H3      | 0.9300     |
| N2—H2B                                 | 0.8600      | C4—Br5     | 1.878 (12) |
| N3—C5                                  | 1.328 (16)  | C4—C5      | 1.394 (18) |
| Br1—Sn1—Br1 <sup>i</sup>               | 180.0       | H2A—N2—H2B | 120.0      |
| Br1 <sup>i</sup> —Sn1—Br3              | 88.56 (5)   | C5—N3—H3A  | 120.0      |
| Br1—Sn1—Br3 <sup>i</sup>               | 88.56 (5)   | C5—N3—H3B  | 120.0      |
| Br1—Sn1—Br3                            | 91.44 (5)   | H3A—N3—H3B | 120.0      |
| Br1 <sup>i</sup> —Sn1—Br3 <sup>i</sup> | 91.44 (5)   | N2—C1—C2   | 123.9 (12) |
| Br2—Sn1—Br1                            | 88.21 (5)   | N2—C1—N1   | 117.8 (11) |
| Br2 <sup>i</sup> —Sn1—Br1 <sup>i</sup> | 88.21 (5)   | C2—C1—N1   | 118.3 (10) |
| Br2 <sup>i</sup> —Sn1—Br1              | 91.79 (5)   | C1—C2—Br4  | 120.9 (9)  |
| Br2—Sn1—Br1 <sup>i</sup>               | 91.79 (5)   | C1—C2—C3   | 119.7 (11) |
| Br2 <sup>i</sup> —Sn1—Br2              | 180.0       | C3—C2—Br4  | 119.3 (9)  |

|  |             |              |            |
|--|-------------|--------------|------------|
| Br2—Sn1—Br3                            | 89.60 (5)   | C2—C3—C4     | 121.6 (11) |
| Br2 <sup>i</sup> —Sn1—Br3 <sup>i</sup> | 89.61 (5)   | C2—C3—H3     | 119.2      |
| Br2 <sup>i</sup> —Sn1—Br3              | 90.39 (5)   | C4—C3—H3     | 119.2      |
| Br2—Sn1—Br3 <sup>i</sup>               | 90.39 (5)   | C3—C4—Br5    | 123.1 (9)  |
| Br3 <sup>i</sup> —Sn1—Br3              | 180.0       | C3—C4—C5     | 118.6 (11) |
| C1—N1—H1                               | 117.6       | C5—C4—Br5    | 118.3 (10) |
| C5—N1—C1                               | 124.9 (10)  | N1—C5—C4     | 116.9 (12) |
| C5—N1—H1                               | 117.6       | N3—C5—N1     | 118.8 (12) |
| C1—N2—H2A                              | 120.0       | N3—C5—C4     | 124.2 (12) |
| C1—N2—H2B                              | 120.0       |              |            |
| N2—C1—N1—C5                            | -179.9 (12) | C2—C3—C4—C5  | -2(2)      |
| C2—C1—N1—C5                            | 2(2)        | C2—C3—C4—Br5 | 177.8 (11) |
| N2—C1—C2—C3                            | 179.9 (14)  | C1—N1—C5—N3  | 179.6 (13) |
| N1—C1—C2—C3                            | -3(2)       | C1—N1—C5—C4  | -2(2)      |
| N2—C1—C2—Br4                           | 4(2)        | C3—C4—C5—N3  | 179.8 (13) |
| N1—C1—C2—Br4                           | -178.7 (10) | Br5—C4—C5—N3 | 0.4 (19)   |
| C1—C2—C3—C4                            | 2(2)        | C3—C4—C5—N1  | 1.2 (19)   |
| Br4—C2—C3—C4                           | 178.5 (11)  | Br5—C4—C5—N1 | -178.2 (9) |

Symmetry codes: (i)  $-x+2, -y+1, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

| <i>D</i> —H $\cdots$ <i>A</i>      | <i>D</i> —H | H $\cdots$ <i>A</i> | <i>D</i> $\cdots$ <i>A</i> | <i>D</i> —H $\cdots$ <i>A</i> |
|------------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N1—H1 $\cdots$ Br3 <sup>ii</sup>   | 0.86        | 2.54                | 3.354 (9)                  | 159                           |
| N2—H2B $\cdots$ Br3 <sup>ii</sup>  | 0.86        | 2.88                | 3.612 (12)                 | 144                           |
| N3—H3A $\cdots$ Br2 <sup>ii</sup>  | 0.86        | 2.79                | 3.608 (10)                 | 160                           |
| N3—H3B $\cdots$ Br1 <sup>iii</sup> | 0.86        | 2.82                | 3.604 (10)                 | 153                           |

Symmetry codes: (ii)  $-x+1, y-1/2, -z+1/2$ ; (iii)  $-x+1, -y+1, -z+1$ .

Fig. 1




Fig. 2

