

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[μ -10,21-Dimethyl-3,6,14,17-tetraaza-tricyclo[17.3.1.1^{8,12}]tetracos-1(23),-8(24),9,11,19,21-hexaene-23,24-diolato- κ^8 N³,N⁶,O²³,O²⁴:N¹⁴,N¹⁷,O²³,O²⁴]-bis[(nitrate- κ^2 O,O')nickel(II)]

Quan-Jun Li, Jian-Fang Ma,* Jie Liu and Ting-Ting Han

Department of Chemistry, Northeast Normal University, Changchun 130024, People's Republic of China

Correspondence e-mail: majf247nenu@yahoo.com.cn


Received 11 March 2009; accepted 19 March 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.043; wR factor = 0.117; data-to-parameter ratio = 12.4.

In the title centrosymmetric dinuclear nickel complex, $[\text{Ni}_2(\text{C}_{22}\text{H}_{30}\text{N}_4\text{O}_2)(\text{NO}_3)_2]$, each of the two Ni^{II} atoms has a distorted octahedral geometry, defined by two N atoms and two O atoms from the macrocyclic ligand and two O atoms from a chelating nitrate anion. The two Ni atoms are bridged by two phenolate O atoms, forming a four-membered Ni_2O_2 ring.

Related literature

For general background, see: Caldwell & Crumbliss (1998); Rosa *et al.* (1998). For related structures, see: Aromi *et al.* (2002). For the ligand synthesis, see: Mandal & Nag (1986).



Experimental

Crystal data

$[\text{Ni}_2(\text{C}_{22}\text{H}_{30}\text{N}_4\text{O}_2)(\text{NO}_3)_2]$
 $M_r = 623.90$
 Trigonal, $R\bar{3}$
 $a = 25.020$ (5) Å
 $c = 10.616$ (5) Å
 $V = 5755$ (3) Å³

$Z = 9$
 Mo $K\alpha$ radiation
 $\mu = 1.53$ mm⁻¹
 $T = 293$ K
 $0.40 \times 0.30 \times 0.25$ mm

Data collection

Bruker APEX CCD diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.495$, $T_{\text{max}} = 0.609$
 (expected range = 0.554–0.682)

9432 measured reflections
 2213 independent reflections
 1745 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.107$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.117$
 $S = 1.03$
 2213 reflections
 178 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 1.06$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ni1—O1	2.000 (2)	Ni1—N1	2.054 (3)
Ni1—O1 ⁱ	2.006 (2)	Ni1—O3	2.134 (3)
Ni1—N2	2.038 (3)	Ni1—O2	2.183 (3)

Symmetry code: (i) $-x + \frac{2}{3}$, $-y + \frac{1}{3}$, $-z + \frac{1}{3}$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

We thank the National Natural Science Foundation of China (grant No. 20471014), the Program for New Century Excellent Talents in Chinese Universities (grant No. NCET-05-0320), the Fok Ying Tung Education Foundation and the Analysis and Testing Foundation of Northeast Normal University for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2188).

References

- Aromi, G., Gamez, P., Roubeau, O., Carrero-Berzal, P., Kooijman, H. L., Spek, A. L., Driesser, W. & Reedijk, J. (2002). *Eur. J. Inorg. Chem.* **5**, 1046–1048.
 Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Caldwell, D. C. & Crumbliss, L. A. (1998). *Inorg. Chem.* **37**, 1906–1912.
 Mandal, S. K. & Nag, K. (1986). *J. Org. Chem.* **51**, 3900–3902.
 Rosa, T. D., Young, G. V. & Coucouvanis, D. (1998). *Inorg. Chem.* **37**, 5042–5043.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m522 [doi:10.1107/S1600536809010174]

[μ -10,21-Dimethyl-3,6,14,17-tetraazatricyclo[17.3.1.1^{8,12}]tetracos-1(23),8(24),9,11,19,21-hexaene-23,24-diolato- κ^8 N³,N⁶,O²³,O²⁴:N¹⁴,N¹⁷,O²³,O²⁴]bis[(nitrate- κ^2 O,O')nickel(II)]

Q.-J. Li, J.-F. Ma, J. Liu and T.-T. Han

Comment

Crown ether compounds have attracted much interest as a result of their significance in biological transport systems (Caldwell & Crumbliss, 1998). In addition, crown ether compounds are found to have functions in selective molecular recognition (Rosa *et al.*, 1998). To further widen the scope of applications of crown ether, there is a need to prepare new series of crown ether compounds. In this work, a new dinuclear nickel(II) compound has been synthesized and its structure is reported here.

As shown in Fig. 1, the title compound is a centrosymmetric dinuclear nickel complex. The coordination environment around each Ni^{II} atom is distorted octahedral, with one N atom and one O atom from the macrocyclic ligand and two O atoms from the nitrate anion occupying the equatorial plane, and the other N atom and O atom from the ligand occupying the axial positions. In the complex molecule, two Ni atoms are bridged by two phenolate O atoms, generating a four-membered Ni₂O₂ ring, with a Ni...Ni distance of 2.9737 (10) Å. The Ni—O and Ni—N distances are normal (Aromi *et al.*, 2002).

Experimental

The macrocyclic ligand, C₂₂H₃₂N₄O₂ (H₂L), was prepared by the reported procedure (Mandal & Nag, 1986). A mixture of H₂L (0.10 g, 0.26 mmol) and Ni(NO₃)₂·6H₂O (0.15 g, 0.52 mmol) in methanol (20 ml) was stirred for 10 min. The resulting solution was filtered. Green single crystals were obtained by slow evaporation of the filtrate at room temperature (yield 56%).

Refinement

H atoms bound to C atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 (aromatic), 0.97 (CH₂) and 0.96 (CH₃) Å and with $U_{\text{iso}} = 1.2(1.5 \text{ for methyl})U_{\text{eq}}(\text{C})$. The imino H atoms were located in a difference Fourier map and refined with $U_{\text{iso}}(\text{H}) = 0.128 \text{ \AA}^2$. The highest residual electron density was found 1.03 Å from Ni1 and the deepest hole 0.76 Å from Ni1.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity. [Symmetry code: (i) $-x + 2/3, -y + 1/3, -z + 1/3$.]

supplementary materials

[μ -10,21-Dimethyl-3,6,14,17-tetraazatricyclo[17.3.1.1^{8,12}]tetracos- 1(23),8(24),9,11,19,21-hexaene-23,24-diolato- $\kappa^8 N^3, N^6, O^{23}, O^{24}; N^{14}, N^{17}, O^{23}, O^{24}$]bis[(nitrate- $\kappa^2 O, O'$)nickel(II)]

Crystal data

[Ni ₂ (C ₂₂ H ₃₀ N ₄ O ₂)(NO ₃) ₂]	$Z = 9$
$M_r = 623.90$	$F_{000} = 2916$
Trigonal, $R\bar{3}$	$D_x = 1.620 \text{ Mg m}^{-3}$
Hall symbol: -R 3	Mo $K\alpha$ radiation
$a = 25.020 (5) \text{ \AA}$	$\lambda = 0.71069 \text{ \AA}$
$b = 25.020 (5) \text{ \AA}$	Cell parameters from 3000 reflections
$c = 10.616 (5) \text{ \AA}$	$\theta = 2.4\text{--}28.4^\circ$
$\alpha = 90^\circ$	$\mu = 1.53 \text{ mm}^{-1}$
$\beta = 90^\circ$	$T = 293 \text{ K}$
$\gamma = 120^\circ$	Block, green
$V = 5755 (3) \text{ \AA}^3$	$0.40 \times 0.30 \times 0.25 \text{ mm}$

Data collection

Bruker APEX CCD diffractometer	2213 independent reflections
Radiation source: fine-focus sealed tube	1745 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.107$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 24.9^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -15 \rightarrow 29$
$T_{\text{min}} = 0.495, T_{\text{max}} = 0.609$	$k = -29 \rightarrow 23$
9432 measured reflections	$l = -12 \rightarrow 12$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.043$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.117$	$w = 1/[\sigma^2(F_o^2) + (0.066P)^2]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
2213 reflections	$(\Delta/\sigma)_{\text{max}} = 0.007$
178 parameters	$\Delta\rho_{\text{max}} = 1.06 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.355561 (19)	0.13534 (2)	0.25357 (4)	0.03150 (19)
C1	0.04882 (19)	-0.0620 (2)	0.3314 (4)	0.0658 (13)
H1A	0.0488	-0.0770	0.4147	0.099*
H1B	0.0372	-0.0948	0.2717	0.099*
H1C	0.0199	-0.0475	0.3276	0.099*
C2	0.11262 (17)	-0.00973 (17)	0.3005 (3)	0.0431 (9)
C3	0.12561 (16)	0.01995 (17)	0.1862 (3)	0.0410 (9)
H3	0.0947	0.0056	0.1255	0.049*
C4	0.18315 (15)	0.07079 (16)	0.1570 (3)	0.0353 (8)
C5	0.22847 (15)	0.09460 (15)	0.2510 (3)	0.0347 (8)
C6	0.16018 (17)	0.01062 (17)	0.3868 (3)	0.0446 (9)
H6	0.1536	-0.0111	0.4617	0.054*
C7	0.21738 (16)	0.06230 (16)	0.3656 (3)	0.0374 (8)
C8	0.26663 (17)	0.08766 (18)	0.4652 (3)	0.0443 (9)
H8A	0.2746	0.1284	0.4892	0.053*
H8B	0.2515	0.0614	0.5391	0.053*
C9	0.37778 (18)	0.13034 (17)	0.5104 (3)	0.0430 (9)
H9A	0.4112	0.1221	0.4949	0.052*
H9B	0.3646	0.1196	0.5972	0.052*
C10	0.40046 (18)	0.19842 (17)	0.4892 (3)	0.0421 (9)
H10A	0.3704	0.2085	0.5224	0.050*
H10B	0.4390	0.2230	0.5342	0.050*
C11	0.47350 (15)	0.23666 (16)	0.3077 (3)	0.0383 (8)
H11A	0.4824	0.2031	0.3091	0.046*
H11B	0.5022	0.2685	0.3644	0.046*
N1	0.32528 (14)	0.09201 (14)	0.4244 (3)	0.0377 (7)
N2	0.41009 (14)	0.21377 (13)	0.3532 (2)	0.0355 (7)
N3	0.37310 (14)	0.05216 (15)	0.1596 (3)	0.0479 (8)
O1	0.28230 (10)	0.14687 (10)	0.23716 (19)	0.0342 (5)
O2	0.41509 (12)	0.09599 (12)	0.2220 (2)	0.0455 (6)
O3	0.32302 (12)	0.05220 (12)	0.1480 (2)	0.0459 (6)
O4	0.38022 (15)	0.01126 (15)	0.1141 (4)	0.0853 (11)
HN1	0.320 (3)	0.055 (3)	0.423 (6)	0.128*
HN2	0.401 (3)	0.242 (2)	0.336 (6)	0.128*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0299 (3)	0.0299 (3)	0.0353 (3)	0.0154 (2)	-0.00202 (16)	0.00065 (16)
C1	0.041 (3)	0.060 (3)	0.081 (3)	0.014 (2)	0.014 (2)	0.004 (2)
C2	0.037 (2)	0.035 (2)	0.052 (2)	0.0137 (17)	0.0119 (16)	-0.0027 (16)
C3	0.033 (2)	0.038 (2)	0.055 (2)	0.0200 (18)	-0.0023 (16)	-0.0069 (16)
C4	0.0299 (19)	0.0322 (19)	0.0451 (18)	0.0165 (16)	0.0011 (14)	-0.0002 (14)
C5	0.0322 (19)	0.033 (2)	0.0424 (18)	0.0193 (17)	0.0038 (14)	-0.0001 (14)

supplementary materials

C6	0.044 (2)	0.042 (2)	0.049 (2)	0.022 (2)	0.0114 (17)	0.0067 (16)
C7	0.037 (2)	0.037 (2)	0.0402 (18)	0.0193 (17)	0.0053 (15)	0.0017 (14)
C8	0.043 (2)	0.051 (2)	0.0380 (18)	0.024 (2)	0.0040 (15)	0.0054 (16)
C9	0.047 (2)	0.045 (2)	0.0366 (18)	0.0226 (19)	-0.0053 (15)	0.0029 (15)
C10	0.048 (2)	0.043 (2)	0.0359 (18)	0.0227 (19)	-0.0046 (15)	-0.0049 (15)
C11	0.033 (2)	0.034 (2)	0.049 (2)	0.0181 (17)	-0.0105 (15)	-0.0059 (15)
N1	0.0404 (18)	0.0370 (18)	0.0381 (14)	0.0211 (16)	-0.0010 (12)	0.0024 (13)
N2	0.0366 (17)	0.0327 (17)	0.0383 (15)	0.0182 (15)	-0.0035 (12)	-0.0001 (12)
N3	0.0349 (19)	0.040 (2)	0.067 (2)	0.0180 (16)	-0.0012 (15)	-0.0111 (16)
O1	0.0289 (13)	0.0311 (13)	0.0408 (12)	0.0137 (11)	-0.0007 (10)	0.0025 (10)
O2	0.0341 (14)	0.0377 (15)	0.0634 (16)	0.0171 (13)	-0.0070 (12)	-0.0076 (12)
O3	0.0329 (15)	0.0434 (16)	0.0570 (15)	0.0159 (13)	-0.0059 (11)	-0.0075 (11)
O4	0.058 (2)	0.062 (2)	0.139 (3)	0.0324 (18)	-0.0003 (19)	-0.048 (2)

Geometric parameters (\AA , $^\circ$)

Ni1—O1	2.000 (2)	C7—C8	1.502 (5)
Ni1—O1 ⁱ	2.006 (2)	C8—N1	1.481 (5)
Ni1—N2	2.038 (3)	C8—H8A	0.9700
Ni1—N1	2.054 (3)	C8—H8B	0.9700
Ni1—O3	2.134 (3)	C9—N1	1.489 (4)
Ni1—O2	2.183 (3)	C9—C10	1.519 (5)
Ni1—Ni1 ⁱ	2.9737 (10)	C9—H9A	0.9700
C1—C2	1.510 (5)	C9—H9B	0.9700
C1—H1A	0.9600	C10—N2	1.483 (4)
C1—H1B	0.9600	C10—H10A	0.9700
C1—H1C	0.9600	C10—H10B	0.9700
C2—C3	1.374 (5)	C11—N2	1.473 (4)
C2—C6	1.382 (5)	C11—C4 ⁱ	1.505 (5)
C3—C4	1.398 (5)	C11—H11A	0.9700
C3—H3	0.9300	C11—H11B	0.9700
C4—C5	1.400 (5)	N1—HN1	0.87 (6)
C4—C11 ⁱ	1.505 (5)	N2—HN2	0.86 (6)
C5—O1	1.336 (4)	N3—O4	1.223 (4)
C5—C7	1.408 (5)	N3—O3	1.260 (4)
C6—C7	1.386 (5)	N3—O2	1.262 (4)
C6—H6	0.9300		
O1—Ni1—O1 ⁱ	84.17 (9)	N1—C8—C7	113.5 (3)
O1—Ni1—N2	97.29 (10)	N1—C8—H8A	108.9
O1 ⁱ —Ni1—N2	87.95 (10)	C7—C8—H8A	108.9
O1—Ni1—N1	91.74 (10)	N1—C8—H8B	108.9
O1 ⁱ —Ni1—N1	172.81 (10)	C7—C8—H8B	108.9
N2—Ni1—N1	86.72 (12)	H8A—C8—H8B	107.7
O1—Ni1—O3	99.46 (9)	N1—C9—C10	110.3 (3)
O1 ⁱ —Ni1—O3	91.39 (10)	N1—C9—H9A	109.6
N2—Ni1—O3	163.09 (11)	C10—C9—H9A	109.6
N1—Ni1—O3	95.11 (11)	N1—C9—H9B	109.6

O1—Ni1—O2	158.83 (9)	C10—C9—H9B	109.6
O1 ⁱ —Ni1—O2	92.93 (9)	H9A—C9—H9B	108.1
N2—Ni1—O2	103.57 (11)	N2—C10—C9	110.8 (3)
N1—Ni1—O2	92.99 (11)	N2—C10—H10A	109.5
O3—Ni1—O2	59.57 (10)	C9—C10—H10A	109.5
O1—Ni1—Ni1 ⁱ	42.16 (6)	N2—C10—H10B	109.5
O1 ⁱ —Ni1—Ni1 ⁱ	42.01 (6)	C9—C10—H10B	109.5
N2—Ni1—Ni1 ⁱ	93.51 (8)	H10A—C10—H10B	108.1
N1—Ni1—Ni1 ⁱ	133.62 (9)	N2—C11—C4 ⁱ	112.7 (3)
O3—Ni1—Ni1 ⁱ	97.29 (7)	N2—C11—H11A	109.0
O2—Ni1—Ni1 ⁱ	131.44 (7)	C4 ⁱ —C11—H11A	109.0
C2—C1—H1A	109.5	N2—C11—H11B	109.0
C2—C1—H1B	109.5	C4 ⁱ —C11—H11B	109.0
H1A—C1—H1B	109.5	H11A—C11—H11B	107.8
C2—C1—H1C	109.5	C8—N1—C9	113.0 (3)
H1A—C1—H1C	109.5	C8—N1—Ni1	112.8 (2)
H1B—C1—H1C	109.5	C9—N1—Ni1	103.2 (2)
C3—C2—C6	117.3 (3)	C8—N1—HN1	108 (4)
C3—C2—C1	121.5 (4)	C9—N1—HN1	108 (4)
C6—C2—C1	121.2 (4)	Ni1—N1—HN1	112 (4)
C2—C3—C4	123.0 (3)	C11—N2—C10	115.1 (3)
C2—C3—H3	118.5	C11—N2—Ni1	106.0 (2)
C4—C3—H3	118.5	C10—N2—Ni1	108.2 (2)
C3—C4—C5	118.4 (3)	C11—N2—HN2	107 (4)
C3—C4—C11 ⁱ	118.1 (3)	C10—N2—HN2	110 (4)
C5—C4—C11 ⁱ	123.5 (3)	Ni1—N2—HN2	110 (4)
O1—C5—C4	122.9 (3)	O4—N3—O3	121.5 (3)
O1—C5—C7	118.0 (3)	O4—N3—O2	121.9 (3)
C4—C5—C7	119.1 (3)	O3—N3—O2	116.6 (3)
C2—C6—C7	122.2 (3)	C5—O1—Ni1	113.40 (19)
C2—C6—H6	118.9	C5—O1—Ni1 ⁱ	125.52 (19)
C7—C6—H6	118.9	Ni1—O1—Ni1 ⁱ	95.83 (9)
C6—C7—C5	119.4 (3)	N3—O2—Ni1	90.8 (2)
C6—C7—C8	121.7 (3)	N3—O3—Ni1	93.11 (19)
C5—C7—C8	118.7 (3)		
C6—C2—C3—C4	-3.1 (5)	O2—Ni1—N2—C11	32.8 (2)
C1—C2—C3—C4	176.2 (4)	Ni1 ⁱ —Ni1—N2—C11	-101.34 (19)
C2—C3—C4—C5	-3.7 (5)	O1—Ni1—N2—C10	92.5 (2)
C2—C3—C4—C11 ⁱ	175.9 (3)	O1 ⁱ —Ni1—N2—C10	176.3 (2)
C3—C4—C5—O1	-172.4 (3)	N1—Ni1—N2—C10	1.1 (2)
C11 ⁱ —C4—C5—O1	8.1 (5)	O3—Ni1—N2—C10	-95.6 (4)
C3—C4—C5—C7	7.8 (5)	O2—Ni1—N2—C10	-91.1 (2)
C11 ⁱ —C4—C5—C7	-171.7 (3)	Ni1 ⁱ —Ni1—N2—C10	134.7 (2)
C3—C2—C6—C7	5.8 (5)	C4—C5—O1—Ni1	-120.4 (3)
C1—C2—C6—C7	-173.5 (4)	C7—C5—O1—Ni1	59.4 (3)

supplementary materials

C2—C6—C7—C5	-1.7 (5)	C4—C5—O1—Ni ⁱ	-4.0 (4)
C2—C6—C7—C8	173.8 (3)	C7—C5—O1—Ni ⁱ	175.8 (2)
O1—C5—C7—C6	174.9 (3)	O1 ⁱ —Ni1—O1—C5	132.9 (2)
C4—C5—C7—C6	-5.2 (5)	N2—Ni1—O1—C5	-139.9 (2)
O1—C5—C7—C8	-0.7 (5)	N1—Ni1—O1—C5	-53.0 (2)
C4—C5—C7—C8	179.1 (3)	O3—Ni1—O1—C5	42.4 (2)
C6—C7—C8—N1	123.4 (4)	O2—Ni1—O1—C5	49.8 (3)
C5—C7—C8—N1	-61.0 (4)	Ni1 ⁱ —Ni1—O1—C5	132.9 (2)
N1—C9—C10—N2	-48.9 (4)	O1 ⁱ —Ni1—O1—Ni1 ⁱ	0.0
C7—C8—N1—C9	166.4 (3)	N2—Ni1—O1—Ni1 ⁱ	87.17 (11)
C7—C8—N1—Ni1	49.9 (4)	N1—Ni1—O1—Ni1 ⁱ	174.08 (11)
C10—C9—N1—C8	-76.0 (4)	O3—Ni1—O1—Ni1 ⁱ	-90.44 (10)
C10—C9—N1—Ni1	46.1 (3)	O2—Ni1—O1—Ni1 ⁱ	-83.0 (3)
N2—Ni1—N1—C8	96.4 (2)	O4—N3—O2—Ni1	-179.5 (4)
O3—Ni1—N1—C8	-100.4 (2)	O3—N3—O2—Ni1	-0.4 (3)
O2—Ni1—N1—C8	-160.1 (2)	O1—Ni1—O2—N3	-8.2 (4)
Ni1 ⁱ —Ni1—N1—C8	4.7 (3)	O1 ⁱ —Ni1—O2—N3	-89.6 (2)
N2—Ni1—N1—C9	-25.8 (2)	N2—Ni1—O2—N3	-178.2 (2)
O3—Ni1—N1—C9	137.3 (2)	N1—Ni1—O2—N3	94.4 (2)
O2—Ni1—N1—C9	77.6 (2)	O3—Ni1—O2—N3	0.24 (19)
Ni1 ⁱ —Ni1—N1—C9	-117.5 (2)	Ni1 ⁱ —Ni1—O2—N3	-71.0 (2)
C4 ⁱ —C11—N2—C10	-169.3 (3)	O4—N3—O3—Ni1	179.5 (4)
C4 ⁱ —C11—N2—Ni1	71.2 (3)	O2—N3—O3—Ni1	0.4 (3)
C9—C10—N2—C11	-94.0 (4)	O1—Ni1—O3—N3	176.7 (2)
C9—C10—N2—Ni1	24.3 (4)	O1 ⁱ —Ni1—O3—N3	92.3 (2)
O1—Ni1—N2—C11	-143.54 (19)	N2—Ni1—O3—N3	4.8 (5)
O1 ⁱ —Ni1—N2—C11	-59.7 (2)	N1—Ni1—O3—N3	-90.7 (2)
N1—Ni1—N2—C11	125.1 (2)	O2—Ni1—O3—N3	-0.24 (19)
O3—Ni1—N2—C11	28.3 (5)	Ni1 ⁱ —Ni1—O3—N3	134.08 (19)

Symmetry codes: (i) $-x+2/3, -y+1/3, -z+1/3$.

