

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,3',5,5'-Tetranitrobiphenyl

N. Hammond,^a P. Carvalho,^a Y. Wu^a and M. A. Avery^{a,b,c,*}

^aUniversity of Mississippi, Department of Medicinal Chemistry, 417 Faser Hall, University, MS 38677, USA, ^bNational Center for Natural Products Research, Research Institute of Pharmaceutical Sciences, School of Pharmacy, University of Mississippi, University, MS 38677, USA, and ^cDepartment of Chemistry and Biochemistry, University of Mississippi, University, MS 38677, USA
Correspondence e-mail: mavery@olemiss.edu


Received 5 March 2009; accepted 25 March 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.030; wR factor = 0.081; data-to-parameter ratio = 10.3.

The title compound, $\text{C}_{12}\text{H}_6\text{N}_4\text{O}_8$, is a biphenyl system that was synthesized as a building block for a new series of antimalarial compounds. The aromatic rings are oriented at a dihedral angle of $45.5(2)^\circ$, and intermolecular short $\text{O}\cdots\text{O}$ contacts form a chain along the b axis. The strength of the interactions involved in this chain cause one of the rings to be slightly distorted, with the torsion angle between the nitro groups being $23.4(2)^\circ$, whereas, in the other ring, both nitro systems are parallel, forming an angle of $9.6(2)^\circ$ with the plane of the aromatic ring to which they are bound. Furthermore, the three ring C atoms around the ring–ring linkage belong to a plane inclined by $4.5(1)^\circ$ in relation to the plane containing the other three C atoms, i.e. $(\text{NO}_2)\text{-C}-\text{C}-\text{C}(\text{NO}_2)$. This distortion of the ring causes uncommonly short intermolecular $\text{O}\cdots\text{O}$ [$3.038(2)$ Å] and $\text{O}\cdots\text{C}$ [$3.000(4)$ and $3.214(1)$ Å] contacts.

Related literature

For the previous synthesis of the title compound and its stability studies, see Case (1942) and Hoffsommer & McCullough (1968). For the use of polynitroaromatic compounds as explosives, see Davis (1941) and Keshavarz & Pouretedal (2005). For their mutagenic and carcinogenic properties, see Debnath *et al.* (1991). For previous studies showing distortions induced in aromatic rings, see Murray-Rust (1982), Allen *et al.* (1998), and Khurstalev *et al.* (2005). For the preparation, see: Goossen *et al.* (2007).



Experimental

Crystal data

$\text{C}_{12}\text{H}_6\text{N}_4\text{O}_8$
 $M_r = 334.21$
 Orthorhombic, $Pbca$
 $a = 10.0683(1)$ Å
 $b = 15.4640(2)$ Å
 $c = 16.3436(2)$ Å
 $V = 2544.64(5)$ Å³
 $Z = 8$
 Cu $K\alpha$ radiation
 $\mu = 1.32$ mm⁻¹
 $T = 100$ K
 $0.18 \times 0.15 \times 0.11$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: none
 30954 measured reflections
 2315 independent reflections
 2294 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.081$
 $S = 1.09$
 2315 reflections
 225 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors thank the Center for Disease Control and Prevention, USA, for providing financial assistance (CDC cooperative agreements 1U01 CI000211-03 and 1U01 CI000362-01). This investigation was conducted in a facility constructed with support from Research Facilities Improvement Program grant No. C06Rr-14503-01 from the National Center for Research Resources, National Institutes of Health.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2105).

References

- Allen, F. H., Baalham, C. A., Lommerse, J. P. M. & Raithby, P. R. (1998). *Acta Cryst.* **B54**, 320–329.
 Bruker (2005). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2008). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
 Case, F. H. (1942). *J. Am. Chem. Soc.* **64**, 1848–1852.

- Davis, T. L. (1941). *Chemistry of Powder and Explosives*. Hollywood, CA: Angriff Press.
- Debnath, A. K., Lopez de Compadre, R. L., Debnath, G., Shusterman, A. J. & Hansch, C. (1991). *J. Med. Chem.* **34**, 786–797.
- Goossen, L. J., Rodriguez, N., Melzer, B., Linder, C., Deng, G. & Levy, L. M. (2007). *J. Am. Chem. Soc.* **129**, 4824–4833.
- Hoffsommer, J. C. & McCullough, J. F. (1968). *J. Chromatogr.* **38**, 508–514.
- Keshavarz, M. H. & Pouretedal, H. R. (2005). *J. Hazard. Mater.* **124**, 27–33.
- Khrustalev, V. N., Vasil'kov, A. Y. & Antipin, M. Y. (2005). *Acta Cryst.* **B61**, 304–311.
- Murray-Rust, P. (1982). *Acta Cryst.* **B38**, 2818–2825.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o1052-o1053 [doi:10.1107/S1600536809011088]

3,3',5,5'-Tetranitrobiphenyl

N. Hammond, P. Carvalho, Y. Wu and M. A. Avery

Comment

Synthesis of the 3,3',5,5'-tetranitrobiphenyl has previously been reported by Case (1942), from the reaction of 3,5-dinitroiodobenzene and copper at 270°C. This paper reports the crystal structure of the title compound, obtained by reaction of 1-bromo-3,5-dinitrobenzene and 3,5-dinitrobenzoic acid in a sealed microwave tube in 1,2-dimethoxyethane. It was found to have activity against *Plasmodium falciparum* with an IC₅₀ of 5.7 μM against the chloroquine resistant D6 clone and 3.9 μM against the W2 clone.

Experimental

The title compound was prepared by decarboxylative coupling as previously published (Goossen *et al.*, 2007). Briefly, 1-bromo-3,5-dinitrobenzene (247 mg, 1 mmol), 3,5-dinitrobenzoic acid (211 mg, 1 mmol), copper(II) bromide (180 mg, 0.8 mmol), 1,2-dimethoxyethane (3 ml), [tetrakis(triphenylphosphine)palladium(0)] (100 mg, 0.086 mmol), [dichlorobis(triphenylphosphine)palladium (II)] (60 mg, 0.86 mmol), potassium carbonate (495 mg, 3 mmol) and water (1 ml) were added together in a microwave vial and microwaved at 160°C for 30 minutes while stirring, in a Biotage Initiator™ Sixty, with variable microwave output. To keep the programmed temperature, the initial output was 100 W for 5 minutes, then varying between 0 and 30 W for the next 25 minutes. The reaction mixture was shaken 3 times with water, then brine and dried over magnesium sulfate and concentrated. The resulting residue was purified by column chromatography on silica gel, eluting with ethyl acetate/hexanes (90:10) to afford 62 mg (20% yield) of the title compound. 3,3',5,5'-Tetranitrobiphenyl was recrystallized from MeOH:CHCl₃ (5:95) by slow evaporation of the solvent at room temperature.

Refinement

All H atoms were located in difference maps, and treated as riding atoms, except H2' (C—H = 0.95 Å) and H4' (C—H = 0.94 Å), with the following distance restraints: C—H = 0.93 Å, $U_{iso}=1.2U_{eq}$ (C) for $Csp2$.

Figures




Fig. 1. Molecular structure of the title compound, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.



Fig. 2. Partial view of the crystal packing of the title compound viewed along the *b* axis, showing molecules linked into chains by short intermolecular O1'...O4' interactions.

3,3',5,5'-Tetranitrobiphenyl

Crystal data

$C_{12}H_6N_4O_8$

$M_r = 334.21$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 10.0683$ (1) Å

$b = 15.4640$ (2) Å

$c = 16.3436$ (2) Å

$V = 2544.64$ (5) Å³

$Z = 8$

$F_{000} = 1360$

$D_x = 1.745$ Mg m⁻³

Melting point: not measured K

Cu *K*α radiation

$\lambda = 1.54178$ Å

Cell parameters from 9828 reflections

$\theta = 3.9$ – 67.6°

$\mu = 1.32$ mm⁻¹

$T = 100$ K

Block, yellow

$0.18 \times 0.15 \times 0.11$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100$ K

$P = 101.325$ kPa

φ and ω scans

Absorption correction: none

30954 measured reflections

2315 independent reflections

2294 reflections with $I > 2\sigma(I)$

$R_{int} = 0.021$

$\theta_{max} = 68.0^\circ$

$\theta_{min} = 5.4^\circ$

$h = -12 \rightarrow 12$

$k = -18 \rightarrow 18$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.081$

$S = 1.09$

2315 reflections

225 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0431P)^2 + 1.2464P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$

$\Delta\rho_{max} = 0.22$ e Å⁻³

$\Delta\rho_{min} = -0.31$ e Å⁻³

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.34665 (12)	0.30780 (8)	0.43269 (7)	0.0157 (3)
C2	0.33233 (12)	0.30138 (8)	0.51737 (7)	0.0156 (3)
H2	0.2894	0.3448	0.5466	0.019*
C3	0.38272 (12)	0.22966 (8)	0.55741 (7)	0.0162 (3)
C4	0.44841 (12)	0.16333 (8)	0.51778 (8)	0.0168 (3)
H4	0.4828	0.1160	0.5458	0.020*
C5	0.45990 (12)	0.17162 (8)	0.43369 (8)	0.0165 (3)
C6	0.41093 (12)	0.24155 (8)	0.39006 (7)	0.0162 (3)
H6	0.4206	0.2443	0.3335	0.019*
N1	0.37026 (10)	0.22503 (7)	0.64699 (6)	0.0182 (2)
N2	0.52978 (10)	0.10245 (7)	0.38839 (7)	0.0189 (2)
O1	0.30268 (11)	0.28031 (6)	0.68111 (5)	0.0259 (2)
O2	0.42893 (10)	0.16675 (6)	0.68273 (5)	0.0236 (2)
O3	0.58781 (10)	0.04674 (6)	0.42849 (6)	0.0276 (2)
O4	0.52572 (10)	0.10442 (6)	0.31365 (6)	0.0253 (2)
C1'	0.30389 (12)	0.38719 (8)	0.38844 (7)	0.0155 (3)
C2'	0.33837 (12)	0.46789 (8)	0.42007 (7)	0.0162 (3)
H2'	0.3828 (15)	0.4741 (10)	0.4712 (9)	0.018 (4)*
C3'	0.31428 (12)	0.54092 (8)	0.37355 (7)	0.0156 (3)
C4'	0.25353 (12)	0.53897 (8)	0.29759 (7)	0.0159 (3)
H4'	0.2391 (15)	0.5895 (9)	0.2666 (9)	0.018 (4)*
C5'	0.21448 (12)	0.45823 (8)	0.27031 (7)	0.0160 (3)
C6'	0.23899 (12)	0.38234 (8)	0.31302 (7)	0.0161 (3)
H6'	0.2127	0.3292	0.2918	0.019*
N1'	0.36202 (10)	0.62466 (7)	0.40437 (6)	0.0169 (2)
N2'	0.14792 (10)	0.45304 (7)	0.18984 (6)	0.0177 (2)
O1'	0.38516 (9)	0.63076 (6)	0.47790 (5)	0.0231 (2)
O2'	0.37639 (9)	0.68355 (6)	0.35486 (6)	0.0212 (2)
O3'	0.12418 (9)	0.52096 (6)	0.15399 (5)	0.0233 (2)
O4'	0.12047 (10)	0.38108 (6)	0.16341 (6)	0.0246 (2)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0138 (6)	0.0161 (6)	0.0173 (6)	-0.0029 (5)	-0.0013 (4)	0.0007 (5)
C2	0.0155 (6)	0.0143 (6)	0.0171 (6)	-0.0015 (5)	0.0002 (5)	-0.0013 (5)
C3	0.0170 (6)	0.0179 (6)	0.0137 (6)	-0.0044 (5)	-0.0009 (4)	0.0009 (5)
C4	0.0162 (6)	0.0149 (6)	0.0193 (6)	-0.0018 (5)	-0.0024 (5)	0.0028 (5)
C5	0.0148 (6)	0.0154 (6)	0.0192 (6)	-0.0021 (5)	0.0010 (5)	-0.0025 (5)
C6	0.0163 (6)	0.0183 (6)	0.0140 (6)	-0.0035 (5)	-0.0005 (4)	-0.0003 (5)
N1	0.0214 (5)	0.0174 (5)	0.0158 (5)	-0.0039 (4)	0.0003 (4)	0.0015 (4)
N2	0.0167 (5)	0.0169 (5)	0.0230 (6)	-0.0012 (4)	0.0017 (4)	-0.0019 (4)
O1	0.0409 (6)	0.0193 (5)	0.0175 (5)	0.0015 (4)	0.0061 (4)	-0.0017 (4)
O2	0.0248 (5)	0.0269 (5)	0.0191 (5)	0.0004 (4)	-0.0015 (4)	0.0084 (4)
O3	0.0309 (5)	0.0200 (5)	0.0320 (5)	0.0079 (4)	0.0017 (4)	0.0022 (4)
O4	0.0263 (5)	0.0304 (5)	0.0193 (5)	0.0025 (4)	0.0033 (4)	-0.0057 (4)
C1'	0.0137 (6)	0.0173 (6)	0.0155 (6)	0.0000 (4)	0.0026 (4)	0.0011 (5)
C2'	0.0146 (6)	0.0194 (6)	0.0147 (6)	0.0005 (5)	0.0003 (5)	0.0008 (5)
C3'	0.0138 (6)	0.0154 (6)	0.0177 (6)	-0.0008 (4)	0.0017 (5)	-0.0011 (5)
C4'	0.0138 (5)	0.0173 (6)	0.0166 (6)	0.0018 (5)	0.0024 (5)	0.0030 (5)
C5'	0.0133 (6)	0.0206 (6)	0.0141 (6)	0.0006 (5)	0.0001 (4)	0.0007 (5)
C6'	0.0154 (6)	0.0164 (6)	0.0165 (6)	-0.0005 (5)	0.0021 (5)	-0.0004 (5)
N1'	0.0144 (5)	0.0172 (5)	0.0192 (5)	0.0008 (4)	-0.0004 (4)	0.0003 (4)
N2'	0.0165 (5)	0.0211 (5)	0.0155 (5)	0.0003 (4)	-0.0004 (4)	0.0013 (4)
O1'	0.0283 (5)	0.0236 (5)	0.0175 (5)	-0.0042 (4)	-0.0031 (4)	-0.0022 (4)
O2'	0.0246 (5)	0.0154 (4)	0.0236 (5)	-0.0015 (4)	-0.0009 (4)	0.0042 (4)
O3'	0.0274 (5)	0.0222 (5)	0.0205 (5)	0.0013 (4)	-0.0059 (4)	0.0063 (4)
O4'	0.0314 (5)	0.0213 (5)	0.0210 (5)	-0.0028 (4)	-0.0074 (4)	-0.0024 (4)

Geometric parameters (\AA , $^\circ$)

C1—C1'	1.4885 (17)	C5'—N2'	1.4783 (16)
C1'—C2'	1.3947 (17)	C5—N2	1.4790 (16)
C1'—C6'	1.3971 (18)	C6—C1	1.3978 (17)
C2—C1	1.3950 (17)	C6—C5	1.3860 (17)
C2—C3	1.3842 (17)	C6—H6	0.9300
C2—H2	0.9300	C6'—H6'	0.9300
C2'—H2'	0.953 (15)	N1'—C3'	1.4702 (15)
C3'—C2'	1.3828 (17)	N1—O1	1.2266 (14)
C3—C4	1.3816 (18)	N1'—O1'	1.2278 (14)
C3—N1	1.4712 (15)	N1—O2	1.2257 (14)
C4'—C3'	1.3843 (17)	N1'—O2'	1.2268 (14)
C4—H4	0.9300	N2'—O3'	1.2263 (14)
C4'—H4'	0.942 (15)	N2—O3	1.2301 (14)
C5'—C4'	1.3828 (17)	N2—O4	1.2226 (15)
C5—C4	1.3851 (18)	N2'—O4'	1.2252 (14)
C5'—C6'	1.3876 (17)		
C1—C2—H2	120.4	C4—C5—N2	118.00 (11)

C1'—C2'—H2'	122.1 (9)	C5'—C4'—C3'	115.76 (11)
C1'—C6'—H6'	120.6	C5—C4—H4	122.1
C1—C6—H6	120.7	C5'—C4'—H4'	122.1 (9)
C2'—C1'—C1	119.08 (11)	C5—C6—C1	118.69 (11)
C2—C1—C1'	120.71 (11)	C5'—C6'—C1'	118.79 (11)
C2—C1—C6	119.36 (11)	C5'—C6'—H6'	120.6
C2'—C1'—C6'	119.44 (11)	C5—C6—H6	120.7
C2'—C3'—C4'	123.55 (11)	C6—C1—C1'	119.75 (11)
C2'—C3'—N1'	118.27 (11)	C6'—C1'—C1	121.31 (11)
C2—C3—N1	118.57 (11)	C6—C5—N2	118.42 (11)
C3'—C2'—C1'	118.90 (11)	C6'—C5'—N2'	118.84 (11)
C3—C2—C1	119.21 (11)	O1'—N1'—C3'	117.71 (10)
C3'—C2'—H2'	118.9 (9)	O1—N1—C3	117.76 (10)
C3—C2—H2	120.4	O2'—N1'—C3'	117.81 (10)
C3—C4—C5	115.88 (11)	O2—N1—C3	117.96 (10)
C3—C4—H4	122.1	O2—N1—O1	124.27 (10)
C3'—C4'—H4'	122.2 (9)	O2'—N1'—O1'	124.48 (11)
C4—C3—C2	123.28 (11)	O3—N2—C5	117.76 (10)
C4'—C3'—N1'	118.09 (10)	O3'—N2'—C5'	117.84 (10)
C4—C3—N1	118.11 (11)	O4'—N2'—C5'	117.72 (10)
C4'—C5'—C6'	123.42 (11)	O4—N2—C5	117.79 (10)
C4—C5—C6	123.57 (11)	O4'—N2'—O3'	124.44 (10)
C4'—C5'—N2'	117.70 (10)	O4—N2—O3	124.45 (11)
C1—C1'—C2'—C3'	171.43 (11)	C5'—C4'—C3'—C2'	1.28 (18)
C1—C1'—C6'—C5'	-172.98 (11)	C5'—C4'—C3'—N1'	177.92 (10)
C1—C2—C3—C4	0.57 (18)	C5—C6—C1—C1'	174.55 (11)
C1—C2—C3—N1	178.25 (11)	C5—C6—C1—C2	-0.67 (17)
C1—C6—C5—C4	0.20 (18)	C6—C1—C1'—C2'	-129.65 (12)
C1—C6—C5—N2	-178.99 (10)	C6—C1—C1'—C6'	45.44 (17)
C2—C1—C1'—C2'	45.51 (17)	C6'—C1'—C2'—C3'	-3.76 (18)
C2—C1—C1'—C6'	-139.40 (12)	C6—C5—C4—C3	0.62 (18)
C2'—C1'—C6'—C5'	2.09 (18)	C6'—C5'—C4'—C3'	-3.07 (18)
C2—C3—C4—C5	-1.01 (18)	C6—C5—N2—O3	170.62 (11)
C2—C3—N1—O1	7.99 (17)	C6'—C5'—N2'—O3'	178.23 (11)
C2—C3—N1—O2	-171.36 (11)	C6'—C5'—N2'—O4'	-1.94 (17)
C3—C2—C1—C1'	-174.87 (11)	C6—C5—N2—O4	-9.53 (16)
C3—C2—C1—C6	0.31 (18)	N1'—C3'—C2'—C1'	-174.56 (11)
C4'—C3'—C2'—C1'	2.08 (19)	N1—C3—C4—C5	-178.70 (10)
C4—C3—N1—O1	-174.20 (11)	N2'—C5'—C4'—C3'	179.39 (10)
C4—C3—N1—O2	6.44 (16)	N2—C5—C4—C3	179.81 (10)
C4'—C5'—C6'—C1'	1.44 (19)	N2'—C5'—C6'—C1'	178.95 (10)
C4'—C5'—N2'—O3'	-4.12 (16)	O1'—N1'—C3'—C2'	-20.79 (16)
C4—C5—N2—O3	-8.62 (16)	O1'—N1'—C3'—C4'	162.39 (11)
C4—C5—N2—O4	171.23 (11)	O2'—N1'—C3'—C2'	158.99 (11)
C4'—C5'—N2'—O4'	175.71 (11)	O2'—N1'—C3'—C4'	-17.83 (16)

Fig. 1




Fig. 2

