

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2,2'-Bipyridine)(2-{1-[2-(dimethyl-amino)ethylimino]ethyl}-4-methoxyphenolato)copper(II) perchlorate

Yueh-Hsuan Tsai, Wen-Chou Hung and Chu-Chieh Lin*

Department of Chemistry, National Chung Hsing University, Taichung 402, Taiwan, Republic of China

Correspondence e-mail: cchlin@mail.nchu.edu.tw


Received 24 March 2009; accepted 20 April 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.038; wR factor = 0.105; data-to-parameter ratio = 15.2.

The Cu atom of the title complex, $[\text{Cu}(\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2)(\text{C}_{10}\text{H}_8\text{N}_2)]\text{ClO}_4$, has a distorted square-pyramidal geometry with all three of the donor atoms from the N,N',O -tridentate Schiff base ligand in the equatorial positions and the bipyridine N atoms in an equatorial-axial binding mode. The Cu atom is 0.1801 (11) Å above the N_3O mean basal plane.

Related literature

For the development of efficient catalytic systems for the coupling of CO_2 with heterocycles into polycarbonates, see: Inoue *et al.* (1969). For the synthesis and catalytic studies of a series of *bis*-(salicylaldiminato)zinc complexes, see: Darensbourg *et al.* (2001). For similar complexes, see: Dhar *et al.* (2006); Shen *et al.* (2003). For the synthesis, see: Hung & Lin (2009); Hung *et al.* (2008); For the chemical activity of complexes, see: Noh *et al.* (2007).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2)(\text{C}_{10}\text{H}_8\text{N}_2)]\text{ClO}_4$ $V = 2472.8$ (4) Å³
 $M_r = 554.49$ $Z = 4$
 Monoclinic, $P2_1/c$ $\text{Mo } K\alpha$ radiation
 $a = 10.1588$ (10) Å $\mu = 1.04$ mm⁻¹
 $b = 18.2163$ (17) Å $T = 293$ K
 $c = 13.3764$ (13) Å $0.34 \times 0.26 \times 0.15$ mm
 $\beta = 92.610$ (2)°

Data collection

Bruker SMART 1000 CCD 13946 measured reflections
 diffractometer 4859 independent reflections
 Absorption correction: multi-scan 3488 reflections with $I > 2\sigma(I)$
 (SADABS; Sheldrick, 1996) $R_{\text{int}} = 0.037$
 $T_{\text{min}} = 0.719$, $T_{\text{max}} = 0.860$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$ 319 parameters
 $wR(F^2) = 0.105$ H-atom parameters constrained
 $S = 0.98$ $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 4859 reflections $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the National Science Council of the Republic of China is gratefully appreciated. Helpful comments from the reviewers are also greatly appreciated.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2135).

References

- Bruker (1999). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Darensbourg, D. J., Rainey, P. & Yarbrough, J. C. (2001). *Inorg. Chem.* **40**, 986–993.
 Dhar, S., Nethaji, M. & Chakravarty, A. R. (2006). *Inorg. Chem.* **45**, 11043–11050.
 Hung, W.-C., Hung, H. & Lin, C.-C. (2008). *J. Polym. Sci. Part A Polym. Chem.* **46**, 6466–6476.
 Hung, W.-C. & Lin, C.-C. (2009). *Inorg. Chem.* **48**, 728–734.
 Inoue, S., Koinuma, H. & Tsuruta, T. (1969). *Makromol. Chem.* **130**, 210–220.
 Noh, E. K., Na, S. J. S. S., Kim, S.-W. & Lee, B. Y. (2007). *J. Am. Chem. Soc.* **129**, 8082–8083.
 Sheldrick, G. M. (1996). *SADABS*, University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shen, Y. M., Duan, W. L. & Shi, M. (2003). *J. Org. Chem.* **68**, 1559–1562.

supplementary materials

Acta Cryst. (2009). E65, m578 [doi:10.1107/S1600536809014573]

(2,2'-Bipyridine)(2-{1-[2-(dimethylamino)ethylimino]ethyl}-4-methoxyphenolato)copper(II) perchlorate

Y.-H. Tsai, W.-C. Hung and C.-C. Lin

Comment

Though many bacteria convert CO₂ into organic compounds by photosynthesis, utilization of CO₂ as a chemical feedstock in industrial and laboratory is rare. Recently, reuse of CO₂ has received great attention because of environmental concern. Polycarbonates (PC) have been widely used in the modern chemical industry. Co-polymerization of CO₂ with olefins may benefit from reducing the release of CO₂ and generating potential industrial useful PCs. Therefore, there has been increasing interest in the development of efficient catalytic systems for the coupling of CO₂ with heterocycles into polycarbonates (Inoue *et al.*, 1969). One of the major successes is the utilization of epoxides and CO₂ as starting materials to prepare PCs and/or cyclic carbonates in the presence of a transition metal catalyst. Recently, Darensbourg *et al.*, (2001) disclosed the synthesis, characterization and catalytic studies of a series of *bis*-(salicylaldiminato)zinc complexes, in which the most active catalyst for co-polymerization of cyclohexene oxide and CO₂ giving poly(cyclohexene carbonate) (>99% carbonate linkages, $M_n = 41000 \text{ g mol}^{-1}$, $M_w/M_n = 10.3$) with a turnover frequency of 6.9 h^{-1} . In addition, Shen *et al.* (2003) reported that binaphthyl-diaminosalen-type Zn, Cu, and Co complexes efficiently catalyzed reactions of epoxides with CO₂ to achieve five-membered ring cyclic carbonates in the presence of various catalytic amounts of organic bases. Noh *et al.*, (2007) disclosed catalytic studies of the binary system of [(*salen*)Co(III)complex] / (quatarnary ammonium salt) for co-polymerization of propylene oxide and CO₂. Most recently, a series of *N,N,O*-tridentate Schiff base zinc- and magnesium-complexes have been reported to be effective initiators / catalyst for *ROP* of lactide (Hung *et al.*, 2008; Hung & Lin, 2009). We report herein the synthesis and crystal structure of [LCu(*bipy*)]ClO₄, where *L* is title tridentate ligand and *bipy* is 2,2'-bipyridine, a potential catalyst for CO₂ / epoxide coupling co-polymerization.

The solid structure of [LCu(*bipy*)]⁺ ion reveals a monomeric Cu^{II} complex containing a six-member and a five-member ring coordinated from the tridentate salicylideneimine ligand and a five-member ring coordinated from the bipyridine ligand. The geometry around Cu atom is penta-coordinated with a slight distorted square pyramidal environment in which all three of the *N,N,O*-tridentate donor atoms and one of the N atoms of the bipyridine ligand sitting on the equatorial plane, and another N atom of the bipyridine ligand at the axial position. The distances between the Cu atom and O1, N1, N2, N3 and N4 are 1.903 (2), 1.964 (2), 2.076 (2), 2.208 (2) and 2.044 (2) Å, respectively which are all within a normal distance for a Cu—O and Cu—N distance. These bond distances are similar to those found in other Schiff base Cu^{II} complexes (Dhar *et al.*, 2006).

Experimental

The ligand, 2-{1-[2-(dimethylamino)ethylimino]ethyl}-4-methoxyphenol was prepared according to the method reported previously (Hung *et al.*, 2008). The title complex was synthesized by the following procedures: Cu(OAc)₂·H₂O (0.197 g, 1.00 mmol) and 2,2'-bipyridine (0.199 g, 1.28 mmol) was stirred in *EtOH* (15 ml) at room temperature for 0.5 h. The

supplementary materials

2-{1-[2-(dimethylamino)ethylimino]ethyl}-4-methoxyphenol (0.298 g, 1.0 mmol) in *EtOH* (10 ml) was added. The reaction mixture was then stirred for another 1 h, and an 10 ml ethanolic solution of NaClO_4 (0.122 g, 1.0 mmol) was added producing green precipitate. The product was isolated by filtration and the resulting precipitate was crystallized from *EtOH* to yield green crystals.

Refinement

The methyl H atoms were located and then constrained to an ideal geometry with C—H distances of 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$, but each group was allowed to rotate freely about its C—C bond. All other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances in the range 0.93 Å and 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. A view of the molecular structure with the atom numbering scheme. The displacement ellipsoids are shown at the 20% probability level. H atoms are presented as a small spheres of arbitrary radius.

(2,2'-Bipyridine)(2-{1-[2-(dimethylamino)ethylimino]ethyl}-4-methoxyphenolato)copper(II) perchlorate

Crystal data

$[\text{Cu}(\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2)(\text{C}_{10}\text{H}_8\text{N}_2)]\text{ClO}_4$

$M_r = 554.49$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.1588$ (10) Å

$b = 18.2163$ (17) Å

$c = 13.3764$ (13) Å

$\beta = 92.610$ (2)°

$V = 2472.8$ (4) Å³

$Z = 4$

$F_{000} = 1148$

$D_x = 1.489$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 4710 reflections

$\theta = 2.3$ – 25.6 °

$\mu = 1.04$ mm⁻¹

$T = 293$ K

Parallelepiped, green

$0.34 \times 0.26 \times 0.15$ mm

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube

4859 independent reflections

3488 reflections with $I > 2\sigma(I)$

Monochromator: graphite $R_{\text{int}} = 0.037$
 $T = 293$ K $\theta_{\text{max}} = 26.0^\circ$
 φ and ω scans $\theta_{\text{min}} = 1.9^\circ$
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996) $h = -12 \rightarrow 12$
 $T_{\text{min}} = 0.719$, $T_{\text{max}} = 0.860$ $k = -22 \rightarrow 16$
 13946 measured reflections $l = -16 \rightarrow 15$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.038$ H-atom parameters constrained
 $wR(F^2) = 0.105$ $w = 1/[\sigma^2(F_o^2) + (0.06P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 0.98$ $(\Delta/\sigma)_{\text{max}} = 0.001$
 4859 reflections $\Delta\rho_{\text{max}} = 0.33 \text{ e } \text{\AA}^{-3}$
 319 parameters $\Delta\rho_{\text{min}} = -0.30 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	0.62390 (3)	0.172001 (16)	0.82971 (2)	0.04095 (12)
O1	0.6832 (2)	0.15346 (10)	0.96441 (14)	0.0532 (5)
O2	0.9049 (3)	-0.10697 (14)	1.0997 (2)	0.1017 (10)
N1	0.5494 (2)	0.07290 (12)	0.81586 (16)	0.0457 (5)
N2	0.4979 (2)	0.19843 (13)	0.70841 (18)	0.0528 (6)
N3	0.8094 (2)	0.16361 (12)	0.75071 (17)	0.0469 (6)
N4	0.6989 (2)	0.27569 (11)	0.84447 (15)	0.0412 (5)
C1	0.7293 (3)	0.08881 (15)	0.9916 (2)	0.0455 (6)
C2	0.8227 (3)	0.08568 (17)	1.0724 (2)	0.0583 (8)
H2A	0.8483	0.1293	1.1039	0.070*
C3	0.8779 (4)	0.02114 (19)	1.1069 (2)	0.0681 (9)

supplementary materials

H3A	0.9395	0.0215	1.1605	0.082*
C4	0.8411 (4)	-0.04469 (18)	1.0612 (2)	0.0665 (9)
C5	0.7497 (3)	-0.04455 (16)	0.9842 (2)	0.0576 (8)
H5A	0.7257	-0.0890	0.9544	0.069*
C6	0.6895 (3)	0.02080 (15)	0.94751 (19)	0.0439 (6)
C7	0.5849 (3)	0.01591 (15)	0.8687 (2)	0.0447 (7)
C8	0.5192 (3)	-0.05759 (16)	0.8497 (2)	0.0605 (8)
H8A	0.5193	-0.0850	0.9111	0.079 (10)*
H8B	0.5666	-0.0844	0.8011	0.102 (13)*
H8C	0.4300	-0.0500	0.8250	0.098 (13)*
C9	0.4456 (3)	0.06785 (18)	0.7363 (2)	0.0629 (8)
H9A	0.3599	0.0752	0.7639	0.075*
H9B	0.4469	0.0197	0.7054	0.075*
C10	0.4709 (3)	0.12658 (16)	0.6597 (2)	0.0603 (8)
H10A	0.5457	0.1125	0.6214	0.072*
H10B	0.3947	0.1309	0.6137	0.072*
C11	0.3779 (4)	0.2319 (2)	0.7440 (3)	0.0901 (13)
H11A	0.3197	0.2441	0.6879	0.135*
H11B	0.4002	0.2758	0.7809	0.135*
H11C	0.3349	0.1980	0.7867	0.135*
C12	0.5537 (4)	0.24885 (18)	0.6336 (2)	0.0728 (10)
H12A	0.4889	0.2579	0.5805	0.109*
H12B	0.6301	0.2267	0.6066	0.109*
H12C	0.5781	0.2944	0.6654	0.109*
C13	0.8909 (5)	-0.1720 (2)	1.0445 (3)	0.1034 (15)
H13A	0.9394	-0.2106	1.0783	0.155*
H13B	0.9244	-0.1648	0.9792	0.155*
H13C	0.7994	-0.1852	1.0380	0.155*
C14	0.8536 (3)	0.10711 (16)	0.6971 (2)	0.0578 (8)
H14A	0.8076	0.0630	0.6976	0.069*
C15	0.9634 (3)	0.11167 (19)	0.6418 (2)	0.0630 (9)
H15A	0.9903	0.0718	0.6045	0.076*
C16	1.0329 (3)	0.1763 (2)	0.6427 (3)	0.0689 (9)
H16A	1.1076	0.1809	0.6055	0.083*
C17	0.9908 (3)	0.23447 (17)	0.6995 (2)	0.0572 (8)
H17A	1.0379	0.2783	0.7021	0.069*
C18	0.8784 (3)	0.22671 (14)	0.75206 (19)	0.0415 (6)
C19	0.8210 (3)	0.28739 (14)	0.81221 (18)	0.0402 (6)
C20	0.8879 (3)	0.35225 (16)	0.8336 (2)	0.0531 (7)
H20A	0.9728	0.3592	0.8123	0.064*
C21	0.8265 (3)	0.40656 (17)	0.8872 (2)	0.0611 (8)
H21A	0.8702	0.4503	0.9025	0.073*
C22	0.7019 (3)	0.39554 (15)	0.9173 (2)	0.0546 (8)
H22A	0.6587	0.4319	0.9520	0.066*
C23	0.6409 (3)	0.32963 (14)	0.8954 (2)	0.0480 (7)
H23A	0.5561	0.3220	0.9166	0.058*
Cl	0.25347 (7)	0.09143 (4)	0.40216 (5)	0.05096 (19)
O3	0.2755 (3)	0.16261 (12)	0.4419 (2)	0.0957 (9)
O4	0.2085 (3)	0.09639 (16)	0.30025 (18)	0.0956 (9)

O5	0.1576 (3)	0.05517 (15)	0.4574 (2)	0.0958 (8)
O6	0.3729 (2)	0.05090 (16)	0.4085 (2)	0.0970 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu	0.0450 (2)	0.03477 (19)	0.04338 (19)	-0.00436 (14)	0.00530 (14)	-0.00633 (13)
O1	0.0719 (14)	0.0383 (11)	0.0488 (11)	-0.0068 (9)	-0.0037 (10)	-0.0082 (8)
O2	0.157 (3)	0.0660 (17)	0.0788 (17)	0.0307 (17)	-0.0360 (18)	0.0011 (13)
N1	0.0480 (14)	0.0429 (13)	0.0461 (12)	-0.0095 (11)	0.0028 (10)	-0.0070 (11)
N2	0.0570 (15)	0.0454 (14)	0.0554 (14)	0.0058 (12)	-0.0036 (12)	-0.0100 (11)
N3	0.0431 (13)	0.0430 (14)	0.0552 (14)	0.0014 (10)	0.0088 (11)	-0.0067 (10)
N4	0.0466 (13)	0.0356 (12)	0.0416 (12)	-0.0007 (10)	0.0033 (10)	-0.0018 (9)
C1	0.0533 (17)	0.0415 (16)	0.0424 (14)	-0.0072 (13)	0.0103 (12)	-0.0023 (12)
C2	0.071 (2)	0.0517 (19)	0.0519 (17)	-0.0082 (16)	-0.0031 (15)	-0.0078 (14)
C3	0.082 (2)	0.070 (2)	0.0515 (18)	-0.0007 (19)	-0.0059 (17)	-0.0052 (16)
C4	0.089 (3)	0.057 (2)	0.0532 (18)	0.0107 (18)	0.0023 (18)	-0.0003 (15)
C5	0.080 (2)	0.0413 (17)	0.0515 (17)	-0.0015 (15)	0.0065 (16)	-0.0040 (13)
C6	0.0526 (16)	0.0415 (15)	0.0383 (14)	-0.0049 (13)	0.0089 (12)	-0.0015 (11)
C7	0.0517 (16)	0.0389 (15)	0.0451 (15)	-0.0108 (13)	0.0191 (13)	-0.0092 (12)
C8	0.074 (2)	0.0480 (18)	0.0608 (19)	-0.0214 (16)	0.0146 (17)	-0.0062 (15)
C9	0.061 (2)	0.060 (2)	0.066 (2)	-0.0155 (16)	-0.0089 (16)	-0.0077 (16)
C10	0.070 (2)	0.0530 (19)	0.0569 (18)	0.0031 (16)	-0.0116 (16)	-0.0116 (14)
C11	0.063 (2)	0.103 (3)	0.103 (3)	0.028 (2)	-0.010 (2)	-0.035 (2)
C12	0.099 (3)	0.055 (2)	0.063 (2)	0.0052 (19)	-0.0126 (19)	0.0072 (16)
C13	0.149 (4)	0.068 (3)	0.092 (3)	0.037 (3)	-0.011 (3)	0.005 (2)
C14	0.0549 (18)	0.0486 (18)	0.070 (2)	0.0049 (14)	0.0082 (15)	-0.0145 (15)
C15	0.060 (2)	0.066 (2)	0.063 (2)	0.0147 (17)	0.0092 (16)	-0.0190 (16)
C16	0.0503 (19)	0.091 (3)	0.067 (2)	0.0089 (18)	0.0195 (16)	-0.0087 (18)
C17	0.0470 (17)	0.059 (2)	0.0658 (19)	-0.0041 (14)	0.0093 (15)	0.0006 (15)
C18	0.0399 (14)	0.0445 (15)	0.0398 (14)	0.0019 (12)	-0.0011 (11)	0.0024 (11)
C19	0.0429 (15)	0.0402 (15)	0.0371 (13)	-0.0018 (12)	-0.0026 (11)	0.0038 (11)
C20	0.0515 (18)	0.0513 (18)	0.0566 (17)	-0.0134 (14)	0.0023 (14)	-0.0014 (14)
C21	0.076 (2)	0.0430 (17)	0.0639 (19)	-0.0151 (16)	-0.0035 (17)	-0.0052 (14)
C22	0.077 (2)	0.0381 (16)	0.0488 (17)	0.0009 (15)	0.0025 (15)	-0.0054 (12)
C23	0.0541 (17)	0.0423 (16)	0.0480 (16)	0.0021 (13)	0.0060 (13)	-0.0046 (12)
Cl	0.0475 (4)	0.0508 (4)	0.0552 (4)	0.0010 (3)	0.0103 (3)	0.0000 (3)
O3	0.138 (3)	0.0527 (15)	0.097 (2)	-0.0073 (15)	0.0050 (18)	-0.0065 (13)
O4	0.0810 (17)	0.146 (3)	0.0593 (15)	0.0118 (17)	-0.0038 (13)	-0.0072 (15)
O5	0.0868 (18)	0.0864 (19)	0.118 (2)	-0.0077 (15)	0.0486 (16)	0.0223 (15)
O6	0.0599 (15)	0.113 (2)	0.118 (2)	0.0323 (15)	0.0115 (14)	0.0019 (17)

Geometric parameters (\AA , $^\circ$)

Cu—O1	1.9037 (19)	C9—H9B	0.9700
Cu—N1	1.963 (2)	C10—H10A	0.9700
Cu—N4	2.043 (2)	C10—H10B	0.9700
Cu—N2	2.077 (2)	C11—H11A	0.9600
Cu—N3	2.207 (2)	C11—H11B	0.9600

supplementary materials

O1—C1	1.313 (3)	C11—H11C	0.9600
O2—C4	1.393 (4)	C12—H12A	0.9600
O2—C13	1.400 (4)	C12—H12B	0.9600
N1—C7	1.298 (3)	C12—H12C	0.9600
N1—C9	1.466 (3)	C13—H13A	0.9600
N2—C11	1.462 (4)	C13—H13B	0.9600
N2—C10	1.482 (4)	C13—H13C	0.9600
N2—C12	1.490 (4)	C14—C15	1.369 (4)
N3—C14	1.343 (3)	C14—H14A	0.9300
N3—C18	1.346 (3)	C15—C16	1.372 (4)
N4—C19	1.348 (3)	C15—H15A	0.9300
N4—C23	1.347 (3)	C16—C17	1.383 (4)
C1—C2	1.407 (4)	C16—H16A	0.9300
C1—C6	1.423 (4)	C17—C18	1.375 (4)
C2—C3	1.373 (4)	C17—H17A	0.9300
C2—H2A	0.9300	C18—C19	1.501 (4)
C3—C4	1.389 (4)	C19—C20	1.386 (4)
C3—H3A	0.9300	C20—C21	1.387 (4)
C4—C5	1.355 (4)	C20—H20A	0.9300
C5—C6	1.416 (4)	C21—C22	1.360 (4)
C5—H5A	0.9300	C21—H21A	0.9300
C6—C7	1.465 (4)	C22—C23	1.376 (4)
C7—C8	1.512 (4)	C22—H22A	0.9300
C8—H8A	0.9600	C23—H23A	0.9300
C8—H8B	0.9600	Cl—O5	1.413 (2)
C8—H8C	0.9600	Cl—O3	1.416 (2)
C9—C10	1.512 (4)	Cl—O6	1.419 (2)
C9—H9A	0.9700	Cl—O4	1.421 (2)
O1—Cu—N1	91.73 (9)	N2—C10—C9	111.1 (3)
O1—Cu—N4	88.40 (8)	N2—C10—H10A	109.4
N1—Cu—N4	179.21 (9)	C9—C10—H10A	109.4
O1—Cu—N2	159.65 (9)	N2—C10—H10B	109.4
N1—Cu—N2	85.30 (9)	C9—C10—H10B	109.4
N4—Cu—N2	94.31 (9)	H10A—C10—H10B	108.0
O1—Cu—N3	101.58 (9)	N2—C11—H11A	109.5
N1—Cu—N3	103.01 (9)	N2—C11—H11B	109.5
N4—Cu—N3	77.72 (8)	H11A—C11—H11B	109.5
N2—Cu—N3	98.71 (9)	N2—C11—H11C	109.5
C1—O1—Cu	121.01 (16)	H11A—C11—H11C	109.5
C4—O2—C13	117.4 (3)	H11B—C11—H11C	109.5
C7—N1—C9	121.2 (2)	N2—C12—H12A	109.5
C7—N1—Cu	125.99 (18)	N2—C12—H12B	109.5
C9—N1—Cu	112.81 (18)	H12A—C12—H12B	109.5
C11—N2—C10	111.8 (3)	N2—C12—H12C	109.5
C11—N2—C12	108.1 (3)	H12A—C12—H12C	109.5
C10—N2—C12	108.5 (2)	H12B—C12—H12C	109.5
C11—N2—Cu	109.6 (2)	O2—C13—H13A	109.5
C10—N2—Cu	103.51 (18)	O2—C13—H13B	109.5
C12—N2—Cu	115.31 (19)	H13A—C13—H13B	109.5

C14—N3—C18	118.4 (2)	O2—C13—H13C	109.5
C14—N3—Cu	128.5 (2)	H13A—C13—H13C	109.5
C18—N3—Cu	112.95 (17)	H13B—C13—H13C	109.5
C19—N4—C23	118.5 (2)	N3—C14—C15	122.8 (3)
C19—N4—Cu	117.37 (17)	N3—C14—H14A	118.6
C23—N4—Cu	123.61 (19)	C15—C14—H14A	118.6
O1—C1—C2	118.0 (2)	C14—C15—C16	118.6 (3)
O1—C1—C6	125.1 (3)	C14—C15—H15A	120.7
C2—C1—C6	116.9 (3)	C16—C15—H15A	120.7
C3—C2—C1	122.9 (3)	C15—C16—C17	119.3 (3)
C3—C2—H2A	118.5	C15—C16—H16A	120.3
C1—C2—H2A	118.5	C17—C16—H16A	120.3
C2—C3—C4	119.6 (3)	C18—C17—C16	119.1 (3)
C2—C3—H3A	120.2	C18—C17—H17A	120.4
C4—C3—H3A	120.2	C16—C17—H17A	120.4
C5—C4—C3	119.6 (3)	N3—C18—C17	121.7 (2)
C5—C4—O2	125.0 (3)	N3—C18—C19	114.9 (2)
C3—C4—O2	115.4 (3)	C17—C18—C19	123.4 (3)
C4—C5—C6	122.4 (3)	N4—C19—C20	121.2 (2)
C4—C5—H5A	118.8	N4—C19—C18	116.2 (2)
C6—C5—H5A	118.8	C20—C19—C18	122.6 (2)
C5—C6—C1	118.5 (3)	C21—C20—C19	119.1 (3)
C5—C6—C7	119.1 (2)	C21—C20—H20A	120.4
C1—C6—C7	122.3 (2)	C19—C20—H20A	120.4
N1—C7—C6	121.2 (2)	C22—C21—C20	119.6 (3)
N1—C7—C8	120.5 (3)	C22—C21—H21A	120.2
C6—C7—C8	118.3 (3)	C20—C21—H21A	120.2
C7—C8—H8A	109.5	C21—C22—C23	118.8 (3)
C7—C8—H8B	109.5	C21—C22—H22A	120.6
H8A—C8—H8B	109.5	C23—C22—H22A	120.6
C7—C8—H8C	109.5	N4—C23—C22	122.8 (3)
H8A—C8—H8C	109.5	N4—C23—H23A	118.6
H8B—C8—H8C	109.5	C22—C23—H23A	118.6
N1—C9—C10	108.0 (2)	O5—Cl—O3	109.42 (18)
N1—C9—H9A	110.1	O5—Cl—O6	109.47 (17)
C10—C9—H9A	110.1	O3—Cl—O6	109.50 (18)
N1—C9—H9B	110.1	O5—Cl—O4	109.41 (17)
C10—C9—H9B	110.1	O3—Cl—O4	109.93 (18)
H9A—C9—H9B	108.4	O6—Cl—O4	109.10 (17)

