

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dichlorido{2-[(4-iodophenyl)imino-methyl]pyridine- κ^2N,N' }copper(II)

 Ali Mahmoudi,^{a*} Mehdi Khalaj,^b Shan Gao,^c Seik Weng Ng^d and Mahmoud Mohammadgholiha^a
^aDepartment of Chemistry, Islamic Azad University, Karaj Branch, Karaj, Iran,

^bDepartment of Chemistry, Islamic Azad University, Buinzahra Branch, Qazvin, Iran,

^cSchool of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: mahmoudi_ali@yahoo.com


Received 22 March 2009; accepted 30 March 2009

 Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.028; wR factor = 0.079; data-to-parameter ratio = 19.4.

The Cu^{II} atom in the title complex, [CuCl₂(C₁₂H₉IN₂)], has a square-planar coordination being N,N' -chelated by the Schiff base ligand, and by two Cl atoms. The geometry is distorted towards square pyramidal owing to a long Cu...Cl interaction of 2.941 (1) Å. This results in the formation of a zigzag chain structure propagating in the c -axis direction.

Related literature

For background to the synthesis and structure of metal complexes of diimines, see: Yamada (1999). For the structure of the zinc chloride complex of the same ligand, see: Dehghanpour *et al.* (2007).



Experimental

Crystal data

 [CuCl₂(C₁₂H₉IN₂)]
 $M_r = 442.55$

 Monoclinic, $P2_1/c$
 $a = 12.2721$ (5) Å

 $b = 15.2159$ (5) Å

 $c = 7.4709$ (2) Å

 $\beta = 94.8913$ (10)°

 $V = 1389.97$ (8) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 4.16$ mm⁻¹
 $T = 295$ K

 $0.31 \times 0.25 \times 0.17$ mm

Data collection

 Rigaku RAXIS-RAPID
 diffractometer

 Absorption correction: multi-scan
 (ABSCOR; Higashi, 1995)

 $T_{\min} = 0.359$, $T_{\max} = 0.538$

(expected range = 0.329–0.493)

21713 measured reflections

3166 independent reflections

 2650 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.079$
 $S = 1.05$

3166 reflections

163 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.88$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.51$ e Å⁻³

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *SHELXL97*.

We thank the Islamic Azad University Research Council and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2104).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Dehghanpour, S., Mahmoudi, A., Khalaj, M., Salmanpour, S. & Adib, M. (2007). *Acta Cryst.* **E63**, m2841.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yamada, S. (1999). *Coord. Chem. Rev.* **190**, 537–555.

supplementary materials

Acta Cryst. (2009). E65, m555 [doi:10.1107/S1600536809011684]

Dichlorido{2-[(4-iodophenyl)iminomethyl]pyridine- κ^2N,N' }copper(II)

A. Mahmoudi, M. Khalaj, S. Gao, S. W. Ng and M. Mohammadgholiha

Comment

The structure of the title complex is illustrated in Fig. 1, and the geometrical parameters are available in the archived CIF. The complex was prepared by the reaction of copper chloride with the Schiff base ligand 2-[(4-iodophenyl)iminomethyl]pyridine (Yamada, 1999). The ligand is slightly twisted with the benzene ring and pyridine ring being inclined to one another by 26.07813° .

A long $\text{Cu}\cdots\text{Cl}2^i$ [symmetry operation (i) = x, 1.5-y, 0.5+z] interaction of 2.941 (1) Å, leads the formation of a zigzag chain propagating in the c direction (Fig. 2). This situation is different to that observed in the ZnCl_2 complex of the same ligand, which is mononuclear (Dehghanpour *et al.*, 2007).

Experimental

To a solution of 2-[(4-iodophenyl)iminomethyl]pyridine (30.8 mg, 0.1 mmol) in 20 ml acetonitrile was added copper(II) chloride (13.4 mg, 0.1 mmol). The mixture was heated to dissolve the reactants and then the solution was filtered and the volume reduced under vacuum to ca. 5 ml. Diffusion of diethyl ether vapor into the solution gave green crystals of the title complex. The crystals were collected and washed with diethylether-dichloromethane (9:1 v/v). Yield 75%. CHN elemental analysis: Calc. for $\text{C}_{12}\text{H}_9\text{Cl}_2\text{CuIN}_2$: C 32.57, H 2.05, N 6.33%; found: C 32.54, H 2.07, N 6.35%.

Refinement

The H-atoms were placed in calculated positions [C—H 0.93 Å] and treated as riding atoms [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$].

Figures




Fig. 1. A view of the structure of the title complex, with displacement ellipsoids drawn at the 50% probability level [H atoms are represented as spheres of arbitrary radius].




Fig. 2. A view of the $\text{Cu}\cdots\text{Cl}2^i$ bridged zigzag bridged chain structure of the title compound [Symmetry operations (i) = x, 1.5-y, -0.5+z]

Dichlorido[2-[(4-iodophenyl)iminomethyl]pyridine- κ^2N,N']copper(II)

Crystal data

[CuCl ₂ (C ₁₂ H ₉ IN ₂)]	$F_{000} = 844$
$M_r = 442.55$	$D_x = 2.115 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 12.2721 (5) \text{ \AA}$	Cell parameters from 13838 reflections
$b = 15.2159 (5) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$c = 7.4709 (2) \text{ \AA}$	$\mu = 4.16 \text{ mm}^{-1}$
$\beta = 94.8913 (10)^\circ$	$T = 295 \text{ K}$
$V = 1389.97 (8) \text{ \AA}^3$	Block, green
$Z = 4$	$0.31 \times 0.25 \times 0.17 \text{ mm}$

Data collection

Rigaku RAXIS-RAPID diffractometer	3166 independent reflections
Radiation source: fine-focus sealed tube	2650 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.037$
Detector resolution: $10.000 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.5^\circ$
$T = 295 \text{ K}$	$\theta_{\text{min}} = 3.1^\circ$
ω scans	$h = -15 \rightarrow 15$
Absorption correction: Multi-scan (ABSCOR; Higashi, 1995)	$k = -19 \rightarrow 19$
$T_{\text{min}} = 0.359$, $T_{\text{max}} = 0.538$	$l = -9 \rightarrow 9$
21713 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.079$	$w = 1/[\sigma^2(F_o^2) + (0.0447P)^2 + 0.6865P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
3166 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
163 parameters	$\Delta\rho_{\text{max}} = 0.88 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.51 \text{ e \AA}^{-3}$
	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	-0.001646 (16)	0.649922 (16)	0.55181 (3)	0.05798 (10)
Cu1	0.60873 (3)	0.72391 (2)	0.87653 (5)	0.03844 (11)
Cl1	0.72032 (6)	0.82416 (5)	1.01818 (12)	0.05137 (19)
Cl2	0.51519 (6)	0.82558 (5)	0.71178 (11)	0.04616 (18)
N1	0.70612 (18)	0.62040 (15)	0.9438 (3)	0.0375 (5)
N2	0.50252 (17)	0.62295 (15)	0.7935 (3)	0.0334 (4)
C1	0.8056 (2)	0.6212 (2)	1.0289 (4)	0.0469 (7)
H1	0.8344	0.6738	1.0752	0.056*
C2	0.8676 (2)	0.5448 (2)	1.0503 (5)	0.0520 (8)
H2	0.9368	0.5463	1.1114	0.062*
C3	0.8259 (3)	0.4672 (2)	0.9809 (5)	0.0489 (7)
H3	0.8677	0.4162	0.9895	0.059*
C4	0.7210 (3)	0.4657 (2)	0.8980 (4)	0.0443 (6)
H4	0.6902	0.4135	0.8530	0.053*
C5	0.6629 (2)	0.54354 (18)	0.8832 (4)	0.0365 (6)
C6	0.5499 (2)	0.54841 (18)	0.8064 (4)	0.0378 (6)
H6	0.5128	0.4977	0.7672	0.045*
C7	0.3893 (2)	0.62851 (18)	0.7297 (4)	0.0350 (5)
C8	0.3398 (2)	0.5679 (2)	0.6095 (4)	0.0446 (6)
H8	0.3814	0.5233	0.5643	0.054*
C9	0.2287 (2)	0.5736 (2)	0.5565 (4)	0.0479 (7)
H9	0.1957	0.5323	0.4775	0.057*
C10	0.1677 (2)	0.64099 (19)	0.6219 (4)	0.0417 (6)
C11	0.2166 (2)	0.70219 (19)	0.7420 (4)	0.0422 (6)
H11	0.1753	0.7475	0.7852	0.051*
C12	0.3266 (2)	0.69542 (18)	0.7964 (4)	0.0385 (6)
H12	0.3591	0.7357	0.8781	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.02714 (12)	0.06908 (18)	0.07514 (18)	0.00149 (8)	-0.01061 (10)	0.00131 (11)
Cu1	0.02850 (18)	0.03088 (18)	0.0536 (2)	0.00103 (12)	-0.00989 (14)	-0.00045 (14)
Cl1	0.0377 (4)	0.0440 (4)	0.0702 (5)	-0.0064 (3)	-0.0082 (3)	-0.0106 (4)
Cl2	0.0382 (4)	0.0366 (3)	0.0624 (4)	0.0042 (3)	-0.0032 (3)	0.0115 (3)
N1	0.0270 (11)	0.0373 (11)	0.0472 (12)	0.0010 (9)	-0.0036 (9)	0.0017 (11)
N2	0.0265 (11)	0.0321 (10)	0.0404 (11)	0.0000 (8)	-0.0033 (8)	0.0008 (9)
C1	0.0306 (14)	0.0500 (16)	0.0578 (17)	0.0032 (12)	-0.0087 (12)	-0.0016 (15)
C2	0.0279 (14)	0.0624 (19)	0.0640 (19)	0.0063 (13)	-0.0055 (13)	0.0086 (16)
C3	0.0358 (16)	0.0470 (16)	0.0640 (18)	0.0124 (12)	0.0054 (13)	0.0129 (15)
C4	0.0390 (15)	0.0367 (14)	0.0570 (17)	0.0040 (11)	0.0022 (13)	0.0056 (13)
C5	0.0287 (13)	0.0353 (13)	0.0449 (14)	0.0019 (10)	-0.0002 (10)	0.0041 (12)
C6	0.0315 (13)	0.0325 (13)	0.0484 (15)	-0.0023 (10)	-0.0027 (11)	0.0010 (12)
C7	0.0264 (12)	0.0349 (12)	0.0429 (13)	0.0000 (10)	-0.0022 (10)	0.0020 (12)

supplementary materials

C8	0.0304 (14)	0.0513 (17)	0.0513 (16)	0.0015 (12)	-0.0014 (11)	-0.0108 (14)
C9	0.0331 (15)	0.0581 (18)	0.0507 (16)	-0.0033 (13)	-0.0069 (12)	-0.0096 (15)
C10	0.0244 (13)	0.0492 (16)	0.0503 (16)	-0.0011 (11)	-0.0046 (11)	0.0049 (13)
C11	0.0320 (14)	0.0387 (14)	0.0556 (16)	0.0054 (11)	0.0017 (11)	0.0012 (13)
C12	0.0326 (14)	0.0360 (13)	0.0457 (14)	-0.0010 (11)	-0.0036 (11)	-0.0015 (12)

Geometric parameters (Å, °)

Il—C10	2.104 (3)	C4—C5	1.383 (4)
Cu1—N1	2.015 (2)	C4—H4	0.9300
Cu1—N2	2.075 (2)	C5—C6	1.457 (4)
Cu1—Cl1	2.253 (1)	C6—H6	0.9300
Cu1—Cl2	2.233 (1)	C7—C8	1.390 (4)
N1—C1	1.328 (4)	C7—C12	1.393 (4)
N1—C5	1.346 (4)	C8—C9	1.389 (4)
N2—C6	1.274 (4)	C8—H8	0.9300
N2—C7	1.433 (3)	C9—C10	1.384 (4)
C1—C2	1.391 (4)	C9—H9	0.9300
C1—H1	0.9300	C10—C11	1.393 (4)
C2—C3	1.371 (5)	C11—C12	1.380 (4)
C2—H2	0.9300	C11—H11	0.9300
C3—C4	1.380 (4)	C12—H12	0.9300
C3—H3	0.9300		
N1—Cu1—N2	80.79 (9)	N1—C5—C4	122.1 (3)
N1—Cu1—Cl2	160.96 (7)	N1—C5—C6	115.0 (2)
N1—Cu1—Cl1	95.07 (7)	C4—C5—C6	122.8 (3)
N2—Cu1—Cl1	169.40 (7)	N2—C6—C5	119.2 (2)
N2—Cu1—Cl2	93.89 (6)	N2—C6—H6	120.4
Cl1—Cu2—Cl1	93.05 (3)	C5—C6—H6	120.4
C1—N1—C5	119.3 (2)	C8—C7—C12	119.5 (2)
C1—N1—Cu1	127.9 (2)	C8—C7—N2	122.2 (2)
C5—N1—Cu1	112.76 (17)	C12—C7—N2	118.3 (2)
C6—N2—C7	120.1 (2)	C7—C8—C9	120.5 (3)
C6—N2—Cu1	111.44 (18)	C7—C8—H8	119.8
C7—N2—Cu1	128.50 (17)	C9—C8—H8	119.8
N1—C1—C2	121.2 (3)	C10—C9—C8	119.5 (3)
N1—C1—H1	119.4	C10—C9—H9	120.3
C2—C1—H1	119.4	C8—C9—H9	120.3
C3—C2—C1	119.6 (3)	C9—C10—C11	120.4 (3)
C3—C2—H2	120.2	C9—C10—H1	120.9 (2)
C1—C2—H2	120.2	C11—C10—H1	118.7 (2)
C2—C3—C4	119.2 (3)	C12—C11—C10	119.8 (3)
C2—C3—H3	120.4	C12—C11—H11	120.1
C4—C3—H3	120.4	C10—C11—H11	120.1
C3—C4—C5	118.5 (3)	C11—C12—C7	120.3 (3)
C3—C4—H4	120.8	C11—C12—H12	119.8
C5—C4—H4	120.8	C7—C12—H12	119.8
N2—Cu1—N1—C1	-175.5 (3)	C3—C4—C5—N1	1.4 (5)
Cl2—Cu1—N1—C1	109.5 (3)	C3—C4—C5—C6	-176.8 (3)

Cl1—Cu1—N1—C1	-5.4 (3)	C7—N2—C6—C5	-175.7 (2)
N2—Cu1—N1—C5	8.14 (19)	Cu1—N2—C6—C5	3.6 (3)
Cl2—Cu1—N1—C5	-66.8 (3)	N1—C5—C6—N2	3.3 (4)
Cl1—Cu1—N1—C5	178.30 (18)	C4—C5—C6—N2	-178.3 (3)
N1—Cu1—N2—C6	-6.3 (2)	C6—N2—C7—C8	-29.4 (4)
Cl2—Cu1—N2—C6	155.25 (19)	Cu1—N2—C7—C8	151.3 (2)
Cl1—Cu1—N2—C6	-74.0 (4)	C6—N2—C7—C12	148.2 (3)
N1—Cu1—N2—C7	172.9 (2)	Cu1—N2—C7—C12	-31.0 (3)
Cl2—Cu1—N2—C7	-25.5 (2)	C12—C7—C8—C9	-0.2 (4)
Cl1—Cu1—N2—C7	105.3 (4)	N2—C7—C8—C9	177.4 (3)
C5—N1—C1—C2	2.6 (5)	C7—C8—C9—C10	1.1 (5)
Cu1—N1—C1—C2	-173.5 (2)	C8—C9—C10—C11	-0.8 (5)
N1—C1—C2—C3	0.6 (5)	C8—C9—C10—I1	-178.1 (2)
C1—C2—C3—C4	-2.9 (5)	C9—C10—C11—C12	-0.2 (5)
C2—C3—C4—C5	1.9 (5)	I1—C10—C11—C12	177.1 (2)
C1—N1—C5—C4	-3.7 (4)	C10—C11—C12—C7	1.0 (4)
Cu1—N1—C5—C4	173.0 (2)	C8—C7—C12—C11	-0.8 (4)
C1—N1—C5—C6	174.7 (3)	N2—C7—C12—C11	-178.6 (3)
Cu1—N1—C5—C6	-8.6 (3)		

Fig. 1




Fig. 2

