

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## N-(3-Bromophenyl)acetamide

 B. Thimme Gowda,<sup>a\*</sup> Sabine Foro,<sup>b</sup> Hiromitsu Terao<sup>c</sup> and Hartmut Fuess<sup>b</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany, and <sup>c</sup>Faculty of Integrated Arts and Sciences, Tokushima University, Minamijosanjima-cho, Tokushima 770-8502, Japan  
Correspondence e-mail: gowdabt@yahoo.com


Received 6 April 2009; accepted 7 April 2009

Key indicators: single-crystal X-ray study;  $T = 299$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.093; data-to-parameter ratio = 17.2.

The conformation of the N—H bond in the structure of the title compound,  $\text{C}_8\text{H}_8\text{BrNO}$ , is *anti* to the C=O bond and to the *meta*-bromo substituent of the aromatic ring in both independent molecules comprising the asymmetric unit. Molecules are linked through N—H...O hydrogen bonding into supramolecular chains with a twisted topology.

## Related literature

For the preparation of the compound, see: Gowda *et al.* (2006). For related structures, see: Gowda *et al.* (2007, 2008, 2009).



## Experimental

## Crystal data

 $\text{C}_8\text{H}_8\text{BrNO}$ 
 $M_r = 214.06$ 

Orthorhombic,  $P2_12_12_1$   
 $a = 4.7836$  (6) Å  
 $b = 18.765$  (1) Å  
 $c = 19.379$  (2) Å  
 $V = 1739.5$  (3) Å<sup>3</sup>

$Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 4.67$  mm<sup>-1</sup>  
 $T = 299$  K  
 $0.44 \times 0.10 \times 0.08$  mm

## Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector  
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2007  
 $T_{\min} = 0.226$ ,  $T_{\max} = 0.685$   
9612 measured reflections  
3449 independent reflections  
2043 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.093$   
 $S = 0.99$   
3449 reflections  
201 parameters  
H-atom parameters constrained

$\Delta\rho_{\max} = 0.31$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.49$  e Å<sup>-3</sup>  
Absolute structure: Flack (1983), 1366 Friedel pairs  
Flack parameter:  $-0.008$  (13)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.86	2.05	2.887 (5)	166
$\text{N2}-\text{H2N}\cdots\text{O1}$	0.86	2.10	2.953 (5)	169

 Symmetry code: (i)  $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2004); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2007); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2417).

## References

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
Gowda, B. T., Foro, S. & Fuess, H. (2008). *Acta Cryst.* **E64**, o381.  
Gowda, B. T., Foro, S., Terao, H. & Fuess, H. (2009). *Acta Cryst.* **E65**, o949.  
Gowda, B. T., Shilpa & Lakshmiathy, J. K. (2006). *Z. Naturforsch. Teil A*, **61**, 595–599.  
Gowda, B. T., Svoboda, I. & Fuess, H. (2007). *Acta Cryst.* **E63**, o3267.  
Oxford Diffraction (2004). *CrysAlis CCD*. Oxford Diffraction Ltd, Köln, Germany.  
Oxford Diffraction (2007). *CrysAlis RED*. Oxford Diffraction Ltd, Köln, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1039 [ doi:10.1107/S1600536809013294 ]

## ***N*-(3-Bromophenyl)acetamide**

**B. T. Gowda, S. Foro, H. Terao and H. Fuess**

### **Comment**

As part of a study of the effect of ring and side-chain substitutions on the crystal structures of aromatic amides (Gowda *et al.*, 2007, 2008, 2009), in the present work, the structure of *N*-(3-bromophenyl)acetamide (I) has been determined. The conformation of the N—H bond in the structure is *anti* to the *meta*-bromo substituent of the aromatic ring (Fig. 1), in both independent molecules comprising the asymmetric unit, similar to that observed in *N*-(3-chlorophenyl)acetamide (Gowda *et al.*, 2008). Further, the conformation of the C=O bond is *anti* to the N—H bond. The two independent molecules in (I) are linked through intermolecular N—H $\cdots$ O hydrogen bonding into a supramolecular chains with a twisted topology (Table 1, Fig. 2).

### **Experimental**

Compound (I) was prepared according to the literature method (Gowda *et al.*, 2006). Single crystals were obtained from an ethanolic solution of (I).

### **Refinement**

The H atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å and N—H = 0.86 Å, and with  $U_{\text{iso}}$  set to 1.2 times  $U_{\text{eq}}$ (parent atom).

### **Figures**




Fig. 1. Molecular structure of (I), showing the atom labeling scheme and displacement ellipsoids drawn at the 50% probability level. The intermolecular N—H $\cdots$ O hydrogen is shown as a dashed line.




Fig. 2. Molecular packing of (I) viewed in projection down the *a*-axis highlighting the supramolecular chains mediated by hydrogen bonding (shown as dashed lines).

## *N*-(3-Bromophenyl)acetamide

### *Crystal data*

$C_8H_8BrNO$	$F_{000} = 848$
$M_r = 214.06$	$D_x = 1.635 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 4.7836 (6) \text{ \AA}$	Cell parameters from 3601 reflections
$b = 18.765 (1) \text{ \AA}$	$\theta = 2.4\text{--}27.6^\circ$
$c = 19.379 (2) \text{ \AA}$	$\mu = 4.67 \text{ mm}^{-1}$
$V = 1739.5 (3) \text{ \AA}^3$	$T = 299 \text{ K}$
$Z = 8$	Long needle, colourless
	$0.44 \times 0.10 \times 0.08 \text{ mm}$

### *Data collection*

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector	3449 independent reflections
Radiation source: fine-focus sealed tube	2043 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.034$
$T = 299 \text{ K}$	$\theta_{\text{max}} = 26.4^\circ$
Rotation method data acquisition using $\omega$ and $\varphi$ scans	$\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2007)	$h = -5 \rightarrow 5$
$T_{\text{min}} = 0.226$ , $T_{\text{max}} = 0.685$	$k = -22 \rightarrow 23$
9612 measured reflections	$l = -20 \rightarrow 24$

### *Refinement*

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.045$	$w = 1/[\sigma^2(F_o^2) + (0.0455P)^2 + 0.0252P]$
$wR(F^2) = 0.093$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.99$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3449 reflections	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
201 parameters	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1366 Friedel pairs
	Flack parameter: $-0.008 (13)$

### *Special details*

**Experimental.** Absorption correction: CrysAlis RED, Oxford Diffraction Ltd. (2007). Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.64658 (17)	0.36438 (3)	0.40336 (3)	0.0961 (3)
O1	1.0611 (8)	0.22271 (18)	0.20168 (19)	0.0786 (12)
N1	0.9531 (7)	0.13129 (19)	0.27225 (19)	0.0531 (10)
H1N	0.9782	0.0863	0.2779	0.064*
C1	0.7804 (10)	0.1647 (2)	0.3206 (2)	0.0492 (12)
C2	0.7903 (10)	0.2377 (2)	0.3338 (2)	0.0524 (12)
H2	0.9111	0.2670	0.3092	0.063*
C3	0.6192 (12)	0.2656 (3)	0.3838 (2)	0.0606 (14)
C4	0.4398 (12)	0.2247 (4)	0.4216 (3)	0.0740 (17)
H4	0.3269	0.2450	0.4553	0.089*
C5	0.4300 (12)	0.1519 (3)	0.4084 (3)	0.0848 (18)
H5	0.3068	0.1234	0.4331	0.102*
C6	0.6000 (11)	0.1215 (3)	0.3592 (3)	0.0691 (15)
H6	0.5951	0.0726	0.3516	0.083*
C7	1.0850 (10)	0.1608 (3)	0.2180 (2)	0.0546 (13)
C8	1.2664 (11)	0.1102 (2)	0.1767 (2)	0.0726 (17)
H8A	1.2240	0.0620	0.1895	0.087*
H8B	1.4597	0.1199	0.1861	0.087*
H8C	1.2303	0.1166	0.1284	0.087*
Br2	0.30944 (16)	0.49664 (3)	-0.03009 (3)	0.0919 (3)
O2	0.9050 (8)	0.48750 (17)	0.18809 (18)	0.0709 (10)
N2	0.8933 (8)	0.36972 (18)	0.16644 (18)	0.0475 (9)
H2N	0.9634	0.3294	0.1785	0.057*
C9	0.6865 (9)	0.3675 (2)	0.11456 (19)	0.0397 (10)
C10	0.6162 (10)	0.4252 (2)	0.0737 (2)	0.0464 (11)
H10	0.7039	0.4690	0.0799	0.056*
C11	0.4147 (10)	0.4168 (2)	0.0239 (2)	0.0499 (12)
C12	0.2819 (9)	0.3533 (3)	0.0127 (2)	0.0528 (12)
H12	0.1476	0.3489	-0.0218	0.063*
C13	0.3519 (12)	0.2959 (3)	0.0536 (2)	0.0576 (13)
H13	0.2619	0.2525	0.0473	0.069*
C14	0.5532 (9)	0.3025 (2)	0.1035 (2)	0.0490 (12)
H14	0.6013	0.2632	0.1302	0.059*
C15	0.9939 (9)	0.4277 (3)	0.1993 (2)	0.0490 (12)
C16	1.2135 (10)	0.4133 (3)	0.2521 (3)	0.0652 (14)

## supplementary materials

---

H16A	1.2663	0.3640	0.2502	0.078*
H16B	1.1416	0.4242	0.2971	0.078*
H16C	1.3739	0.4426	0.2429	0.078*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.1400 (6)	0.0715 (4)	0.0768 (4)	0.0319 (4)	-0.0121 (4)	-0.0259 (3)
O1	0.110 (3)	0.046 (2)	0.080 (2)	0.013 (2)	0.022 (2)	0.0251 (19)
N1	0.067 (3)	0.035 (2)	0.057 (2)	0.0055 (19)	0.003 (2)	0.012 (2)
C1	0.058 (3)	0.045 (3)	0.044 (3)	0.005 (2)	-0.005 (2)	0.006 (2)
C2	0.065 (3)	0.045 (3)	0.047 (2)	0.004 (2)	-0.003 (3)	0.003 (2)
C3	0.075 (4)	0.064 (3)	0.043 (3)	0.021 (3)	-0.007 (3)	-0.006 (3)
C4	0.068 (4)	0.108 (5)	0.046 (3)	0.021 (3)	0.008 (3)	-0.009 (3)
C5	0.089 (5)	0.096 (5)	0.070 (4)	-0.020 (3)	0.014 (4)	0.002 (4)
C6	0.083 (4)	0.059 (3)	0.065 (3)	-0.009 (3)	0.003 (3)	0.009 (3)
C7	0.067 (4)	0.045 (3)	0.052 (3)	0.007 (2)	0.002 (3)	0.008 (2)
C8	0.086 (5)	0.065 (3)	0.066 (3)	0.005 (3)	0.012 (3)	0.002 (3)
Br2	0.1222 (5)	0.0670 (4)	0.0866 (4)	0.0074 (4)	-0.0273 (4)	0.0277 (3)
O2	0.091 (3)	0.0351 (18)	0.086 (2)	0.0039 (19)	-0.014 (2)	-0.0108 (17)
N2	0.058 (3)	0.0299 (19)	0.054 (2)	0.0034 (19)	0.000 (2)	0.0001 (18)
C9	0.044 (3)	0.035 (2)	0.040 (2)	0.000 (2)	0.004 (2)	-0.003 (2)
C10	0.051 (3)	0.035 (2)	0.053 (3)	-0.002 (2)	-0.001 (3)	0.005 (2)
C11	0.059 (3)	0.044 (3)	0.047 (3)	0.008 (2)	0.003 (3)	0.007 (2)
C12	0.048 (3)	0.058 (3)	0.052 (3)	-0.002 (2)	-0.003 (2)	-0.006 (2)
C13	0.056 (3)	0.051 (3)	0.066 (3)	-0.002 (3)	-0.007 (3)	-0.011 (3)
C14	0.053 (3)	0.035 (3)	0.058 (3)	0.000 (2)	0.006 (3)	-0.002 (2)
C15	0.050 (3)	0.047 (3)	0.050 (3)	-0.001 (3)	-0.001 (3)	-0.007 (3)
C16	0.065 (4)	0.062 (3)	0.069 (3)	-0.009 (3)	-0.007 (3)	-0.012 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Br1—C3	1.897 (5)	Br2—C11	1.895 (4)
O1—C7	1.209 (5)	O2—C15	1.220 (5)
N1—C7	1.346 (5)	N2—C15	1.349 (5)
N1—C1	1.398 (5)	N2—C9	1.411 (5)
N1—H1N	0.8600	N2—H2N	0.8600
C1—C2	1.394 (6)	C9—C10	1.383 (5)
C1—C6	1.400 (6)	C9—C14	1.392 (6)
C2—C3	1.372 (6)	C10—C11	1.374 (6)
C2—H2	0.9300	C10—H10	0.9300
C3—C4	1.364 (7)	C11—C12	1.369 (6)
C4—C5	1.390 (7)	C12—C13	1.377 (6)
C4—H4	0.9300	C12—H12	0.9300
C5—C6	1.377 (7)	C13—C14	1.371 (6)
C5—H5	0.9300	C13—H13	0.9300
C6—H6	0.9300	C14—H14	0.9300
C7—C8	1.514 (6)	C15—C16	1.490 (6)
C8—H8A	0.9600	C16—H16A	0.9600

C8—H8B	0.9600	C16—H16B	0.9600
C8—H8C	0.9600	C16—H16C	0.9600
C7—N1—C1	128.1 (4)	C15—N2—C9	127.6 (4)
C7—N1—H1N	116.0	C15—N2—H2N	116.2
C1—N1—H1N	116.0	C9—N2—H2N	116.2
C2—C1—N1	123.0 (4)	C10—C9—C14	119.1 (4)
C2—C1—C6	119.5 (5)	C10—C9—N2	123.7 (4)
N1—C1—C6	117.5 (4)	C14—C9—N2	117.2 (4)
C3—C2—C1	119.0 (5)	C11—C10—C9	118.9 (4)
C3—C2—H2	120.5	C11—C10—H10	120.5
C1—C2—H2	120.5	C9—C10—H10	120.5
C4—C3—C2	122.7 (5)	C12—C11—C10	122.5 (4)
C4—C3—Br1	119.1 (4)	C12—C11—Br2	118.5 (4)
C2—C3—Br1	118.2 (4)	C10—C11—Br2	119.0 (3)
C3—C4—C5	118.3 (5)	C11—C12—C13	118.4 (4)
C3—C4—H4	120.8	C11—C12—H12	120.8
C5—C4—H4	120.8	C13—C12—H12	120.8
C6—C5—C4	121.0 (5)	C14—C13—C12	120.4 (4)
C6—C5—H5	119.5	C14—C13—H13	119.8
C4—C5—H5	119.5	C12—C13—H13	119.8
C5—C6—C1	119.6 (5)	C13—C14—C9	120.6 (4)
C5—C6—H6	120.2	C13—C14—H14	119.7
C1—C6—H6	120.2	C9—C14—H14	119.7
O1—C7—N1	123.7 (4)	O2—C15—N2	122.2 (4)
O1—C7—C8	121.3 (4)	O2—C15—C16	122.3 (4)
N1—C7—C8	115.0 (4)	N2—C15—C16	115.5 (4)
C7—C8—H8A	109.5	C15—C16—H16A	109.5
C7—C8—H8B	109.5	C15—C16—H16B	109.5
H8A—C8—H8B	109.5	H16A—C16—H16B	109.5
C7—C8—H8C	109.5	C15—C16—H16C	109.5
H8A—C8—H8C	109.5	H16A—C16—H16C	109.5
H8B—C8—H8C	109.5	H16B—C16—H16C	109.5
C7—N1—C1—C2	-22.3 (7)	C15—N2—C9—C10	21.5 (6)
C7—N1—C1—C6	160.8 (4)	C15—N2—C9—C14	-160.2 (4)
N1—C1—C2—C3	-177.9 (4)	C14—C9—C10—C11	0.7 (6)
C6—C1—C2—C3	-1.0 (7)	N2—C9—C10—C11	178.9 (4)
C1—C2—C3—C4	0.4 (7)	C9—C10—C11—C12	-0.5 (7)
C1—C2—C3—Br1	177.6 (3)	C9—C10—C11—Br2	178.0 (3)
C2—C3—C4—C5	-0.4 (8)	C10—C11—C12—C13	0.7 (7)
Br1—C3—C4—C5	-177.6 (4)	Br2—C11—C12—C13	-177.8 (4)
C3—C4—C5—C6	1.0 (8)	C11—C12—C13—C14	-1.1 (7)
C4—C5—C6—C1	-1.7 (8)	C12—C13—C14—C9	1.2 (7)
C2—C1—C6—C5	1.7 (7)	C10—C9—C14—C13	-1.0 (6)
N1—C1—C6—C5	178.7 (4)	N2—C9—C14—C13	-179.4 (4)
C1—N1—C7—O1	-3.1 (8)	C9—N2—C15—O2	2.2 (7)
C1—N1—C7—C8	177.9 (4)	C9—N2—C15—C16	-179.9 (4)

## supplementary materials

---

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N···O2 <sup>i</sup>	0.86	2.05	2.887 (5)	166
N2—H2N···O1	0.86	2.10	2.953 (5)	169

Symmetry codes: (i)  $-x+2, y-1/2, -z+1/2$ .

Fig. 1




Fig. 2

