

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Aquabis(methacrylato- κ O)bis(pyridine- κ N)copper(II)

Bin Wu* and Haizhen Yao

Department of Chemistry, Zhejiang Sci-Tech University, Hangzhou 310018, People's Republic of China

Correspondence e-mail: chemdpwu@yahoo.com.cn


Received 27 February 2009; accepted 2 April 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.021; wR factor = 0.055; data-to-parameter ratio = 14.6.

In the crystal structure of the title complex, $[\text{Cu}(\text{C}_4\text{H}_5\text{O}_2)_2(\text{C}_5\text{H}_5\text{N})_2(\text{H}_2\text{O})]$, the Cu^{II} cation is located on a twofold rotation axis and coordinated by two methacrylate anions, two pyridine ligands and one water molecule in a distorted square-pyramidal geometry. The coordinated water molecule is also located on the twofold axis. In the crystal structure $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming chains along the c axis.

Related literature

For general background to copper complexes, see: Du *et al.* (2004); Hu *et al.* (2004); Zhu *et al.* (2007). For a related structure, see: Wu & Wang (2004).



Experimental

Crystal data

 $[\text{Cu}(\text{C}_4\text{H}_5\text{O}_2)_2(\text{C}_5\text{H}_5\text{N})_2(\text{H}_2\text{O})]$ $M_r = 409.92$ Orthorhombic, $Fdd2$ $a = 15.619$ (3) Å $b = 40.200$ (8) Å $c = 6.0576$ (12) Å $V = 3803.4$ (13) Å³ $Z = 8$ Mo $K\alpha$ radiation $\mu = 1.18$ mm⁻¹ $T = 293$ K $0.50 \times 0.36 \times 0.08$ mm

Data collection

Rigaku R-Axis RAPID IP diffractometer
Absorption correction: multi-scan *ABSCOR* (Higashi, 1995)
 $T_{\text{min}} = 0.612$, $T_{\text{max}} = 0.913$ 7925 measured reflections
1808 independent reflections
1740 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.021$ $wR(F^2) = 0.055$ $S = 1.09$

1808 reflections

124 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Absolute structure: Flack (1983), 797 Friedel pairs

Flack parameter: 0.006 (13)

Table 1

Selected bond lengths (Å).

Cu—O1	2.281 (2)	Cu—N1	2.0254 (14)
Cu—O2	1.9389 (12)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots O3 ⁱ	0.83 (3)	1.96 (3)	2.783 (2)	178 (2)

Symmetry code: (i) $x, y, z + 1$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the Natural Science Foundation of Zhejiang Province, China (M203105).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2491).

References

- Du, M., Cai, H. & Zhao, X.-J. (2004). *Acta Cryst.* **E60**, m1139–m1141.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Hu, R.-Z., Liu, Z.-D., Tan, M.-Y. & Zhu, H.-L. (2004). *Acta Cryst.* **E60**, m946–m947.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Wu, B. & Wang, G. (2004). *Acta Cryst.* **E60**, m1764–m1765.
Zhu, X.-F., Zhao, L.-M., Hou, G.-F. & Gao, J.-S. (2007). *Acta Cryst.* **E63**, m1809.

supplementary materials

Acta Cryst. (2009). E65, m494 [doi:10.1107/S1600536809012422]

Aquabis(methacrylato- κO)bis(pyridine- κN)copper(II)

B. Wu and H. Yao

Comment

Copper complexes with organic acids and other donor ligands exist extensively in living things, playing an important role in a vast range of chemical and biochemical catalytic systems. A series of copper-carboxylate complexes has been reported (Du *et al.*, 2004; Hu *et al.*, 2004; Zhu *et al.*, 2007).

The molecular structure is shown in Fig. 1. The Cu atom is located on a twofold axis and coordinated with two methacrylate, two pyridine ligands and one coordinated water molecule in a distorted square-pyramidal geometry (Table 1).

The compound is an infinite one-dimensional network structure connected by hydrogen bonds. It forms hydrogen bonds between coordination waters and carboxy group (Table 2).

The corresponding complex with one pyridine ligand has binuclear cage structural unit, two Cu atoms are bridged by four μ_2 -O, $O'\alpha$ -methacrylate groups, forming a cage structure (Wu *et al.*, 2004).

Experimental

HL, $\text{CH}_2\text{C}(\text{CH}_3)\text{COOH}$, (0.5 ml, 6.0 mmol) and $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (240 mg, 1.0 mmol) were dissolved in 60 ml H_2O , and the pH adjusted to 4.0 using 0.5 M NaOH. Two milliliters of 1.0 M pyridine solution were added into the mixed solution with stirring. After filtration, the filtrate was allowed to stand at room temperature and single crystals were obtained after one week.

Refinement

Methyl H atoms were constrained to an ideal geometry with C—H distances of 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$, but each group was allowed to rotate freely about its C—C bond. The methylene H atoms and aromatic H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances of 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title molecule showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level for non-H atoms.

Aquabis(methacrylato- κ O)bis(pyridine- κ N)copper(II)

Crystal data

[Cu(C ₄ H ₅ O ₂) ₂ (C ₅ H ₅ N) ₂ (H ₂ O)]	$F_{000} = 1704$
$M_r = 409.92$	$D_x = 1.432 \text{ Mg m}^{-3}$
Orthorhombic, $Fdd2$	Mo $K\alpha$ radiation
Hall symbol: F 2 -2d	$\lambda = 0.71069 \text{ \AA}$
$a = 15.619 (3) \text{ \AA}$	Cell parameters from 8692 reflections
$b = 40.200 (8) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$c = 6.0576 (12) \text{ \AA}$	$\mu = 1.18 \text{ mm}^{-1}$
$V = 3803.4 (13) \text{ \AA}^3$	$T = 293 \text{ K}$
$Z = 8$	Platelet, blue
	$0.50 \times 0.36 \times 0.08 \text{ mm}$

Data collection

Rigaku R-Axis RAPID IP diffractometer	1808 independent reflections
Radiation source: fine-focus sealed tube	1740 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
Detector resolution: 10.00 pixels mm^{-1}	$\theta_{\text{max}} = 25.8^\circ$
$T = 293 \text{ K}$	$\theta_{\text{min}} = 3.3^\circ$
ω scans	$h = -18 \rightarrow 18$
Absorption correction: multi-scan ABSCOR (Higashi, 1995)	$k = -48 \rightarrow 48$
$T_{\text{min}} = 0.612, T_{\text{max}} = 0.913$	$l = -7 \rightarrow 7$
7925 measured reflections	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.021$	$w = 1/[\sigma^2(F_o^2) + (0.0325P)^2 + 0.6515P]$
$wR(F^2) = 0.055$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\text{max}} < 0.001$
1808 reflections	$\Delta\rho_{\text{max}} = 0.15 \text{ e \AA}^{-3}$
124 parameters	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 797 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.006 (13)

Special details

Experimental. Analysis: calculated C 52.74, H 5.41, N 6.83%; found C 52.61, H 5.22, N 6.69%. Spectroscopic analysis: IR (KBr, ν cm^{-1}): 700, 832, 936, 1036, 1214, 1243, 1368, 1417, 1599, 1642.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	1.0000	0.0000	0.11357 (7)	0.03375 (10)
O1	1.0000	0.0000	0.4901 (4)	0.0450 (5)
O2	1.05574 (8)	0.04310 (3)	0.1163 (3)	0.0442 (3)
O3	1.03022 (11)	0.05452 (4)	-0.2378 (3)	0.0529 (4)
N1	0.88531 (9)	0.02292 (3)	0.0789 (3)	0.0361 (3)
C1	1.05091 (11)	0.06268 (5)	-0.0494 (3)	0.0371 (4)
C2	1.07255 (14)	0.09857 (5)	-0.0019 (4)	0.0460 (5)
C3	1.0791 (3)	0.10857 (7)	0.2115 (5)	0.0967 (13)
H3A	1.0922	0.1306	0.2439	0.116*
H3B	1.0705	0.0934	0.3252	0.116*
C4	1.0850 (2)	0.12071 (5)	-0.1829 (5)	0.0694 (7)
H4A	1.1000	0.1423	-0.1281	0.104*
H4B	1.0331	0.1222	-0.2672	0.104*
H4C	1.1302	0.1125	-0.2752	0.104*
C5	0.83858 (12)	0.01819 (5)	-0.1017 (3)	0.0431 (5)
H5	0.8607	0.0050	-0.2143	0.052*
C6	0.75841 (12)	0.03207 (5)	-0.1284 (8)	0.0485 (4)
H6	0.7275	0.0286	-0.2576	0.058*
C7	0.72506 (13)	0.05110 (5)	0.0387 (4)	0.0493 (5)
H7	0.6705	0.0602	0.0262	0.059*
C8	0.77349 (15)	0.05649 (5)	0.2250 (5)	0.0523 (6)
H8	0.7525	0.0695	0.3397	0.063*
C9	0.85367 (14)	0.04220 (4)	0.2390 (4)	0.0443 (4)
H9	0.8868	0.0461	0.3640	0.053*
H1	1.0098 (15)	0.0163 (6)	0.569 (6)	0.052 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu	0.02640 (14)	0.04033 (15)	0.03451 (15)	0.00130 (13)	0.000	0.000

supplementary materials

O1	0.0619 (14)	0.0422 (12)	0.0309 (11)	-0.0041 (9)	0.000	0.000
O2	0.0355 (6)	0.0454 (6)	0.0516 (8)	-0.0031 (5)	-0.0055 (7)	0.0079 (7)
O3	0.0684 (10)	0.0493 (7)	0.0410 (8)	-0.0091 (7)	0.0044 (8)	-0.0081 (7)
N1	0.0290 (7)	0.0405 (7)	0.0389 (9)	0.0003 (6)	0.0015 (7)	0.0002 (7)
C1	0.0287 (9)	0.0394 (9)	0.0432 (10)	0.0040 (7)	0.0049 (8)	-0.0027 (8)
C2	0.0534 (12)	0.0379 (9)	0.0468 (11)	0.0101 (8)	-0.0090 (10)	-0.0030 (9)
C3	0.185 (4)	0.0516 (14)	0.0534 (16)	0.0172 (18)	-0.027 (2)	-0.0070 (12)
C4	0.109 (2)	0.0436 (10)	0.0552 (16)	-0.0013 (12)	-0.0059 (15)	0.0058 (11)
C5	0.0357 (9)	0.0541 (10)	0.0394 (12)	0.0002 (8)	-0.0002 (8)	-0.0046 (8)
C6	0.0362 (9)	0.0556 (10)	0.0538 (11)	-0.0010 (8)	-0.0128 (11)	0.0031 (17)
C7	0.0319 (9)	0.0450 (10)	0.0711 (15)	0.0064 (8)	-0.0002 (10)	0.0046 (10)
C8	0.0475 (12)	0.0461 (11)	0.0634 (14)	0.0101 (9)	0.0061 (11)	-0.0088 (11)
C9	0.0419 (11)	0.0457 (9)	0.0453 (11)	0.0022 (8)	-0.0027 (9)	-0.0038 (9)

Geometric parameters (\AA , $^\circ$)

Cu—O1	2.281 (2)	C3—H3B	0.9300
Cu—O2	1.9389 (12)	C4—H4A	0.9600
Cu—O2 ⁱ	1.9391 (12)	C4—H4B	0.9600
Cu—N1	2.0254 (14)	C4—H4C	0.9600
Cu—N1 ⁱ	2.0254 (14)	C5—C6	1.380 (3)
O1—H1	0.82 (3)	C5—H5	0.9300
O2—C1	1.278 (2)	C6—C7	1.371 (5)
O3—C1	1.230 (3)	C6—H6	0.9300
N1—C5	1.329 (3)	C7—C8	1.376 (4)
N1—C9	1.336 (3)	C7—H7	0.9300
C1—C2	1.510 (3)	C8—C9	1.380 (3)
C2—C3	1.357 (4)	C8—H8	0.9300
C2—C4	1.425 (3)	C9—H9	0.9300
C3—H3A	0.9300		
O2—Cu—O2 ⁱ	179.03 (10)	H3A—C3—H3B	120.0
O2—Cu—N1	89.51 (5)	C2—C4—H4A	109.5
O2 ⁱ —Cu—N1	90.59 (5)	C2—C4—H4B	109.5
O2—Cu—N1 ⁱ	90.59 (5)	H4A—C4—H4B	109.5
O2 ⁱ —Cu—N1 ⁱ	89.51 (5)	C2—C4—H4C	109.5
N1—Cu—N1 ⁱ	168.08 (10)	H4A—C4—H4C	109.5
O2—Cu—O1	89.51 (5)	H4B—C4—H4C	109.5
O2 ⁱ —Cu—O1	89.51 (5)	N1—C5—C6	122.5 (3)
N1—Cu—O1	95.96 (5)	N1—C5—H5	118.8
N1 ⁱ —Cu—O1	95.96 (5)	C6—C5—H5	118.8
Cu—O1—H1	126 (2)	C7—C6—C5	118.9 (3)
C1—O2—Cu	121.15 (13)	C7—C6—H6	120.5
C5—N1—C9	118.52 (16)	C5—C6—H6	120.5
C5—N1—Cu	120.41 (13)	C6—C7—C8	119.0 (2)
C9—N1—Cu	121.03 (14)	C6—C7—H7	120.5
O3—C1—O2	125.43 (17)	C8—C7—H7	120.5
O3—C1—C2	119.36 (18)	C7—C8—C9	118.9 (2)

O2—C1—C2	115.21 (18)	C7—C8—H8	120.5
C3—C2—C4	122.5 (2)	C9—C8—H8	120.5
C3—C2—C1	118.8 (2)	N1—C9—C8	122.1 (2)
C4—C2—C1	118.7 (2)	N1—C9—H9	118.9
C2—C3—H3A	120.0	C8—C9—H9	118.9
C2—C3—H3B	120.0		

Symmetry codes: (i) $-x+2, -y, z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O3 ⁱⁱ	0.83 (3)	1.96 (3)	2.783 (2)	178 (2)

Symmetry codes: (ii) $x, y, z+1$.

Fig. 1

