

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dimethyl [(4-fluorophenyl)(6-methoxybenzothiazol-2-ylamino)methyl]-phosphonate

 Yan-Ping Hong,^{a,b*} Bao-An Song^b and Xin-Chen Shangguan^a
^aCollege of Food Science and Engineering, Jiangxi Agricultural University, Nanchang 330045, People's Republic of China, and ^bKey Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, Guizhou University, Guiyang 550025, People's Republic of China

Correspondence e-mail: jxndspxy74@yahoo.cn


Received 19 April 2009; accepted 24 April 2009

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.041; wR factor = 0.115; data-to-parameter ratio = 14.4.

In the molecule of title compound, $\text{C}_{17}\text{H}_{18}\text{FN}_2\text{O}_4\text{PS}$, both the benzene ring with its conjunction C atom and the benzothiazole ring with its conjunction N atom are close to planar (the maximum deviations are 0.0267 and 0.0427 Å for the benzene and benzothiazole rings, respectively), the dihedral angle between the planes of the benzothiazole and benzene rings is 119.05 (3)°. The molecular packing is stabilized by intermolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{F}$ hydrogen bonding, and by $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ stacking interactions [centroid-centroid distances = 2.99 (2), 2.96 (3), 2.88 (2) and 3.773 (4) Å].

Related literature

For the biological activity of α -aminophosphonate derivatives, see: Kafarski & Lejczak (2001); De Lombaert *et al.* (1995); Du *et al.*, (1999). For activities of α -aminophosphonate derivatives containing an F atom and benzothiazole or isoxazole units, see: Yang *et al.* (2005); Song *et al.* (2005); Jin *et al.* (2006). For related structures, see: Fang *et al.* (2009); Yang *et al.* (2005); Jin *et al.* (2006); Song *et al.* (2005); Alvarez *et al.* (2005); Chen & Li (1987); Li *et al.* (2008).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{FN}_2\text{O}_4\text{PS}$
 $M_r = 396.36$
 Monoclinic, $C2/c$
 $a = 19.761$ (5) Å
 $b = 15.781$ (4) Å
 $c = 14.395$ (4) Å
 $\beta = 122.109$ (3)°
 $V = 3802.4$ (16) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.29$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.26 \times 0.22$ mm

Data collection

Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.919$, $T_{\max} = 0.939$
 9669 measured reflections
 3428 independent reflections
 2477 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.115$
 $S = 1.04$
 3428 reflections
 238 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.86	1.99	2.793 (3)	156
$\text{C16}-\text{H16C}\cdots\text{F1}^{\text{ii}}$	0.96	2.51	3.2496 (133)	133
$\text{C15}-\text{H15B}\cdots\text{Cg1}^{\text{iii}}$	0.96	2.99	3.608 (4)	123
$\text{C16}-\text{H16B}\cdots\text{Cg3}^{\text{iv}}$	0.96	2.96	3.549 (4)	121
$\text{C17}-\text{H17C}\cdots\text{Cg1}^{\text{v}}$	0.96	2.88	3.625 (3)	136

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z$; (ii) $-x + 1, y, -z + \frac{1}{2}$; (iii) $-x, y, -z - \frac{1}{2}$; (iv) $-x + \frac{1}{2}, y + \frac{1}{2}, -z - \frac{1}{2}$; (v) $-x, y, -z + \frac{1}{2}$. Cg1 and Cg3 are the centroids of the S1/N2/C4-C6 and C9-C14 rings, respectively.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors acknowledge Jiangxi Agricultural University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2769).

References

- Alvarez, R. G., Kennedy, A. R., Khalaf, A. I., Suckling, C. J. & Waigh, R. D. (2005). *Acta Cryst.* **E61**, o569–o570.
 Bruker (2004). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chen, R. Y. & Li, Y. G. (1987). *Chemistry of Organic Phosphorus*, p.38. Beijing: High Education Press.
 De Lombaert, S., Blanchard, L., Tan, J., Sakane, Y., Berry, C. & Ghai, R. D. (1995). *Bioorg. Med. Chem. Lett.* **5**, 145–150.
 Du, S. C., Faiger, H., Belakhov, V. & Timor Bassov, T. (1999). *Bioorg. Med. Chem.* **7**, 2671–2682.
 Fang, H., Fang, M.-J., Xu, Y., Yu, W.-C. & Zhao, Y.-F. (2009). *Acta Cryst.* **E65**, o642.
 Jin, L. H., Song, B. A., Zhang, G. P., Xu, R. Q., Zhang, S. M., Gao, X. W., Hu, D. Y. & Yang, S. (2006). *Bioorg. Med. Chem. Lett.* **16**, 1537–1543.
 Kafarski, P. & Lejczak, B. (2001). *Curr. Med. Chem. Anti-Cancer Agents*, **1**, 301–312.

Li, B., Zhang, S., Wang, Y. & Luo, S. (2008). *Acta Cryst.* **E64**, o1549.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Song, B. A., Chen, C. J., Yang, S., Jin, L. H., Xue, W., Zhang, S. M., Zou, Z. H.,
Hu, D. Y. & Liu, G. (2005). *Acta Chim. Sin.* **63**, 1720–1726.
Yang, S., Song, B. A., Hong, Y. P., Jin, L. H. & Hu, D. Y. (2005). *J. Chem.*
Crystallogr. **11**, 891–895.

supporting information

Acta Cryst. (2009). E65, o1199–o1200 [doi:10.1107/S1600536809015384]

Dimethyl [(4-fluorophenyl)(6-methoxybenzothiazol-2-ylamino)methyl]phosphonate

Yan-Ping Hong, Bao-An Song and Xin-Chen Shangguan

S1. Comment

α -Aminophosphonate derivatives, as isosteres of natural aminocarboxylic acids, have attracted much attention in medicinal and pesticide chemistry in recent years due to their wide range of biological activities (Kafarski *et al.*, 2001; De Lombaert *et al.*, 1995; Du *et al.*, 1999). In our previous work, a plenty of α -aminophosphonate derivatives containing fluorine atom and benzothiazole or isoxazole moieties had been synthesized and some of them showed fungicidal (Yang *et al.*, 2005) and antitumor activities (Song *et al.*, 2005; Jin *et al.*, 2006). And some crystals of α -aminophosphonate derivatives containing dialkyl had been reported in our previous works. However, It is noteworthy that a crystal of α -aminophosphonate containing dimethyl moiety was obtained for the first time. We report herein the crystal structure of the title compound.

As illustrated in Fig. 1, both the benzene ring with its conjunction carbon atom C8 and the benzothiazole ring with its conjunction nitrogen atom N1 are fairly planar, the dihedral angle between the planes of the benzothiazole group (C1–C7/N2/S1) and the benzene ring (C9–C14) is 119.05°. The P atom exhibits a distorted tetrahedral configuration because the bond angles of O2–P1–O3 = 116.07 (11)° and O2–P1–O4 = 114.04 (10)° are significantly larger than that of O3–P1–O4 = 103.65 (10)°. The P1–C8 = 1.807 (2) Å is similar to the corresponding P–C value of 1.809 (3) Å found in diisopropyl [(benzoylamino)(phenyl)methyl] phosphonate (Fang *et al.*, 2009) and a little shorter than normal P–C single bond length of 1.850 Å (Chen *et al.*, 1987). The C5–N1 = 1.355 (3) Å is remarkably shorter than normal C–N 1.471 (3) Å (Alvarez *et al.*, 2005) and close to the C = N = 1.343 (2) Å, similar to the corresponding bond length of 1.358 (3) Å of the 2-(1-piperidinyl)-1,3-benzothiazole (Alvarez *et al.*, 2005). Meanwhile, the S1–C6 and S1–C5 with bond lengths of 1.742 (2) Å and 1.760 (2) Å, respectively, are shorter than the typical C–S of 1.811 (9) Å (Li *et al.*, 2008). These indicate that the N1 atom and benzothiazole ring form a considerable delocalization of the electron density in which the N1 atom is sp^2 hybridized. The molecular packing is consolidated through weak inter/intra molecular N–H \cdots O, C–H \cdots N, and C–H \cdots F hydrogen bonding, C–H \cdots π and π – π stacking interactions (Table 2, Fig. 2, Cg1 = ring (S1/N2/C4–C6); Cg3 = ring(C9–C14)). C–H \cdots π interactions of methyl H atoms are established towards the π -systems of neighboring aromatic groups from 4-fluorophenyl and five rings of 6-methoxybenzothiazol-2-ylamino, with centroid-to-centroid distance are 2.99 (2), 2.96 (3) and 2.88 (2) Å, respectively. In the π – π stacking interactions between two close-by phenyl rings of 4-fluorophenyl, the centroid-to-centroid distance is 3.773 (4) Å.

S2. Experimental

A mixture of 2-amino-6-methoxybenzothiazole (0.9010 g, 5 mmol) and 4-fluorobenzaldehyde (0.6205 g, 5 mmol) and 20 ml toluene in a 50 ml dry flask affiliated a water separator was refluxed and stirred for 6 h. Toluene was removed by water separator and then dimethylphosphate (0.88 g, 8 mmol) was added into the flask, the reaction mixture was refluxed in a nonsolvent condition for 5 h. Residue was washed with water, filtered, dried, and crystallized three times from

ethanol yielding title compound as colorless solid. The crystal suitable for X-ray analysis was obtained by slow evaporation of an anhydrous ethanol at room temperature over a period of ten days.

S3. Refinement

H atoms were placed in calculated positions and were treated as riding on the parent C atoms with $C-H = 0.93 - 0.987$ Å; $N-H = 0.86$ Å, and with $U_{iso}(H) = 1.2U_{eq}(C)$ for CH_2 , CH , NH , and $1.5U_{eq}(C)$ for CH_3 .




Figure 1

The structure of the title compound, showing as 50% probability displacement ellipsoids.




Figure 2

A packing view of the title compound. $N-H\cdots O$, $C-H\cdots O$, $C-H\cdots N$ and $C-H\cdots F$ hydrogen bonds, $C-H\cdots\pi$ and $\pi-\pi$ stacking interactions are shown as dashed lines.

Dimethyl [(4-fluorophenyl)(6-methoxybenzothiazol-2-ylamino)methyl]phosphonate

Crystal data

$C_{17}H_{18}FN_2O_4PS$

$M_r = 396.36$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 19.761$ (5) Å

$b = 15.781$ (4) Å

$c = 14.395$ (4) Å

$\beta = 122.109$ (3)°

$V = 3802.4$ (16) Å³

$Z = 8$

$F(000) = 1648$
 $D_x = 1.385 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 5837 reflections
 $\theta = 2.8\text{--}27.9^\circ$

$\mu = 0.29 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Block, colourless
 $0.30 \times 0.26 \times 0.22 \text{ mm}$

Data collection

Bruker APEXII area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.919$, $T_{\max} = 0.939$

9669 measured reflections
 3428 independent reflections
 2477 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -23 \rightarrow 23$
 $k = -18 \rightarrow 18$
 $l = -15 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.115$
 $S = 1.04$
 3428 reflections
 238 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0545P)^2 + 1.1262P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.07360 (4)	0.28313 (4)	0.06124 (5)	0.0510 (2)
P1	0.20273 (4)	0.09893 (4)	-0.09266 (5)	0.0483 (2)
N2	0.06939 (11)	0.11776 (12)	0.06276 (16)	0.0477 (5)
C5	0.10708 (13)	0.18124 (14)	0.05470 (17)	0.0421 (5)
C4	0.00865 (13)	0.14714 (14)	0.07713 (18)	0.0445 (5)
C6	0.00177 (13)	0.23554 (14)	0.07921 (17)	0.0438 (5)
C7	-0.05321 (14)	0.27298 (16)	0.09682 (19)	0.0524 (6)
H7	-0.0568	0.3317	0.0981	0.063*
C1	-0.10315 (14)	0.22182 (16)	0.11256 (19)	0.0506 (6)
C3	-0.04236 (14)	0.09767 (16)	0.0913 (2)	0.0571 (7)
H3	-0.0395	0.0390	0.0890	0.069*
C2	-0.09822 (15)	0.13484 (16)	0.1089 (2)	0.0563 (6)

H2	-0.1326	0.1009	0.1184	0.068*
N1	0.16956 (11)	0.17525 (12)	0.04083 (17)	0.0524 (5)
H1	0.1923	0.2209	0.0382	0.063*
C8	0.19929 (13)	0.09419 (14)	0.03023 (19)	0.0452 (6)
H8	0.1605	0.0505	0.0196	0.054*
C9	0.27995 (14)	0.06879 (14)	0.12740 (18)	0.0452 (6)
C11	0.37489 (18)	-0.04111 (18)	0.2316 (2)	0.0659 (8)
H11	0.3884	-0.0982	0.2439	0.079*
C10	0.30051 (17)	-0.01584 (16)	0.1484 (2)	0.0560 (6)
H10	0.2630	-0.0567	0.1049	0.067*
C14	0.33614 (15)	0.12855 (17)	0.1936 (2)	0.0625 (7)
H14	0.3237	0.1859	0.1810	0.075*
C12	0.42812 (17)	0.0196 (2)	0.2955 (2)	0.0684 (8)
C13	0.41102 (17)	0.1034 (2)	0.2788 (3)	0.0751 (9)
H13	0.4489	0.1435	0.3238	0.090*
F1	0.50137 (11)	-0.00510 (13)	0.37934 (15)	0.1064 (7)
O1	-0.15552 (10)	0.26361 (12)	0.13111 (15)	0.0658 (5)
C17	-0.20925 (16)	0.21423 (19)	0.1466 (2)	0.0707 (8)
H17A	-0.2437	0.1822	0.0813	0.106*
H17B	-0.2409	0.2511	0.1617	0.106*
H17C	-0.1794	0.1762	0.2071	0.106*
O2	0.24876 (11)	0.17048 (11)	-0.09456 (13)	0.0612 (5)
O4	0.23331 (10)	0.01050 (10)	-0.10441 (13)	0.0555 (4)
O3	0.11234 (11)	0.09795 (13)	-0.18384 (15)	0.0752 (6)
C16	0.31296 (19)	-0.0033 (2)	-0.0807 (3)	0.0886 (10)
H16A	0.3292	0.0444	-0.1056	0.133*
H16B	0.3138	-0.0536	-0.1177	0.133*
H16C	0.3490	-0.0101	-0.0031	0.133*
C15	0.0861 (2)	0.1088 (3)	-0.2974 (3)	0.1293 (16)
H15A	0.1166	0.1532	-0.3040	0.194*
H15B	0.0305	0.1236	-0.3386	0.194*
H15C	0.0938	0.0569	-0.3254	0.194*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0591 (4)	0.0372 (4)	0.0646 (4)	-0.0040 (3)	0.0382 (3)	0.0005 (3)
P1	0.0512 (4)	0.0440 (4)	0.0452 (3)	-0.0112 (3)	0.0226 (3)	-0.0035 (3)
N2	0.0461 (11)	0.0395 (11)	0.0582 (12)	-0.0059 (9)	0.0282 (10)	-0.0038 (9)
C5	0.0426 (13)	0.0393 (13)	0.0389 (12)	-0.0061 (10)	0.0180 (11)	-0.0032 (9)
C4	0.0451 (13)	0.0396 (14)	0.0455 (13)	-0.0073 (10)	0.0219 (11)	-0.0049 (10)
C6	0.0455 (13)	0.0416 (14)	0.0425 (12)	-0.0052 (11)	0.0221 (11)	-0.0035 (10)
C7	0.0572 (15)	0.0428 (14)	0.0582 (15)	-0.0044 (12)	0.0313 (13)	-0.0054 (11)
C1	0.0490 (14)	0.0532 (16)	0.0514 (14)	-0.0053 (12)	0.0278 (12)	-0.0090 (11)
C3	0.0579 (16)	0.0412 (15)	0.0777 (18)	-0.0081 (12)	0.0398 (15)	-0.0053 (12)
C2	0.0549 (15)	0.0535 (17)	0.0678 (16)	-0.0135 (12)	0.0374 (14)	-0.0064 (12)
N1	0.0529 (12)	0.0390 (11)	0.0741 (14)	-0.0097 (9)	0.0396 (11)	-0.0039 (10)
C8	0.0488 (14)	0.0367 (13)	0.0547 (14)	-0.0089 (11)	0.0306 (12)	-0.0056 (10)

C9	0.0538 (14)	0.0431 (14)	0.0464 (13)	-0.0045 (11)	0.0319 (12)	-0.0008 (10)
C11	0.087 (2)	0.0563 (18)	0.0570 (16)	0.0167 (16)	0.0403 (17)	0.0113 (14)
C10	0.0756 (18)	0.0459 (16)	0.0494 (14)	-0.0029 (13)	0.0352 (14)	-0.0002 (11)
C14	0.0611 (17)	0.0487 (16)	0.0647 (17)	-0.0037 (13)	0.0247 (15)	-0.0009 (13)
C12	0.0643 (18)	0.079 (2)	0.0573 (16)	0.0179 (16)	0.0290 (15)	0.0111 (15)
C13	0.0598 (18)	0.070 (2)	0.0721 (19)	-0.0062 (15)	0.0192 (16)	-0.0051 (15)
F1	0.0756 (12)	0.1209 (17)	0.0865 (13)	0.0313 (11)	0.0186 (10)	0.0189 (11)
O1	0.0625 (11)	0.0652 (12)	0.0842 (13)	-0.0062 (9)	0.0488 (11)	-0.0148 (10)
C17	0.0617 (18)	0.087 (2)	0.0781 (19)	-0.0073 (15)	0.0467 (16)	-0.0076 (15)
O2	0.0787 (12)	0.0477 (10)	0.0626 (11)	-0.0209 (9)	0.0412 (10)	-0.0049 (8)
O4	0.0680 (11)	0.0442 (10)	0.0622 (10)	-0.0108 (8)	0.0399 (9)	-0.0105 (8)
O3	0.0579 (11)	0.0853 (14)	0.0560 (11)	-0.0091 (10)	0.0125 (10)	0.0023 (10)
C16	0.093 (2)	0.086 (2)	0.125 (3)	0.0037 (19)	0.084 (2)	0.003 (2)
C15	0.120 (3)	0.149 (4)	0.054 (2)	-0.030 (3)	0.003 (2)	0.009 (2)

Geometric parameters (Å, °)

S1—C6	1.742 (2)	C9—C14	1.381 (3)
S1—C5	1.760 (2)	C9—C10	1.382 (3)
P1—O2	1.4593 (17)	C11—C12	1.357 (4)
P1—O3	1.5560 (19)	C11—C10	1.372 (4)
P1—O4	1.5653 (18)	C11—H11	0.9300
P1—C8	1.807 (2)	C10—H10	0.9300
N2—C5	1.290 (3)	C14—C13	1.386 (4)
N2—C4	1.401 (3)	C14—H14	0.9300
C5—N1	1.355 (3)	C12—C13	1.355 (4)
C4—C3	1.373 (3)	C12—F1	1.359 (3)
C4—C6	1.404 (3)	C13—H13	0.9300
C6—C7	1.374 (3)	O1—C17	1.427 (3)
C7—C1	1.384 (3)	C17—H17A	0.9600
C7—H7	0.9300	C17—H17B	0.9600
C1—O1	1.368 (3)	C17—H17C	0.9600
C1—C2	1.379 (3)	O4—C16	1.439 (3)
C3—C2	1.388 (3)	O3—C15	1.438 (4)
C3—H3	0.9300	C16—H16A	0.9600
C2—H2	0.9300	C16—H16B	0.9600
N1—C8	1.449 (3)	C16—H16C	0.9600
N1—H1	0.8600	C15—H15A	0.9600
C8—C9	1.513 (3)	C15—H15B	0.9600
C8—H8	0.9800	C15—H15C	0.9600
C6—S1—C5	88.47 (10)	C14—C9—C8	121.5 (2)
O2—P1—O3	116.07 (11)	C10—C9—C8	120.1 (2)
O2—P1—O4	114.04 (10)	C12—C11—C10	118.1 (3)
O3—P1—O4	103.65 (10)	C12—C11—H11	120.9
O2—P1—C8	113.48 (10)	C10—C11—H11	120.9
O3—P1—C8	101.64 (11)	C11—C10—C9	121.7 (3)
O4—P1—C8	106.64 (10)	C11—C10—H10	119.1

C5—N2—C4	109.72 (19)	C9—C10—H10	119.1
N2—C5—N1	125.0 (2)	C9—C14—C13	120.3 (3)
N2—C5—S1	116.95 (17)	C9—C14—H14	119.9
N1—C5—S1	118.01 (17)	C13—C14—H14	119.9
C3—C4—N2	126.0 (2)	C13—C12—C11	122.6 (3)
C3—C4—C6	118.4 (2)	C13—C12—F1	119.0 (3)
N2—C4—C6	115.62 (19)	C11—C12—F1	118.4 (3)
C7—C6—C4	121.8 (2)	C12—C13—C14	119.0 (3)
C7—C6—S1	128.96 (18)	C12—C13—H13	120.5
C4—C6—S1	109.24 (16)	C14—C13—H13	120.5
C6—C7—C1	118.8 (2)	C1—O1—C17	118.1 (2)
C6—C7—H7	120.6	O1—C17—H17A	109.5
C1—C7—H7	120.6	O1—C17—H17B	109.5
O1—C1—C2	124.3 (2)	H17A—C17—H17B	109.5
O1—C1—C7	115.5 (2)	O1—C17—H17C	109.5
C2—C1—C7	120.2 (2)	H17A—C17—H17C	109.5
C4—C3—C2	120.3 (2)	H17B—C17—H17C	109.5
C4—C3—H3	119.8	C16—O4—P1	123.11 (18)
C2—C3—H3	119.8	C15—O3—P1	121.1 (2)
C1—C2—C3	120.5 (2)	O4—C16—H16A	109.5
C1—C2—H2	119.8	O4—C16—H16B	109.5
C3—C2—H2	119.8	H16A—C16—H16B	109.5
C5—N1—C8	121.96 (19)	O4—C16—H16C	109.5
C5—N1—H1	119.0	H16A—C16—H16C	109.5
C8—N1—H1	119.0	H16B—C16—H16C	109.5
N1—C8—C9	115.05 (18)	O3—C15—H15A	109.5
N1—C8—P1	107.14 (15)	O3—C15—H15B	109.5
C9—C8—P1	110.28 (15)	H15A—C15—H15B	109.5
N1—C8—H8	108.0	O3—C15—H15C	109.5
C9—C8—H8	108.0	H15A—C15—H15C	109.5
P1—C8—H8	108.0	H15B—C15—H15C	109.5
C14—C9—C10	118.3 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O2 ⁱ	0.86	1.99	2.793 (3)	156
C8—H8...N2	0.98	2.44	2.868 (4)	106
C14—H14...N1	0.93	2.62	2.919 (5)	100
C16—H16C...F1 ⁱⁱ	0.96	2.51	3.2496	133
C15—H15B...Cg1 ⁱⁱⁱ	0.96	2.99	3.608 (4)	123
C16—H16B...Cg3 ^{iv}	0.96	2.96	3.549 (4)	121
C17—H17C...Cg1 ^v	0.96	2.88	3.625 (3)	136

Symmetry codes: (i) $-x+1/2, -y+1/2, -z$; (ii) $-x+1, y, -z+1/2$; (iii) $-x, y, -z-1/2$; (iv) $-x+1/2, y+1/2, -z-1/2$; (v) $-x, y, -z+1/2$.