

(1-Adamantyl){4-[(2-chloro-9-isopropyl-9H-purin-6-yl)aminomethyl]phenyl}-methanone trichloromethane solvate

Michal Rouchal,^a Marek Nečas^b and Robert Vícha^{a*}

^aDepartment of Chemistry, Faculty of Technology, Tomas Bata University in Zlín, Nám. T. G. Masaryka 275, Zlín 762 72, Czech Republic, and ^bDepartment of Chemistry, Faculty of Science, Masaryk University in Brno, Kamenice 5, Brno-Bohunice 625 00, Czech Republic
Correspondence e-mail: rvicha@ft.utb.cz


Received 27 March 2009; accepted 4 May 2009

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.050; wR factor = 0.177; data-to-parameter ratio = 14.9.

In the title compound, $\text{C}_{26}\text{H}_{30}\text{ClN}_5\text{O}\cdot\text{CHCl}_3$, the purine molecule consists of essentially planar benzene and purine ring systems [maximum deviation 0.010 (4) Å for both ring systems] forming a dihedral angle of 85.52 (9)°. Intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds link adjacent molecules into centrosymmetric dimers. The structure also contains intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ interactions. The benzene rings form offset face-to-face $\pi-\pi$ stacking interactions with an interplanar distance of 3.541 (4) Å and a centroid-to-centroid distance of 4.022 (4) Å.

Related literature

The title compound was prepared according to a modified literature procedure (Fiorini & Abel, 1998). For the synthesis and/or biological activity of related compounds, see: Legraverend & Grierson (2006); Long *et al.* (2007). For related structures, see: Trávníček & Kryštof (2004); Trávníček & Zatloukal (2004); Trávníček & Popa (2007a,b) Rouchal *et al.* (2009).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{30}\text{ClN}_5\text{O}\cdot\text{CHCl}_3$
 $M_r = 583.39$
Orthorhombic, $Pbca$
 $a = 19.434$ (12) Å
 $b = 13.186$ (7) Å
 $c = 22.149$ (11) Å
 $V = 5676$ (5) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.45$ mm⁻¹
 $T = 120$ K
0.45 × 0.40 × 0.20 mm

Data collection

Kuma KM4 CCD diffractometer
Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2006)
 $T_{\min} = 0.738$, $T_{\max} = 0.917$
32219 measured reflections
4984 independent reflections
2981 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.177$
 $S = 1.12$
4984 reflections
334 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.71$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.54$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{N5}^i$	0.88	2.22	3.013 (5)	150
$\text{C27}-\text{H27A}\cdots\text{N2}^{ii}$	1.00	2.59	3.553 (6)	161
$\text{C5}-\text{H5B}\cdots\text{N3}^{iii}$	1.00	2.66	3.641 (5)	166
$\text{C23}-\text{H23A}\cdots\text{O1}^{iv}$	0.95	2.23	3.179 (5)	175

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $-x + \frac{1}{2}, -y + 2, z + \frac{1}{2}$; (iii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (iv) $x, y - 1, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Financial support of this work by the Science Foundation of the Czech Republic (grant No. 203/06/P362) and the Czech Ministry of Education (project No. MSM 7088352101) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BI2361).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Fiorini, M. T. & Abel, Ch. (1998). *Tetrahedron Lett.* **39**, 1827–1830.
Legraverend, M. & Grierson, D. S. (2006). *Bioorg. Med. Chem.* **14**, 3987–4006.
Long, J., Manchandia, T., Ban, K., Gao, S., Miller, C. & Chandra, J. (2007). *Cancer Chemother. Pharmacol.* **59**, 527–535.
Oxford Diffraction (2006). *CrysAlis CCD and CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
Rouchal, M., Nečas, M., de Carvalho, F. P. & Vícha, R. (2009). *Acta Cryst.* **E65**, o298–o299.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Trávníček, Z. & Kryštof, V. (2004). *Acta Cryst.* **E60**, o2324–o2327.
Trávníček, Z. & Popa, I. (2007a). *Acta Cryst.* **E63**, o629–o631.
Trávníček, Z. & Popa, I. (2007b). *Acta Cryst.* **E63**, o728–o730.
Trávníček, Z. & Zatloukal, M. (2004). *Acta Cryst.* **E60**, o924–o926.

supplementary materials

Acta Cryst. (2009). E65, o1268 [doi:10.1107/S1600536809016596]

(1-Adamantyl){4-[(2-chloro-9-isopropyl-9*H*-purin-6-yl)aminomethyl]phenyl}methanone trichloromethane solvate

M. Rouchal, M. Necas and R. Vícha

Comment

Purine molecules substituted at C2, C6 and N9 are known as potential ATP-competitive inhibitors of cyclin-dependent kinases (key regulators of the cell division cycle) and due to this fact, they can show an anticancer activity (Legraverend & Grierson, 2006). The adamantane scaffold is often used in order to improve the biological properties of potential or known drugs and a number of compounds with various biological activity have been described, *e.g.* antitumor agents published by Long *et al.* (2007).

The asymmetric unit of the title compound (Fig. 1) consists of a trisubstituted purine molecule with trichloromethane solvent in the ratio of one to one. Both benzene and purine rings are essentially planar with maximum deviation from the best plane being 0.010 (4) Å for C19 of the purine ring and 0.010 (4) Å for C16 of the benzene ring. The dihedral angle between purine and benzene rings is 85.52 (9)°. The torsion angles C18/N1/C19/C22, C19/N1/C18/C15, N1/C18/C15/C16 and H24A/C24/N4/C21 are 173.42 (4), -79.52 (4), -36.80 (5) and -34.28 (5)° respectively. Adjacent molecules are linked into centrosymmetric pairs by intermolecular N1—H1A···N5ⁱ hydrogen bonds (Table 2, Fig. 2; symmetry code: (i) -x, 1 - y, 1 - z). The trichloromethane molecule forms intermolecular C—H···N and C—H···Cl interactions. Additional intermolecular interactions (Table 2) include C23—H23A···O1, C5—H5B···N3, C—H···Cl contacts and offset face-to-face π - π interactions with an interplanar distance of 3.541 (4) Å and a centroid-centroid distance of 4.022 (4) Å.

Experimental

The title compound was prepared according to a slightly modified literature procedure (Fiorini & Abel, 1998). 2,6-Dichloro-9-(propan-2-yl)-9*H*-purine (0.53 mmol, 122.5 mg) and (1-adamantyl)-[4-(aminomethyl)phenyl]methanone hydrochloride (0.56 mmol, 170 mg) were dissolved in a mixture of DMF (2 ml) and triethylamine (1.06 mmol, 0.15 ml). The resulting solution was stirred under reflux for 2 h (the reaction progress was monitored by TLC). After this period, the mixture was diluted with water and extracted five times with 15 ml of diethyl ether. The combined organic layers were washed twice with brine and dried over sodium sulfate. The desired product was obtained by evaporation of the solvent in vacuum and purified by column chromatography (silica gel; petroleum ether/ethyl acetate, *v/v* 1:1) to give a colourless crystalline powder (210 mg, 86%, mp 190–192°C). The crystal used for data collection was acquired by evaporation from a chloroform solution at room temperature.

Figures




Fig. 1. The molecular structure with displacement ellipsoids drawn at 50% probability for non-H atoms.




Fig. 2. Centrosymmetric dimer linked by N—H...N hydrogen bonds (dashed lines). Displacement ellipsoids are drawn at 50% probability for non-H atoms.

(1-Adamantyl){4-[(2-chloro-9-isopropyl-9H-purin-6-yl)aminomethyl]phenyl}methanone trichloromethane solvate

Crystal data

$C_{26}H_{30}ClN_5O \cdot CHCl_3$

$M_r = 583.39$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 19.434$ (12) Å

$b = 13.186$ (7) Å

$c = 22.149$ (11) Å

$V = 5676$ (5) Å³

$Z = 8$

$F_{000} = 2432$

$D_x = 1.365$ Mg m⁻³

Melting point: 191 K

Mo *K*α radiation

$\lambda = 0.71073$ Å

Cell parameters from 4984 reflections

$\theta = 2.6$ – 25.0°

$\mu = 0.45$ mm⁻¹

$T = 120$ K

Block, colourless

$0.45 \times 0.40 \times 0.20$ mm

Data collection

Kuma KM4 CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 0.06 pixels mm⁻¹

$T = 120$ K

ω scans

Absorption correction: multi-scan
(CrysAlisRED; Oxford Diffraction, 2006)

$T_{\min} = 0.738$, $T_{\max} = 0.917$

32219 measured reflections

4984 independent reflections

2981 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\max} = 25.0^\circ$

$\theta_{\min} = 2.6^\circ$

$h = -23 \rightarrow 21$

$k = -15 \rightarrow 15$

$l = -26 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.177$

$S = 1.12$

4984 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0729P)^2 + 9.6676P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.71$ e Å⁻³

334 parameters

$$\Delta\rho_{\min} = -0.54 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.15090 (5)	0.68048 (7)	0.25476 (4)	0.0305 (3)
O1	0.13811 (13)	1.1399 (2)	0.50691 (13)	0.0320 (7)
N1	0.00385 (14)	0.6307 (2)	0.43338 (14)	0.0230 (7)
H1A	-0.0135	0.5931	0.4625	0.028*
N2	0.07497 (15)	0.6451 (2)	0.34920 (14)	0.0226 (7)
N3	0.15847 (14)	0.5170 (2)	0.32042 (13)	0.0217 (7)
N4	0.15553 (15)	0.3704 (2)	0.38744 (13)	0.0233 (7)
N5	0.07206 (15)	0.4204 (2)	0.45195 (14)	0.0256 (7)
C1	0.21373 (18)	1.0428 (3)	0.56796 (17)	0.0254 (9)
C2	0.2555 (2)	1.1416 (3)	0.56949 (19)	0.0307 (9)
H2B	0.2251	1.1982	0.5819	0.037*
H2C	0.2729	1.1566	0.5284	0.037*
C3	0.3165 (2)	1.1347 (3)	0.61349 (18)	0.0322 (10)
H3B	0.3428	1.2000	0.6132	0.039*
C4	0.2899 (2)	1.1132 (3)	0.67731 (19)	0.0355 (10)
H4A	0.2600	1.1696	0.6908	0.043*
H4B	0.3292	1.1080	0.7056	0.043*
C5	0.2487 (2)	1.0128 (3)	0.67746 (18)	0.0322 (10)
H5B	0.2315	0.9988	0.7192	0.039*
C6	0.2956 (2)	0.9260 (3)	0.65710 (19)	0.0361 (10)
H6A	0.2695	0.8615	0.6572	0.043*
H6B	0.3346	0.9192	0.6856	0.043*
C7	0.3232 (2)	0.9470 (3)	0.59318 (18)	0.0299 (9)
H7A	0.3539	0.8901	0.5805	0.036*
C8	0.26304 (19)	0.9563 (3)	0.54841 (17)	0.0270 (9)
H8A	0.2375	0.8914	0.5469	0.032*
H8B	0.2811	0.9705	0.5075	0.032*
C9	0.18688 (19)	1.0227 (3)	0.63391 (17)	0.0282 (9)
H9A	0.1570	1.0794	0.6469	0.034*
H9B	0.1593	0.9596	0.6347	0.034*

supplementary materials

C10	0.3642 (2)	1.0465 (3)	0.59343 (19)	0.0340 (10)
H10A	0.3824	1.0603	0.5525	0.041*
H10B	0.4036	1.0408	0.6216	0.041*
C11	0.15273 (18)	1.0551 (3)	0.52472 (17)	0.0239 (9)
C12	0.10965 (18)	0.9675 (3)	0.50235 (17)	0.0243 (9)
C13	0.07214 (18)	0.9841 (3)	0.44859 (17)	0.0271 (9)
H13A	0.0749	1.0479	0.4288	0.033*
C14	0.03137 (19)	0.9082 (3)	0.42450 (18)	0.0300 (9)
H14A	0.0066	0.9209	0.3883	0.036*
C15	0.02571 (18)	0.8139 (3)	0.45191 (17)	0.0241 (8)
C16	0.06142 (19)	0.7977 (3)	0.50545 (18)	0.0282 (9)
H16A	0.0571	0.7345	0.5256	0.034*
C17	0.10368 (19)	0.8731 (3)	0.53013 (18)	0.0266 (9)
H17A	0.1285	0.8598	0.5662	0.032*
C18	-0.02097 (19)	0.7331 (3)	0.42573 (17)	0.0248 (9)
H18A	-0.0668	0.7388	0.4450	0.030*
H18B	-0.0270	0.7465	0.3821	0.030*
C19	0.05284 (17)	0.5906 (3)	0.39739 (16)	0.0206 (8)
C20	0.12500 (18)	0.6035 (3)	0.31567 (16)	0.0242 (9)
C21	0.13385 (17)	0.4650 (3)	0.36843 (16)	0.0213 (8)
C22	0.08232 (17)	0.4945 (3)	0.40833 (16)	0.0203 (8)
C23	0.11642 (19)	0.3486 (3)	0.43759 (17)	0.0251 (9)
H23A	0.1209	0.2874	0.4599	0.030*
C24	0.21277 (18)	0.3110 (3)	0.35970 (18)	0.0279 (9)
H24A	0.2129	0.3250	0.3153	0.033*
C25	0.2812 (2)	0.3476 (4)	0.3856 (2)	0.0502 (13)
H25A	0.2864	0.4203	0.3779	0.075*
H25B	0.3191	0.3106	0.3664	0.075*
H25C	0.2821	0.3352	0.4292	0.075*
C26	0.2007 (2)	0.1985 (3)	0.3687 (2)	0.0460 (12)
H26A	0.1564	0.1796	0.3508	0.069*
H26B	0.2001	0.1831	0.4120	0.069*
H26C	0.2377	0.1602	0.3491	0.069*
C27	0.5087 (2)	1.1979 (4)	0.7415 (2)	0.0520 (14)
H27A	0.4751	1.2334	0.7683	0.062*
Cl11	0.49438 (8)	1.06661 (13)	0.74624 (7)	0.0720 (5)
Cl12	0.49582 (6)	1.23926 (13)	0.66603 (5)	0.0630 (4)
Cl13	0.59279 (6)	1.22798 (13)	0.76534 (6)	0.0635 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0316 (5)	0.0309 (6)	0.0291 (5)	0.0028 (4)	0.0083 (4)	0.0085 (4)
O1	0.0316 (15)	0.0228 (15)	0.0416 (18)	0.0013 (12)	-0.0007 (12)	0.0059 (13)
N1	0.0249 (16)	0.0199 (17)	0.0242 (17)	-0.0002 (13)	0.0038 (13)	0.0003 (14)
N2	0.0203 (15)	0.0249 (17)	0.0224 (17)	0.0011 (13)	0.0007 (13)	0.0029 (14)
N3	0.0206 (15)	0.0228 (17)	0.0216 (17)	0.0001 (13)	0.0018 (13)	-0.0025 (14)
N4	0.0219 (16)	0.0286 (19)	0.0194 (17)	0.0041 (13)	0.0006 (13)	-0.0001 (14)

N5	0.0247 (16)	0.0278 (19)	0.0243 (18)	0.0027 (14)	-0.0008 (14)	-0.0014 (15)
C1	0.025 (2)	0.025 (2)	0.026 (2)	0.0004 (16)	0.0031 (16)	-0.0034 (17)
C2	0.033 (2)	0.026 (2)	0.033 (2)	-0.0035 (18)	0.0026 (18)	0.0040 (18)
C3	0.032 (2)	0.029 (2)	0.035 (2)	-0.0052 (18)	-0.0028 (18)	-0.0020 (19)
C4	0.040 (2)	0.035 (3)	0.031 (2)	-0.0030 (19)	0.0001 (19)	-0.0044 (19)
C5	0.038 (2)	0.031 (2)	0.027 (2)	-0.0051 (19)	0.0010 (18)	-0.0044 (19)
C6	0.041 (2)	0.032 (2)	0.035 (3)	-0.0031 (19)	-0.0089 (19)	0.000 (2)
C7	0.026 (2)	0.030 (2)	0.034 (2)	0.0021 (17)	0.0001 (17)	-0.0043 (18)
C8	0.027 (2)	0.027 (2)	0.027 (2)	0.0045 (17)	0.0045 (16)	-0.0048 (17)
C9	0.029 (2)	0.027 (2)	0.029 (2)	-0.0016 (17)	0.0082 (17)	-0.0007 (18)
C10	0.025 (2)	0.043 (3)	0.034 (2)	-0.0083 (19)	0.0008 (18)	-0.003 (2)
C11	0.0235 (19)	0.021 (2)	0.028 (2)	0.0037 (16)	0.0087 (16)	0.0004 (17)
C12	0.0178 (18)	0.021 (2)	0.034 (2)	0.0045 (15)	0.0049 (16)	-0.0014 (18)
C13	0.0251 (19)	0.026 (2)	0.030 (2)	0.0045 (17)	0.0018 (17)	0.0063 (18)
C14	0.025 (2)	0.037 (3)	0.029 (2)	0.0029 (18)	-0.0020 (17)	0.0032 (19)
C15	0.0221 (19)	0.025 (2)	0.025 (2)	0.0013 (16)	0.0078 (15)	-0.0009 (17)
C16	0.029 (2)	0.029 (2)	0.027 (2)	-0.0014 (17)	0.0030 (17)	0.0037 (18)
C17	0.029 (2)	0.022 (2)	0.029 (2)	-0.0009 (16)	-0.0039 (17)	0.0028 (17)
C18	0.0267 (19)	0.025 (2)	0.023 (2)	0.0044 (16)	0.0016 (16)	0.0000 (17)
C19	0.0153 (17)	0.021 (2)	0.025 (2)	-0.0035 (15)	-0.0049 (15)	-0.0059 (17)
C20	0.0238 (19)	0.028 (2)	0.021 (2)	0.0000 (16)	0.0008 (16)	0.0018 (17)
C21	0.0197 (18)	0.025 (2)	0.020 (2)	-0.0020 (15)	-0.0032 (15)	-0.0002 (17)
C22	0.0219 (18)	0.021 (2)	0.0178 (19)	-0.0035 (15)	-0.0017 (14)	-0.0018 (16)
C23	0.030 (2)	0.026 (2)	0.020 (2)	0.0019 (16)	0.0021 (16)	0.0025 (17)
C24	0.0207 (19)	0.037 (2)	0.026 (2)	0.0063 (17)	0.0015 (16)	-0.0011 (18)
C25	0.028 (2)	0.070 (4)	0.053 (3)	0.012 (2)	-0.006 (2)	-0.027 (3)
C26	0.048 (3)	0.036 (3)	0.054 (3)	0.011 (2)	0.023 (2)	0.007 (2)
C27	0.035 (2)	0.087 (4)	0.034 (3)	0.009 (3)	-0.003 (2)	-0.013 (3)
Cl11	0.0769 (10)	0.0811 (11)	0.0581 (9)	0.0080 (8)	0.0082 (7)	-0.0164 (8)
Cl12	0.0379 (7)	0.1163 (13)	0.0348 (7)	0.0103 (7)	-0.0040 (5)	-0.0052 (7)
Cl13	0.0324 (6)	0.1206 (13)	0.0377 (7)	0.0056 (7)	-0.0026 (5)	-0.0095 (7)

Geometric parameters (Å, °)

Cl1—C20	1.762 (4)	C8—H8A	0.990
O1—C11	1.219 (4)	C8—H8B	0.990
N1—C19	1.349 (5)	C9—H9A	0.990
N1—C18	1.445 (5)	C9—H9B	0.990
N1—H1A	0.880	C10—H10A	0.990
N2—C20	1.341 (5)	C10—H10B	0.990
N2—C19	1.357 (5)	C11—C12	1.511 (5)
N3—C20	1.316 (5)	C12—C17	1.392 (5)
N3—C21	1.353 (5)	C12—C13	1.413 (5)
N4—C23	1.376 (5)	C13—C14	1.384 (5)
N4—C21	1.383 (5)	C13—H13A	0.950
N4—C24	1.493 (5)	C14—C15	1.388 (5)
N5—C23	1.320 (5)	C14—H14A	0.950
N5—C22	1.388 (5)	C15—C16	1.390 (5)
C1—C11	1.532 (5)	C15—C18	1.515 (5)

supplementary materials

C1—C2	1.535 (5)	C16—C17	1.401 (5)
C1—C8	1.552 (5)	C16—H16A	0.950
C1—C9	1.574 (5)	C17—H17A	0.950
C2—C3	1.538 (6)	C18—H18A	0.990
C2—H2B	0.990	C18—H18B	0.990
C2—H2C	0.990	C19—C22	1.411 (5)
C3—C4	1.531 (6)	C21—C22	1.391 (5)
C3—C10	1.551 (6)	C23—H23A	0.950
C3—H3B	1.000	C24—C26	1.515 (6)
C4—C5	1.548 (6)	C24—C25	1.526 (6)
C4—H4A	0.990	C24—H24A	1.000
C4—H4B	0.990	C25—H25A	0.980
C5—C6	1.531 (6)	C25—H25B	0.980
C5—C9	1.546 (6)	C25—H25C	0.980
C5—H5B	1.000	C26—H26A	0.980
C6—C7	1.539 (6)	C26—H26B	0.980
C6—H6A	0.990	C26—H26C	0.980
C6—H6B	0.990	C27—C11	1.756 (6)
C7—C10	1.535 (6)	C27—C113	1.763 (5)
C7—C8	1.537 (5)	C27—C112	1.775 (5)
C7—H7A	1.000	C27—H27A	1.000
C19—N1—C18	122.2 (3)	H10A—C10—H10B	108.2
C19—N1—H1A	118.9	O1—C11—C12	117.8 (3)
C18—N1—H1A	118.9	O1—C11—C1	118.7 (3)
C20—N2—C19	116.7 (3)	C12—C11—C1	123.5 (3)
C20—N3—C21	109.1 (3)	C17—C12—C13	117.9 (3)
C23—N4—C21	105.4 (3)	C17—C12—C11	125.8 (3)
C23—N4—C24	129.4 (3)	C13—C12—C11	116.3 (3)
C21—N4—C24	125.1 (3)	C14—C13—C12	120.5 (4)
C23—N5—C22	104.1 (3)	C14—C13—H13A	119.7
C11—C1—C2	109.4 (3)	C12—C13—H13A	119.7
C11—C1—C8	112.4 (3)	C13—C14—C15	121.7 (4)
C2—C1—C8	107.7 (3)	C13—C14—H14A	119.2
C11—C1—C9	110.0 (3)	C15—C14—H14A	119.2
C2—C1—C9	107.3 (3)	C14—C15—C16	118.1 (4)
C8—C1—C9	109.9 (3)	C14—C15—C18	120.7 (3)
C1—C2—C3	111.8 (3)	C16—C15—C18	121.2 (3)
C1—C2—H2B	109.3	C15—C16—C17	121.1 (4)
C3—C2—H2B	109.3	C15—C16—H16A	119.4
C1—C2—H2C	109.3	C17—C16—H16A	119.4
C3—C2—H2C	109.3	C12—C17—C16	120.7 (4)
H2B—C2—H2C	107.9	C12—C17—H17A	119.7
C4—C3—C2	109.6 (3)	C16—C17—H17A	119.7
C4—C3—C10	109.1 (3)	N1—C18—C15	114.4 (3)
C2—C3—C10	108.9 (3)	N1—C18—H18A	108.7
C4—C3—H3B	109.8	C15—C18—H18A	108.7
C2—C3—H3B	109.8	N1—C18—H18B	108.7
C10—C3—H3B	109.8	C15—C18—H18B	108.7
C3—C4—C5	109.6 (3)	H18A—C18—H18B	107.6

C3—C4—H4A	109.8	N1—C19—N2	118.8 (3)
C5—C4—H4A	109.8	N1—C19—C22	122.5 (3)
C3—C4—H4B	109.8	N2—C19—C22	118.8 (3)
C5—C4—H4B	109.8	N3—C20—N2	131.9 (3)
H4A—C4—H4B	108.2	N3—C20—C11	114.8 (3)
C6—C5—C9	110.0 (3)	N2—C20—C11	113.3 (3)
C6—C5—C4	109.3 (3)	N3—C21—N4	126.1 (3)
C9—C5—C4	109.2 (3)	N3—C21—C22	127.8 (3)
C6—C5—H5B	109.4	N4—C21—C22	106.1 (3)
C9—C5—H5B	109.4	N5—C22—C21	110.4 (3)
C4—C5—H5B	109.4	N5—C22—C19	133.8 (3)
C5—C6—C7	110.1 (3)	C21—C22—C19	115.7 (3)
C5—C6—H6A	109.6	N5—C23—N4	113.9 (3)
C7—C6—H6A	109.6	N5—C23—H23A	123.0
C5—C6—H6B	109.6	N4—C23—H23A	123.0
C7—C6—H6B	109.6	N4—C24—C26	110.1 (3)
H6A—C6—H6B	108.2	N4—C24—C25	109.2 (3)
C10—C7—C8	109.2 (3)	C26—C24—C25	113.3 (4)
C10—C7—C6	109.4 (3)	N4—C24—H24A	108.0
C8—C7—C6	110.0 (3)	C26—C24—H24A	108.0
C10—C7—H7A	109.4	C25—C24—H24A	108.0
C8—C7—H7A	109.4	C24—C25—H25A	109.5
C6—C7—H7A	109.4	C24—C25—H25B	109.5
C7—C8—C1	110.4 (3)	H25A—C25—H25B	109.5
C7—C8—H8A	109.6	C24—C25—H25C	109.5
C1—C8—H8A	109.6	H25A—C25—H25C	109.5
C7—C8—H8B	109.6	H25B—C25—H25C	109.5
C1—C8—H8B	109.6	C24—C26—H26A	109.5
H8A—C8—H8B	108.1	C24—C26—H26B	109.5
C5—C9—C1	109.6 (3)	H26A—C26—H26B	109.5
C5—C9—H9A	109.7	C24—C26—H26C	109.5
C1—C9—H9A	109.7	H26A—C26—H26C	109.5
C5—C9—H9B	109.7	H26B—C26—H26C	109.5
C1—C9—H9B	109.7	C111—C27—C113	110.5 (3)
H9A—C9—H9B	108.2	C111—C27—C112	109.7 (3)
C7—C10—C3	109.4 (3)	C113—C27—C112	110.1 (3)
C7—C10—H10A	109.8	C111—C27—H27A	108.8
C3—C10—H10A	109.8	C113—C27—H27A	108.8
C7—C10—H10B	109.8	C112—C27—H27A	108.8
C3—C10—H10B	109.8		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots N5 ⁱ	0.88	2.22	3.013 (5)	150
C27—H27A \cdots N2 ⁱⁱ	1.00	2.59	3.553 (6)	161
C5—H5B \cdots N3 ⁱⁱⁱ	1.00	2.66	3.641 (5)	166
C23—H23A \cdots O1 ^{iv}	0.95	2.23	3.179 (5)	175

supplementary materials

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+1/2, -y+2, z+1/2$; (iii) $x, -y+3/2, z+1/2$; (iv) $x, y-1, z$.

Fig. 1




Fig. 2

