

# {*N,N'*-Bis[(*E*)-3-phenylprop-2-en-1-ylidene]propane-1,3-diamine- $\kappa^2$ *N,N'*]-dichloridocobalt(II)}

Morteza Montazerzohori,<sup>a</sup> Mohammad Hossein Habibi,<sup>b\*</sup> Mehdi Amirnasr,<sup>c</sup> Keita Ariyoshi<sup>d</sup> and Takayoshi Suzuki<sup>d</sup>

<sup>a</sup>Department of Chemistry, Yasouj University, Yasouj 75914-353, Iran, <sup>b</sup>Catalysis Division, Department of Chemistry, University of Isfahan, Isfahan 81746-73441, Iran, <sup>c</sup>Department of Chemistry, Isfahan University of Technology, Isfahan, Iran, and <sup>d</sup>Department of Chemistry, Faculty of Science, Okayama University, Tushima-naka 3-1-1, Okayama 700-8530, Japan  
Correspondence e-mail: mhhabibi@yahoo.com


Received 27 April 2009; accepted 30 April 2009

Key indicators: single-crystal X-ray study;  $T = 193$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.036;  $wR$  factor = 0.091; data-to-parameter ratio = 24.7.

The  $\text{Co}^{\text{II}}$  atom in the title monomeric Schiff base complex,  $[\text{CoCl}_2(\text{C}_{21}\text{H}_{22}\text{N}_2)]$ , is bonded to two Cl atoms and to two N atoms of the Schiff base ligand *N,N'*-bis[(*E*)-3-phenylprop-2-en-1-ylidene]propane-1,3-diamine in a distorted tetrahedral geometry. The molecule has an idealised mirror symmetry, but is not located on a crystallographic mirror plane.

## Related literature

For transition metal complexes with Schiff base ligands, see: Yamada (1999). For related structures, see: Amirnasr *et al.* (2003); Blonk *et al.* (1985); Habibi *et al.* (2007*a,b*); Meghdadi *et al.* (2002); Scheidt *et al.* (1969).



## Experimental

### Crystal data

$[\text{CoCl}_2(\text{C}_{21}\text{H}_{22}\text{N}_2)]$   
 $M_r = 432.24$   
Monoclinic,  $P2_1/n$

$a = 7.4976$  (5) Å  
 $b = 16.1594$  (8) Å  
 $c = 16.6238$  (10) Å

$\beta = 91.531$  (2)°  
 $V = 2013.4$  (2) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 1.13$  mm<sup>-1</sup>  
 $T = 193$  K  
 $0.30 \times 0.30 \times 0.20$  mm

### Data collection

Rigaku R-Axis RAPID diffractometer  
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)  
 $T_{\text{min}} = 0.729$ ,  $T_{\text{max}} = 0.806$

23596 measured reflections  
5826 independent reflections  
4806 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.091$   
 $S = 1.09$   
5826 reflections

236 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.53$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.46$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Co1—N1	2.0368 (13)	Co1—Cl1	2.2399 (5)
Co1—N2	2.0392 (13)	Co1—Cl2	2.2559 (5)
N1—Co1—N2	93.21 (5)	N1—Co1—Cl2	106.31 (4)
N1—Co1—Cl1	117.36 (4)	N2—Co1—Cl2	106.71 (4)
N2—Co1—Cl1	118.07 (4)	Cl1—Co1—Cl2	112.965 (19)

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2004); program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Partial support of this work by Yasouj University is acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2939).

## References

- Amirnasr, M., Schenk, K. J., Salavati, M., Dehghanpour, S., Taeb, A. & Tadjarodi, A. (2003). *J. Coord. Chem.* **56**, 231–243.  
Blonk, H. I., Driessen, W. L. & Reedijk, J. (1985). *J. Chem. Soc. Dalton Trans.* pp. 1699–1704.  
Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Habibi, M. H., Lalegani, A., Mokhtari, R. & Suzuki, T. (2007*a*). *Acta Cryst.* **E63**, m2472.  
Habibi, M. H., Lalegani, A., Mokhtari, R. & Suzuki, T. (2007*b*). *Acta Cryst.* **E63**, m2580.  
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
Meghdadi, S., Amirnasr, M., Schenk, K. J. & Dehghanpour, S. (2002). *Helv. Chim. Acta*, **85**, 2807–2816.  
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.  
Rigaku/MS (2004). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.  
Scheidt, W. R., Hanson, J. C. & Rasmussen, P. G. (1969). *Inorg. Chem.* **8**, 2398–2401.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Yamada, S. (1999). *Coord. Chem. Rev.* **190**, 537–555.

**supplementary materials**

*Acta Cryst.* (2009). E65, m617 [ doi:10.1107/S1600536809016274 ]

**{*N,N'*-Bis[(*E*)-3-phenylprop-2-en-1-ylidene]propane-1,3-diamine- $\kappa^2$ *N,N'*]dichloridocobalt(II)}**

**M. Montazerzohori, M. H. Habibi, M. Amirnasr, K. Ariyoshi and T. Suzuki**

**Comment**

Transition metal complexes with Schiff base ligands have attracted substantial interest for many years (Yamada, 1999). Cinnamaldehyde and its substituted derivatives condense with diamines to supply a range of Schiff base compounds; a small number of such bis(cinnamaldehyde)ethylenediimine ligands have been used to prepare adducts with transition metals. Among such complexes whose structures have been described are, for example, the copper(I) iodide (Habibi *et al.*, 2007a), (triphenylphosphine)(halogen/pseudohalogeno)-copper(I) (Habibi *et al.*, 2007b), copper(I) perchlorate (Meghdadi *et al.*, 2002), and the cobalt(II) chloride, cobalt(II) bromide and nickel bromide (Amirnasr *et al.*, 2003) adducts. The title complex, (I), was prepared by the reaction of  $\text{CoCl}_2$  with the bidentate ligand *N,N'*-bis[(*E*)-3-phenylprop-2-en-1-ylidene]propane-1,3-diamine (*ca*<sub>2</sub>pn). The molecular structure of complex (I) and the *ORTEP* structure are shown in Fig. 1. The metal centre has a tetrahedral coordination which shows significant distortion, mainly due to the presence of the six-membered chelate ring (Table 1): the endocyclic N1—Co1—N2 angle is much narrower than the ideal tetrahedral angle of 109.5° whereas the opposite Cl1—Co1—Cl2 angle is much wider than the ideal tetrahedral angle. The Co1—Cl1 and Co1—Cl2 bond lengths are in good agreement with Co—Cl distances in other tetrahedral cobalt complexes, *e.g.* 2.229 (3) Å in Co(ethylenedimorpholine)Cl<sub>2</sub> (Scheidt *et al.*, 1969), and 2.2434 (8) and 2.2266 (8) Å in Co[*N,N*-bis(3,5-dimethylpyrazol-1-ylmethyl)-aminobenzene]Cl<sub>2</sub> (Blonk *et al.*, 1985).  $\pi$ -Conjugation within the azadiene fragments is consistent with the observed pattern of C—C bond distances; the predominantly double C7=C8 and C14=C15 bonds are substantially shorter than the C8—C9 and C13—C14 bonds, which have a significant  $\pi$ -component; the latter bonds in their turn are much shorter than the single C10—C11 and C11—C12 bonds in the propylene bridge.

**Experimental**

The bidentate Schiff base ligand of *N,N'*-bis[(*E*)-3-phenyl-propenylidene]-1,3-diaminopropane was synthesized by the condensation reaction of 2 mmol of (*E*)-3-phenylpropenal and 1 mmol 1,3-diaminopropane in 10 ml dichloromethane in an ice bath for 1 h. The solution then was added drop wise to a solution of 1 mmol anhydrous  $\text{CoCl}_2$  in 10 ml dichloromethane under nitrogen atmosphere. The mixture was stirred for 3 h and then filtered. To the filtrate, 20 ml chloroform was added and kept overnight. The crystals suitable for X-ray were filtered off and washed with chloroform (68% yield). Elemental analysis for  $\text{C}_{21}\text{H}_{22}\text{Cl}_2\text{CoN}_2$ : Calcd.: C, 58.35; H, 5.13; N, 6.48; Found: C, 58.31; H, 5.11; N, 6.42.

**Refinement**

All H atoms were placed in calculated positions and refined using a riding-model, with C—H = 0.95–0.99 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

## Figures




Fig. 1. A view of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.

## {*N,N'*-Bis[(*E*)-3-phenylprop-2-en-1-ylidene]propane-1,3-diamine- $\kappa^2$ *N,N'*}dichloridocobalt(II)

### Crystal data

[CoCl<sub>2</sub>(C<sub>21</sub>H<sub>22</sub>N<sub>2</sub>)]

$M_r = 432.24$

Monoclinic,  $P2_1/n$

$a = 7.4976$  (5) Å

$b = 16.1594$  (8) Å

$c = 16.6238$  (10) Å

$\beta = 91.531$  (2)°

$V = 2013.4$  (2) Å<sup>3</sup>

$Z = 4$

$F_{000} = 892$

$D_x = 1.426$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71075$  Å

Cell parameters from 16882 reflections

$\theta = 3.0$ – $29.9$ °

$\mu = 1.13$  mm<sup>-1</sup>

$T = 193$  K

Cubic, green

$0.30 \times 0.30 \times 0.20$  mm

### Data collection

Rigaku R-Axis RAPID  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 10.00 pixels mm<sup>-1</sup>

$T = 193$  K

$\omega$  scans

Absorption correction: multi-scan  
(ABSCOR; Higashi, 1995)

$T_{\min} = 0.729$ ,  $T_{\max} = 0.806$

23596 measured reflections

5826 independent reflections

4806 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\max} = 30.0$ °

$\theta_{\min} = 3.0$ °

$h = -10 \rightarrow 10$

$k = -22 \rightarrow 22$

$l = -23 \rightarrow 23$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.091$

$S = 1.09$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0428P)^2 + 0.5035P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

5826 reflections  $\Delta\rho_{\max} = 0.53 \text{ e } \text{\AA}^{-3}$   
 236 parameters  $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.25070 (3)	0.193360 (13)	0.243448 (12)	0.02719 (7)
Cl1	0.28692 (6)	0.05592 (2)	0.25023 (2)	0.03655 (10)
Cl2	0.49842 (6)	0.26032 (3)	0.20514 (3)	0.03852 (11)
N1	0.04670 (18)	0.23684 (8)	0.17195 (8)	0.0291 (3)
N2	0.16292 (18)	0.25318 (8)	0.34289 (8)	0.0291 (3)
C1	-0.2001 (2)	-0.01609 (10)	0.05017 (9)	0.0304 (3)
C2	-0.1150 (2)	-0.07521 (11)	0.09975 (11)	0.0377 (4)
H2	-0.0448	-0.0578	0.1451	0.045*
C3	-0.1329 (3)	-0.15844 (12)	0.08302 (12)	0.0426 (4)
H3	-0.0745	-0.1981	0.1167	0.051*
C4	-0.2360 (3)	-0.18448 (11)	0.01710 (12)	0.0419 (4)
H4	-0.2488	-0.2419	0.0063	0.050*
C5	-0.3203 (2)	-0.12722 (10)	-0.03290 (11)	0.0364 (4)
H5	-0.3900	-0.1452	-0.0781	0.044*
C6	-0.3020 (2)	-0.04375 (10)	-0.01645 (10)	0.0330 (3)
H6	-0.3595	-0.0045	-0.0509	0.040*
C7	-0.1865 (2)	0.07272 (11)	0.06595 (10)	0.0325 (3)
H7	-0.2680	0.1076	0.0373	0.039*
C8	-0.0703 (2)	0.10972 (10)	0.11696 (9)	0.0304 (3)
H8	0.0122	0.0768	0.1473	0.037*
C9	-0.0678 (2)	0.19798 (10)	0.12681 (10)	0.0320 (3)
H9	-0.1558	0.2296	0.0984	0.038*
C10	0.0290 (2)	0.32742 (10)	0.17794 (10)	0.0342 (3)
H10A	-0.0620	0.3469	0.1380	0.041*
H10B	0.1440	0.3538	0.1653	0.041*
C11	-0.0252 (2)	0.35316 (10)	0.26220 (10)	0.0340 (3)
H11A	-0.0595	0.4123	0.2608	0.041*
H11B	-0.1321	0.3210	0.2767	0.041*

## supplementary materials

---

C12	0.1177 (2)	0.34122 (9)	0.32822 (10)	0.0342 (3)
H12A	0.2269	0.3712	0.3131	0.041*
H12B	0.0753	0.3659	0.3788	0.041*
C13	0.1347 (2)	0.22441 (10)	0.41378 (9)	0.0299 (3)
H13	0.0819	0.2604	0.4515	0.036*
C14	0.1777 (2)	0.14149 (10)	0.43997 (9)	0.0301 (3)
H14	0.2322	0.1041	0.4041	0.036*
C15	0.1409 (2)	0.11698 (10)	0.51492 (10)	0.0299 (3)
H15	0.0783	0.1550	0.5474	0.036*
C16	0.1881 (2)	0.03716 (10)	0.55142 (9)	0.0303 (3)
C17	0.1433 (3)	0.02236 (11)	0.63139 (10)	0.0383 (4)
H17	0.0757	0.0621	0.6596	0.046*
C18	0.1975 (3)	-0.05030 (12)	0.66951 (12)	0.0470 (5)
H18	0.1663	-0.0600	0.7237	0.056*
C19	0.2957 (3)	-0.10810 (12)	0.62942 (13)	0.0460 (5)
H19	0.3343	-0.1572	0.6561	0.055*
C20	0.3386 (2)	-0.09457 (11)	0.54975 (13)	0.0420 (4)
H20	0.4057	-0.1348	0.5219	0.050*
C21	0.2844 (2)	-0.02314 (10)	0.51072 (11)	0.0345 (3)
H21	0.3126	-0.0149	0.4559	0.041*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.02769 (12)	0.02684 (12)	0.02693 (12)	0.00036 (7)	-0.00153 (8)	0.00003 (8)
C11	0.0445 (2)	0.02843 (19)	0.0363 (2)	0.00604 (15)	-0.00552 (17)	-0.00140 (15)
C12	0.0328 (2)	0.0447 (2)	0.0381 (2)	-0.00778 (17)	0.00149 (16)	0.00277 (18)
N1	0.0300 (7)	0.0313 (6)	0.0258 (6)	0.0031 (5)	-0.0003 (5)	-0.0008 (5)
N2	0.0313 (7)	0.0270 (6)	0.0287 (6)	-0.0016 (5)	-0.0015 (5)	-0.0009 (5)
C1	0.0258 (7)	0.0354 (8)	0.0300 (7)	-0.0005 (6)	0.0016 (6)	-0.0011 (6)
C2	0.0357 (9)	0.0430 (9)	0.0344 (8)	-0.0002 (7)	-0.0020 (7)	0.0037 (7)
C3	0.0413 (10)	0.0407 (10)	0.0461 (10)	0.0037 (8)	0.0050 (8)	0.0120 (8)
C4	0.0403 (10)	0.0341 (9)	0.0518 (11)	-0.0037 (7)	0.0119 (8)	-0.0011 (8)
C5	0.0299 (8)	0.0413 (9)	0.0381 (9)	-0.0053 (7)	0.0050 (7)	-0.0070 (7)
C6	0.0288 (8)	0.0394 (8)	0.0308 (8)	0.0007 (6)	0.0003 (6)	0.0001 (7)
C7	0.0285 (8)	0.0363 (8)	0.0324 (8)	0.0030 (6)	-0.0030 (6)	-0.0008 (7)
C8	0.0274 (7)	0.0356 (8)	0.0283 (7)	0.0021 (6)	0.0002 (6)	-0.0009 (6)
C9	0.0282 (8)	0.0388 (9)	0.0288 (7)	0.0036 (6)	-0.0016 (6)	-0.0010 (6)
C10	0.0386 (9)	0.0302 (8)	0.0335 (8)	0.0038 (7)	-0.0050 (7)	0.0028 (7)
C11	0.0364 (9)	0.0265 (7)	0.0390 (9)	0.0030 (6)	-0.0007 (7)	-0.0016 (7)
C12	0.0450 (10)	0.0237 (7)	0.0339 (8)	-0.0014 (6)	-0.0025 (7)	-0.0018 (6)
C13	0.0297 (8)	0.0315 (8)	0.0285 (7)	0.0010 (6)	-0.0012 (6)	-0.0030 (6)
C14	0.0307 (8)	0.0319 (7)	0.0277 (7)	0.0000 (6)	-0.0007 (6)	-0.0016 (6)
C15	0.0288 (8)	0.0325 (8)	0.0282 (7)	-0.0005 (6)	-0.0016 (6)	-0.0024 (6)
C16	0.0273 (8)	0.0332 (8)	0.0302 (7)	-0.0047 (6)	-0.0041 (6)	0.0007 (6)
C17	0.0447 (10)	0.0380 (9)	0.0322 (8)	-0.0032 (7)	-0.0010 (7)	0.0012 (7)
C18	0.0596 (13)	0.0467 (11)	0.0343 (9)	-0.0085 (9)	-0.0065 (8)	0.0105 (8)
C19	0.0425 (10)	0.0384 (9)	0.0563 (12)	-0.0044 (8)	-0.0131 (9)	0.0140 (9)

C20	0.0330 (9)	0.0337 (9)	0.0593 (12)	-0.0006 (7)	-0.0008 (8)	0.0020 (8)
C21	0.0310 (8)	0.0341 (8)	0.0384 (9)	-0.0038 (6)	0.0015 (6)	0.0022 (7)

*Geometric parameters (Å, °)*

Co1—N1	2.0368 (13)	C10—C11	1.527 (2)
Co1—N2	2.0392 (13)	C10—H10A	0.9900
Co1—C11	2.2399 (5)	C10—H10B	0.9900
Co1—C12	2.2559 (5)	C11—C12	1.525 (2)
N1—C9	1.289 (2)	C11—H11A	0.9900
N1—C10	1.473 (2)	C11—H11B	0.9900
N2—C13	1.289 (2)	C12—H12A	0.9900
N2—C12	1.481 (2)	C12—H12B	0.9900
C1—C6	1.402 (2)	C13—C14	1.443 (2)
C1—C2	1.404 (2)	C13—H13	0.9500
C1—C7	1.462 (2)	C14—C15	1.343 (2)
C2—C3	1.379 (3)	C14—H14	0.9500
C2—H2	0.9500	C15—C16	1.465 (2)
C3—C4	1.389 (3)	C15—H15	0.9500
C3—H3	0.9500	C16—C21	1.398 (2)
C4—C5	1.385 (3)	C16—C17	1.401 (2)
C4—H4	0.9500	C17—C18	1.390 (2)
C5—C6	1.382 (2)	C17—H17	0.9500
C5—H5	0.9500	C18—C19	1.373 (3)
C6—H6	0.9500	C18—H18	0.9500
C7—C8	1.340 (2)	C19—C20	1.389 (3)
C7—H7	0.9500	C19—H19	0.9500
C8—C9	1.436 (2)	C20—C21	1.380 (2)
C8—H8	0.9500	C20—H20	0.9500
C9—H9	0.9500	C21—H21	0.9500
N1—Co1—N2	93.21 (5)	N1—C10—H10B	109.4
N1—Co1—C11	117.36 (4)	C11—C10—H10B	109.4
N2—Co1—C11	118.07 (4)	H10A—C10—H10B	108.0
N1—Co1—C12	106.31 (4)	C12—C11—C10	115.27 (14)
N2—Co1—C12	106.71 (4)	C12—C11—H11A	108.5
C11—Co1—C12	112.965 (19)	C10—C11—H11A	108.5
C9—N1—C10	117.61 (13)	C12—C11—H11B	108.5
C9—N1—Co1	130.55 (11)	C10—C11—H11B	108.5
C10—N1—Co1	111.78 (10)	H11A—C11—H11B	107.5
C13—N2—C12	117.02 (13)	N2—C12—C11	113.16 (13)
C13—N2—Co1	129.36 (11)	N2—C12—H12A	108.9
C12—N2—Co1	113.53 (10)	C11—C12—H12A	108.9
C6—C1—C2	118.44 (15)	N2—C12—H12B	108.9
C6—C1—C7	119.26 (14)	C11—C12—H12B	108.9
C2—C1—C7	122.30 (15)	H12A—C12—H12B	107.8
C3—C2—C1	120.34 (16)	N2—C13—C14	124.76 (15)
C3—C2—H2	119.8	N2—C13—H13	117.6
C1—C2—H2	119.8	C14—C13—H13	117.6
C2—C3—C4	120.22 (17)	C15—C14—C13	120.29 (15)

## supplementary materials

---

C2—C3—H3	119.9	C15—C14—H14	119.9
C4—C3—H3	119.9	C13—C14—H14	119.9
C5—C4—C3	120.42 (17)	C14—C15—C16	126.20 (15)
C5—C4—H4	119.8	C14—C15—H15	116.9
C3—C4—H4	119.8	C16—C15—H15	116.9
C6—C5—C4	119.47 (16)	C21—C16—C17	118.72 (16)
C6—C5—H5	120.3	C21—C16—C15	122.38 (15)
C4—C5—H5	120.3	C17—C16—C15	118.82 (15)
C5—C6—C1	121.10 (16)	C18—C17—C16	120.16 (18)
C5—C6—H6	119.4	C18—C17—H17	119.9
C1—C6—H6	119.4	C16—C17—H17	119.9
C8—C7—C1	126.34 (15)	C19—C18—C17	120.49 (19)
C8—C7—H7	116.8	C19—C18—H18	119.8
C1—C7—H7	116.8	C17—C18—H18	119.8
C7—C8—C9	121.45 (15)	C18—C19—C20	119.78 (17)
C7—C8—H8	119.3	C18—C19—H19	120.1
C9—C8—H8	119.3	C20—C19—H19	120.1
N1—C9—C8	123.85 (15)	C21—C20—C19	120.49 (18)
N1—C9—H9	118.1	C21—C20—H20	119.8
C8—C9—H9	118.1	C19—C20—H20	119.8
N1—C10—C11	111.07 (13)	C20—C21—C16	120.33 (17)
N1—C10—H10A	109.4	C20—C21—H21	119.8
C11—C10—H10A	109.4	C16—C21—H21	119.8
N2—Co1—N1—C9	-125.65 (15)	C10—N1—C9—C8	-178.17 (15)
Cl1—Co1—N1—C9	-1.63 (16)	Co1—N1—C9—C8	-1.2 (3)
Cl2—Co1—N1—C9	125.90 (14)	C7—C8—C9—N1	-176.41 (17)
N2—Co1—N1—C10	51.43 (11)	C9—N1—C10—C11	112.10 (16)
Cl1—Co1—N1—C10	175.45 (9)	Co1—N1—C10—C11	-65.40 (15)
Cl2—Co1—N1—C10	-57.02 (11)	N1—C10—C11—C12	69.65 (18)
N1—Co1—N2—C13	128.77 (14)	C13—N2—C12—C11	-118.87 (16)
Cl1—Co1—N2—C13	5.30 (16)	Co1—N2—C12—C11	58.15 (16)
Cl2—Co1—N2—C13	-123.14 (14)	C10—C11—C12—N2	-65.56 (19)
N1—Co1—N2—C12	-47.80 (11)	C12—N2—C13—C14	-177.22 (14)
Cl1—Co1—N2—C12	-171.27 (9)	Co1—N2—C13—C14	6.3 (2)
Cl2—Co1—N2—C12	60.29 (11)	N2—C13—C14—C15	-179.13 (16)
C6—C1—C2—C3	-0.3 (3)	C13—C14—C15—C16	-175.37 (14)
C7—C1—C2—C3	179.37 (17)	C14—C15—C16—C21	1.8 (3)
C1—C2—C3—C4	-0.3 (3)	C14—C15—C16—C17	178.41 (16)
C2—C3—C4—C5	0.7 (3)	C21—C16—C17—C18	1.4 (3)
C3—C4—C5—C6	-0.4 (3)	C15—C16—C17—C18	-175.39 (16)
C4—C5—C6—C1	-0.2 (3)	C16—C17—C18—C19	0.2 (3)
C2—C1—C6—C5	0.6 (2)	C17—C18—C19—C20	-1.2 (3)
C7—C1—C6—C5	-179.12 (15)	C18—C19—C20—C21	0.5 (3)
C6—C1—C7—C8	-166.72 (17)	C19—C20—C21—C16	1.1 (3)
C2—C1—C7—C8	13.6 (3)	C17—C16—C21—C20	-2.0 (2)
C1—C7—C8—C9	179.17 (16)	C15—C16—C21—C20	174.62 (16)

Fig. 1

