

Poly[[μ_4 -tartrato-cadmium(II)] 0.167-hydrate]Li-Zhen Zhao,^a Ping Li,^a Bao-Liang Cao^a and Seik Weng Ng^{b*}^aDepartment of Chemistry, Jining Normal College, Wulanchabu, Inner Mongolia 012000, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 1 May 2009; accepted 5 May 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.022; wR factor = 0.060; data-to-parameter ratio = 13.3.

The title compound, $\{[\text{Cd}(\text{C}_4\text{H}_4\text{O}_6)] \cdot 0.167\text{H}_2\text{O}\}_n$, adopts a three-dimensional network structure in which each Cd^{II} ion is chelated by two pairs of carboxylate and hydroxyl O atoms from two tartrate anions, and is additionally linked to two O atoms of two carboxylate groups that are not involved in chelation. The asymmetric unit has four independent cadmium atoms, two of which lie on special positions of 2 site symmetry. The tartrate anions all lie on general positions. All hydroxyl groups are engaged in $\text{O}-\text{H} \cdots \text{O}$ hydrogen-bonds, one of which is also bifurcated. The non-coordinating water molecule is situated on a site with half-occupation.

Related literature

For the structure of cadmium tartrate trihydrate, see: González-Silgo *et al.* (1999).



Experimental

Crystal data

$[\text{Cd}(\text{C}_4\text{H}_4\text{O}_6)] \cdot 0.167\text{H}_2\text{O}$	$V = 3696.3$ (3) Å ³
$M_r = 263.47$	$Z = 24$
Orthorhombic, $C222_1$	Mo $K\alpha$ radiation
$a = 10.7901$ (4) Å	$\mu = 3.53$ mm ⁻¹
$b = 11.1995$ (5) Å	$T = 293$ K
$c = 30.588$ (1) Å	$0.37 \times 0.22 \times 0.15$ mm

Data collection

Bruker APEXII area-detector diffractometer	13041 measured reflections
Absorption correction: multi-scan (SADABS, Sheldrick, 1996)	4095 independent reflections
$T_{\text{min}} = 0.505$, $T_{\text{max}} = 0.780$	4073 reflections with $I > 2\sigma(I)$
(expected range = 0.382–0.589)	$R_{\text{int}} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.022$	H-atom parameters constrained
$wR(F^2) = 0.060$	$\Delta\rho_{\text{max}} = 1.34$ e Å ⁻³
$S = 1.02$	$\Delta\rho_{\text{min}} = -1.18$ e Å ⁻³
4095 reflections	Absolute structure: Flack (1983),
308 parameters	1733 Friedel pairs
6 restraints	Flack parameter: -0.02 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O3}-\text{H3} \cdots \text{O18}^{\text{i}}$	0.82	1.96	2.740 (4)	159
$\text{O4}-\text{H4} \cdots \text{O10}^{\text{ii}}$	0.82	2.50	3.236 (6)	149
$\text{O9}-\text{H9} \cdots \text{O11}^{\text{iii}}$	0.82	2.17	2.797 (5)	134
$\text{O9}-\text{H9} \cdots \text{O1w}$	0.82	2.12	2.68 (2)	125
$\text{O10}-\text{H10} \cdots \text{O15}^{\text{ii}}$	0.82	2.15	2.938 (4)	160
$\text{O15}-\text{H15} \cdots \text{O1}^{\text{iv}}$	0.82	2.13	2.717 (4)	128
$\text{O16}-\text{H16} \cdots \text{O7}^{\text{v}}$	0.82	1.84	2.609 (4)	155
$\text{O1W}-\text{H1W1} \cdots \text{O14}^{\text{iii}}$	0.82	2.26	3.034 (19)	157

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (iv) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (v) $x - 1, y, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We acknowledge support from the Scientific Research Projects of Higher Education of Inner Mongolia (NJzy08217) and the University of Malaya.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2796).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 González-Silgo, C., González-Platas, J., Ruiz-Pérez, C., López, T. & Torres, M. E. (1999). *Acta Cryst.* **C55**, 710–712.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). publCIF. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m629 [doi:10.1107/S1600536809016882]

Poly[[μ_4 -tartrato-cadmium(II)] 0.167-hydrate]

L.-Z. Zhao, P. Li, B.-L. Cao and S. W. Ng

Experimental

Cadmium chloride 2.5 hydrate (0.23 g), *R,R*-tartaric acid (0.48 g), sodium hydroxide (0.39 g), imidazole (0.12 g) and water (0.4 ml) were sealed in a 25-ml Teflon-lined stainless-steel vessel. This was heated at 393 K for 3 d. The colourless crystals found in the cooled vessel were picked out manually.

Refinement

C-bound H-atoms were placed in calculated positions (C-H = 0.98 Å) and were included in the refinement in the riding model approximation, with $U(H)$ fixed at $1.2U(C)$. The hydroxy H-atoms were generated geometrically by assuming an sp^3 type of hybridization (O-H = 0.82 Å); these were included in the refinement. At this stage, the difference Fourier map had a peak at about 1.5 Å from hydroxyl H9 atom and it was refined as a water molecule of half-site occupancy as the peak was 2.5 Å from the symmetry-related atom. The two water H atoms were placed in chemically sensible positions; only one of these H atom forms a hydrogen bond to an acceptor oxygen atom. The final difference Fourier map had a peak at 2 Å from O1W and hole at 1.2 Å from Cd1.

Figures




Fig. 1. Displacement ellipsoid plot (Barbour, 2001) of a portion of polymeric cadmium tartrate 1/6 hydrate at the 70% probability level; H atoms are drawn as spheres of arbitrary radius. The disordered water molecule is not shown.

Poly[[μ_4 -tartrato-cadmium(II)] 0.167-hydrate]

Crystal data

[Cd(C₄H₄O₆)]·0.167H₂O

$M_r = 263.47$

Orthorhombic, C222₁

Hall symbol: C 2c 2

$a = 10.7901$ (4) Å

$b = 11.1995$ (5) Å

$c = 30.588$ (1) Å

$V = 3696.3$ (3) Å³

$Z = 24$

$F_{000} = 3016$

$D_x = 2.841$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 4135 reflections

$\theta = 2.3$ – 27.2°

$\mu = 3.53$ mm⁻¹

$T = 293$ K

Block, colourless

$0.37 \times 0.22 \times 0.15$ mm

supplementary materials

Data collection

Bruker APEXII area-detector diffractometer diffractometer	4095 independent reflections
Radiation source: fine-focus sealed tube	4073 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.018$
$T = 293$ K	$\theta_{\text{max}} = 27.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.7^\circ$
Absorption correction: multi-scan (SADABS, Sheldrick, 1996)	$h = -13 \rightarrow 12$
$T_{\text{min}} = 0.505$, $T_{\text{max}} = 0.780$	$k = -14 \rightarrow 14$
13041 measured reflections	$l = -38 \rightarrow 39$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.022$	$w = 1/[\sigma^2(F_o^2) + (0.0411P)^2 + 9.1485P]$
$wR(F^2) = 0.060$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4095 reflections	$\Delta\rho_{\text{max}} = 1.34 \text{ e } \text{\AA}^{-3}$
308 parameters	$\Delta\rho_{\text{min}} = -1.18 \text{ e } \text{\AA}^{-3}$
6 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1733 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: -0.02 (2)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cd1	1.0000	0.70141 (4)	0.7500	0.01735 (8)	
Cd2	0.95531 (3)	0.81768 (3)	0.578406 (9)	0.02161 (7)	
Cd3	0.41091 (3)	0.79800 (2)	0.601933 (8)	0.01865 (7)	
Cd4	0.0000	0.19902 (4)	0.7500	0.03003 (11)	
O1	1.0079 (3)	0.7619 (3)	0.67881 (10)	0.0233 (6)	
O2	1.1079 (3)	0.8548 (3)	0.62512 (10)	0.0302 (7)	
O3	1.1495 (3)	0.8566 (3)	0.74078 (9)	0.0208 (6)	
H3	1.2130	0.8367	0.7536	0.031*	
O4	1.0508 (4)	1.0602 (3)	0.69188 (15)	0.0482 (11)	
H4	1.0337	1.0916	0.6684	0.072*	
O5	1.3661 (3)	1.0768 (3)	0.72008 (12)	0.0317 (7)	
O6	1.1993 (3)	1.1322 (3)	0.75792 (11)	0.0293 (7)	
O7	0.9280 (3)	0.6190 (3)	0.58705 (11)	0.0295 (7)	
O8	0.8178 (3)	0.4562 (3)	0.57088 (11)	0.0253 (6)	
O9	0.7827 (3)	0.7584 (3)	0.53809 (12)	0.0310 (7)	

H9	0.7944	0.7771	0.5125	0.047*	
O10	0.6037 (3)	0.6797 (2)	0.59740 (9)	0.0195 (5)	
H10	0.6630	0.7235	0.6023	0.029*	
O11	0.4479 (3)	0.7476 (3)	0.53234 (10)	0.0320 (7)	
O12	0.5479 (4)	0.6380 (4)	0.48342 (11)	0.0463 (10)	
O13	0.2972 (3)	0.6471 (3)	0.62762 (11)	0.0276 (7)	
O14	0.3822 (3)	0.5056 (3)	0.58715 (11)	0.0327 (8)	
O15	0.2955 (3)	0.3293 (3)	0.63798 (9)	0.0199 (5)	
H15	0.3205	0.2918	0.6592	0.030*	
O16	0.0760 (3)	0.4462 (2)	0.61037 (9)	0.0213 (6)	
H16	0.0493	0.5109	0.6021	0.032*	
O17	-0.0251 (2)	0.3105 (3)	0.67293 (9)	0.0217 (5)	
O18	0.1216 (3)	0.3518 (3)	0.72046 (10)	0.0273 (7)	
O1W	0.6901 (17)	0.9073 (17)	0.4779 (6)	0.145 (7)	0.50
H1W1	0.7356	0.9133	0.4565	0.218*	0.50
H1W2	0.6878	0.9711	0.4910	0.218*	0.50
C1	1.0921 (4)	0.8279 (3)	0.66460 (12)	0.0178 (7)	
C2	1.1821 (3)	0.8874 (3)	0.69695 (12)	0.0160 (7)	
H2C	1.2666	0.8599	0.6909	0.019*	
C3	1.1755 (4)	1.0224 (4)	0.69036 (15)	0.0257 (9)	
H3C	1.2103	1.0423	0.6617	0.031*	
C4	1.2533 (4)	1.0834 (3)	0.72602 (13)	0.0220 (8)	
C5	0.8399 (4)	0.5650 (4)	0.56782 (13)	0.0168 (7)	
C6	0.7517 (4)	0.6364 (3)	0.53935 (12)	0.0174 (7)	
H6C	0.7558	0.6047	0.5095	0.021*	
C7	0.6197 (3)	0.6243 (3)	0.55551 (12)	0.0160 (7)	
H7C	0.6016	0.5390	0.5588	0.019*	
C8	0.5297 (4)	0.6755 (4)	0.52122 (13)	0.0234 (8)	
C9	0.3210 (4)	0.5408 (4)	0.61868 (13)	0.0191 (8)	
C10	0.2672 (3)	0.4479 (3)	0.65126 (12)	0.0155 (7)	
H10C	0.3020	0.4622	0.6804	0.019*	
C11	0.1258 (4)	0.4612 (3)	0.65322 (13)	0.0168 (7)	
H11C	0.1054	0.5415	0.6638	0.020*	
C12	0.0704 (4)	0.3690 (3)	0.68453 (12)	0.0175 (7)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.01892 (17)	0.01774 (17)	0.01540 (16)	0.000	-0.00203 (12)	0.000
Cd2	0.02609 (15)	0.01939 (14)	0.01935 (13)	-0.00727 (12)	0.00123 (10)	-0.00092 (10)
Cd3	0.02465 (14)	0.01520 (12)	0.01611 (12)	-0.00012 (11)	-0.00171 (9)	0.00103 (11)
Cd4	0.0214 (2)	0.0270 (2)	0.0417 (2)	0.000	0.00595 (16)	0.000
O1	0.0222 (14)	0.0262 (14)	0.0216 (13)	-0.0078 (12)	-0.0076 (11)	0.0021 (11)
O2	0.0414 (19)	0.0331 (17)	0.0163 (13)	-0.0116 (15)	-0.0006 (12)	-0.0019 (12)
O3	0.0218 (14)	0.0251 (14)	0.0156 (13)	-0.0048 (11)	-0.0076 (10)	0.0027 (11)
O4	0.043 (2)	0.0263 (16)	0.075 (3)	0.0144 (16)	-0.039 (2)	-0.0157 (17)
O5	0.0260 (16)	0.0282 (16)	0.0409 (19)	-0.0114 (13)	0.0012 (14)	-0.0064 (15)
O6	0.0278 (16)	0.0292 (15)	0.0308 (17)	-0.0002 (13)	-0.0033 (12)	-0.0115 (13)

supplementary materials

O7	0.0274 (17)	0.0166 (12)	0.0447 (19)	0.0003 (12)	-0.0195 (14)	-0.0032 (13)
O8	0.0229 (14)	0.0149 (13)	0.0383 (18)	0.0001 (12)	-0.0082 (12)	0.0011 (12)
O9	0.0288 (18)	0.0210 (14)	0.0432 (19)	-0.0068 (12)	-0.0084 (14)	0.0166 (13)
O10	0.0226 (13)	0.0196 (12)	0.0162 (11)	0.0005 (11)	-0.0020 (10)	-0.0045 (11)
O11	0.0330 (17)	0.0416 (18)	0.0213 (14)	0.0254 (15)	-0.0083 (13)	-0.0069 (13)
O12	0.047 (2)	0.069 (3)	0.0224 (15)	0.038 (2)	-0.0175 (15)	-0.0207 (16)
O13	0.0279 (16)	0.0182 (14)	0.0366 (17)	-0.0058 (12)	0.0073 (13)	0.0002 (13)
O14	0.042 (2)	0.0219 (14)	0.0345 (17)	-0.0030 (14)	0.0219 (15)	0.0005 (13)
O15	0.0222 (14)	0.0177 (13)	0.0200 (13)	0.0019 (11)	-0.0014 (10)	0.0040 (11)
O16	0.0235 (15)	0.0197 (13)	0.0206 (13)	-0.0014 (11)	-0.0066 (11)	0.0055 (10)
O17	0.0200 (14)	0.0240 (14)	0.0211 (12)	-0.0063 (12)	0.0015 (10)	0.0025 (12)
O18	0.0236 (16)	0.0360 (17)	0.0223 (14)	-0.0060 (13)	-0.0009 (11)	0.0037 (13)
O1W	0.130 (10)	0.170 (10)	0.136 (10)	-0.007 (8)	0.010 (8)	0.065 (8)
C1	0.0201 (17)	0.0161 (16)	0.0172 (16)	-0.0026 (15)	-0.0031 (13)	-0.0002 (14)
C2	0.0157 (16)	0.0166 (17)	0.0156 (17)	-0.0012 (14)	0.0003 (13)	-0.0013 (14)
C3	0.036 (2)	0.0153 (18)	0.026 (2)	-0.0032 (16)	-0.0151 (17)	0.0011 (16)
C4	0.028 (2)	0.0134 (16)	0.0244 (19)	-0.0045 (16)	-0.0079 (16)	0.0042 (15)
C5	0.0158 (18)	0.0179 (17)	0.0167 (17)	0.0023 (14)	-0.0004 (12)	-0.0023 (14)
C6	0.0195 (18)	0.0172 (17)	0.0156 (16)	0.0016 (15)	-0.0008 (13)	0.0016 (13)
C7	0.0149 (18)	0.0166 (16)	0.0165 (17)	0.0009 (14)	-0.0041 (13)	-0.0034 (14)
C8	0.023 (2)	0.027 (2)	0.0199 (17)	0.0100 (17)	-0.0087 (14)	-0.0056 (16)
C9	0.0132 (17)	0.0214 (19)	0.0228 (19)	-0.0043 (15)	-0.0052 (14)	-0.0020 (16)
C10	0.0140 (17)	0.0171 (16)	0.0154 (16)	-0.0027 (14)	-0.0022 (13)	0.0002 (13)
C11	0.0162 (18)	0.0117 (15)	0.0224 (19)	-0.0016 (14)	-0.0006 (13)	-0.0008 (14)
C12	0.0182 (19)	0.0187 (16)	0.0157 (16)	-0.0001 (14)	0.0046 (13)	-0.0041 (14)

Geometric parameters (Å, °)

Cd1—O5 ⁱ	2.207 (3)	O8—Cd3 ^{xi}	2.247 (3)
Cd1—O5 ⁱⁱ	2.207 (3)	O9—C6	1.408 (5)
Cd1—O1	2.282 (3)	O9—H9	0.82
Cd1—O1 ⁱⁱⁱ	2.282 (3)	O10—C7	1.434 (4)
Cd1—O3	2.388 (3)	O10—H10	0.82
Cd1—O3 ⁱⁱⁱ	2.388 (3)	O11—C8	1.245 (5)
Cd2—O12 ^{iv}	2.196 (3)	O12—C8	1.245 (5)
Cd2—O2	2.220 (3)	O12—Cd2 ^{xii}	2.196 (3)
Cd2—O7	2.260 (3)	O13—C9	1.248 (5)
Cd2—O14 ^v	2.264 (3)	O14—C9	1.233 (5)
Cd2—O9	2.330 (3)	O14—Cd2 ⁱ	2.264 (3)
Cd2—O15 ^v	2.512 (3)	O15—C10	1.422 (5)
Cd3—O13	2.232 (3)	O15—Cd2 ⁱ	2.512 (3)
Cd3—O11	2.238 (3)	O15—H15	0.82
Cd3—O8 ^{vi}	2.247 (3)	O16—C11	1.426 (4)
Cd3—O17 ^v	2.283 (3)	O16—Cd3 ⁱ	2.449 (3)
Cd3—O16 ^v	2.449 (3)	O16—H16	0.82
Cd3—O10	2.470 (3)	O17—C12	1.272 (5)

Cd4—O6 ^{vii}	2.290 (3)	O17—Cd3 ⁱ	2.283 (3)
Cd4—O6 ^{viii}	2.290 (3)	O18—C12	1.245 (5)
Cd4—O18	2.338 (3)	O1W—H1W1	0.82
Cd4—O18 ^{ix}	2.338 (3)	O1W—H1W2	0.82
Cd4—O4 ^{viii}	2.424 (4)	C1—C2	1.538 (5)
Cd4—O4 ^{vii}	2.424 (4)	C2—C3	1.526 (5)
O1—C1	1.250 (5)	C2—H2C	0.98
O2—C1	1.257 (5)	C3—C4	1.536 (5)
O3—C2	1.428 (5)	C3—H3C	0.98
O3—H3	0.82	C5—C6	1.518 (5)
O4—C3	1.412 (6)	C6—C7	1.514 (5)
O4—Cd4 ^x	2.424 (4)	C6—H6C	0.98
O4—H4	0.82	C7—C8	1.540 (5)
O5—C4	1.233 (6)	C7—H7C	0.98
O5—Cd1 ^v	2.207 (3)	C9—C10	1.554 (5)
O6—C4	1.261 (5)	C10—C11	1.534 (5)
O6—Cd4 ^x	2.290 (3)	C10—H10C	0.98
O7—C5	1.271 (5)	C11—C12	1.530 (5)
O8—C5	1.246 (5)	C11—H11C	0.98
O5 ⁱ —Cd1—O5 ⁱⁱ	101.55 (19)	C6—O9—H9	108.1
O5 ⁱ —Cd1—O1	79.43 (12)	Cd2—O9—H9	108.1
O5 ⁱⁱ —Cd1—O1	123.99 (12)	C7—O10—Cd3	112.6 (2)
O5 ⁱ —Cd1—O1 ⁱⁱⁱ	123.99 (12)	C7—O10—H10	109.1
O5 ⁱⁱ —Cd1—O1 ⁱⁱⁱ	79.43 (12)	Cd3—O10—H10	109.1
O1—Cd1—O1 ⁱⁱⁱ	145.46 (15)	C8—O11—Cd3	123.4 (3)
O5 ⁱ —Cd1—O3	148.58 (11)	C8—O12—Cd2 ^{xii}	130.7 (3)
O5 ⁱⁱ —Cd1—O3	93.84 (12)	C9—O13—Cd3	122.1 (3)
O1—Cd1—O3	69.28 (10)	C9—O14—Cd2 ⁱ	125.2 (3)
O1 ⁱⁱⁱ —Cd1—O3	85.51 (10)	C10—O15—Cd2 ⁱ	113.8 (2)
O5 ⁱ —Cd1—O3 ⁱⁱⁱ	93.84 (12)	C10—O15—H15	108.8
O5 ⁱⁱ —Cd1—O3 ⁱⁱⁱ	148.58 (11)	Cd2 ⁱ —O15—H15	108.8
O1—Cd1—O3 ⁱⁱⁱ	85.51 (10)	C11—O16—Cd3 ⁱ	116.8 (2)
O1 ⁱⁱⁱ —Cd1—O3 ⁱⁱⁱ	69.28 (9)	C11—O16—H16	108.1
O3—Cd1—O3 ⁱⁱⁱ	86.61 (15)	Cd3 ⁱ —O16—H16	108.1
O12 ^{iv} —Cd2—O2	100.04 (13)	C12—O17—Cd3 ⁱ	122.8 (2)
O12 ^{iv} —Cd2—O7	112.50 (14)	C12—O18—Cd4	101.8 (3)
O2—Cd2—O7	101.90 (11)	H1W1—O1W—H1W2	109.7
O12 ^{iv} —Cd2—O14 ^v	92.86 (14)	O1—C1—O2	125.1 (4)
O2—Cd2—O14 ^v	90.47 (13)	O1—C1—C2	119.5 (3)
O7—Cd2—O14 ^v	148.89 (13)	O2—C1—C2	115.4 (3)
O12 ^{iv} —Cd2—O9	88.41 (13)	O3—C2—C3	110.6 (3)
O2—Cd2—O9	170.78 (11)	O3—C2—C1	110.1 (3)
O7—Cd2—O9	71.16 (10)	C3—C2—C1	108.3 (3)

supplementary materials

O14 ^v —Cd2—O9	92.81 (13)	O3—C2—H2C	109.3
O12 ^{iv} —Cd2—O15 ^v	157.71 (14)	C3—C2—H2C	109.3
O2—Cd2—O15 ^v	91.88 (10)	C1—C2—H2C	109.3
O7—Cd2—O15 ^v	82.91 (12)	O4—C3—C2	109.7 (4)
O14 ^v —Cd2—O15 ^v	68.10 (10)	O4—C3—C4	111.3 (4)
O9—Cd2—O15 ^v	81.37 (11)	C2—C3—C4	108.8 (3)
O13—Cd3—O11	104.01 (13)	O4—C3—H3C	109.0
O13—Cd3—O8 ^{vi}	119.99 (12)	C2—C3—H3C	109.0
O11—Cd3—O8 ^{vi}	82.89 (12)	C4—C3—H3C	109.0
O13—Cd3—O17 ^v	82.97 (11)	O5—C4—O6	126.6 (4)
O11—Cd3—O17 ^v	150.04 (10)	O5—C4—C3	114.1 (4)
O8 ^{vi} —Cd3—O17 ^v	119.27 (11)	O6—C4—C3	119.4 (4)
O13—Cd3—O16 ^v	151.21 (11)	O8—C5—O7	125.0 (4)
O11—Cd3—O16 ^v	98.13 (12)	O8—C5—C6	116.0 (3)
O8 ^{vi} —Cd3—O16 ^v	80.54 (10)	O7—C5—C6	119.0 (4)
O17 ^v —Cd3—O16 ^v	68.79 (10)	O9—C6—C7	108.6 (3)
O13—Cd3—O10	94.40 (11)	O9—C6—C5	112.2 (3)
O11—Cd3—O10	70.20 (10)	C7—C6—C5	110.8 (3)
O8 ^{vi} —Cd3—O10	140.87 (11)	O9—C6—H6C	108.4
O17 ^v —Cd3—O10	80.31 (9)	C7—C6—H6C	108.4
O16 ^v —Cd3—O10	75.93 (9)	C5—C6—H6C	108.4
O6 ^{vii} —Cd4—O6 ^{viii}	141.86 (17)	O10—C7—C6	111.5 (3)
O6 ^{vii} —Cd4—O18	136.50 (11)	O10—C7—C8	111.8 (3)
O6 ^{viii} —Cd4—O18	75.69 (12)	C6—C7—C8	109.7 (3)
O6 ^{vii} —Cd4—O18 ^{ix}	75.69 (12)	O10—C7—H7C	107.9
O6 ^{viii} —Cd4—O18 ^{ix}	136.50 (11)	C6—C7—H7C	107.9
O18—Cd4—O18 ^{ix}	85.92 (17)	C8—C7—H7C	107.9
O6 ^{vii} —Cd4—O4 ^{viii}	85.73 (13)	O11—C8—O12	125.8 (4)
O6 ^{viii} —Cd4—O4 ^{viii}	69.85 (11)	O11—C8—C7	120.2 (3)
O18—Cd4—O4 ^{viii}	93.37 (14)	O12—C8—C7	114.1 (4)
O18 ^{ix} —Cd4—O4 ^{viii}	151.60 (11)	O14—C9—O13	125.9 (4)
O6 ^{vii} —Cd4—O4 ^{vii}	69.85 (11)	O14—C9—C10	119.2 (4)
O6 ^{viii} —Cd4—O4 ^{vii}	85.73 (13)	O13—C9—C10	114.9 (4)
O18—Cd4—O4 ^{vii}	151.60 (11)	O15—C10—C11	108.4 (3)
O18 ^{ix} —Cd4—O4 ^{vii}	93.37 (14)	O15—C10—C9	111.2 (3)
O4 ^{viii} —Cd4—O4 ^{vii}	100.2 (2)	C11—C10—C9	109.4 (3)
C1—O1—Cd1	122.3 (2)	O15—C10—H10C	109.3
C1—O2—Cd2	118.2 (3)	C11—C10—H10C	109.3
C2—O3—Cd1	116.9 (2)	C9—C10—H10C	109.3
C2—O3—H3	108.1	O16—C11—C12	110.4 (3)
Cd1—O3—H3	108.1	O16—C11—C10	109.1 (3)
C3—O4—Cd4 ^x	115.6 (3)	C12—C11—C10	110.3 (3)

C3—O4—H4	108.4	O16—C11—H11C	109.0
Cd4 ^x —O4—H4	108.4	C12—C11—H11C	109.0
C4—O5—Cd1 ^v	123.2 (3)	C10—C11—H11C	109.0
C4—O6—Cd4 ^x	119.6 (3)	O18—C12—O17	121.7 (4)
C5—O7—Cd2	120.8 (3)	O18—C12—C11	118.9 (3)
C5—O8—Cd3 ^{xi}	135.9 (3)	O17—C12—C11	119.3 (3)
C6—O9—Cd2	116.9 (2)		
O5 ⁱ —Cd1—O1—C1	-165.4 (3)	C1—C2—C3—C4	173.4 (3)
O5 ⁱⁱ —Cd1—O1—C1	-68.6 (3)	Cd1 ^v —O5—C4—O6	-23.0 (6)
O1 ⁱⁱⁱ —Cd1—O1—C1	57.5 (3)	Cd1 ^v —O5—C4—C3	158.2 (3)
O3—Cd1—O1—C1	11.7 (3)	Cd4 ^x —O6—C4—O5	155.8 (3)
O3 ⁱⁱⁱ —Cd1—O1—C1	99.8 (3)	Cd4 ^x —O6—C4—C3	-25.5 (5)
O12 ^{iv} —Cd2—O2—C1	-173.2 (3)	O4—C3—C4—O5	-163.6 (4)
O7—Cd2—O2—C1	-57.5 (3)	C2—C3—C4—O5	75.5 (5)
O14 ^v —Cd2—O2—C1	93.8 (3)	O4—C3—C4—O6	17.6 (5)
O15 ^v —Cd2—O2—C1	25.7 (3)	C2—C3—C4—O6	-103.4 (4)
O5 ⁱ —Cd1—O3—C2	-6.4 (4)	Cd3 ^{xi} —O8—C5—O7	-6.2 (7)
O5 ⁱⁱ —Cd1—O3—C2	113.2 (3)	Cd3 ^{xi} —O8—C5—C6	175.0 (3)
O1—Cd1—O3—C2	-11.8 (2)	Cd2—O7—C5—O8	-179.7 (3)
O1 ⁱⁱⁱ —Cd1—O3—C2	-167.7 (3)	Cd2—O7—C5—C6	-0.9 (5)
O3 ⁱⁱⁱ —Cd1—O3—C2	-98.3 (3)	Cd2—O9—C6—C7	123.0 (3)
O12 ^{iv} —Cd2—O7—C5	-79.3 (4)	Cd2—O9—C6—C5	0.1 (4)
O2—Cd2—O7—C5	174.4 (3)	O8—C5—C6—O9	179.4 (4)
O14 ^v —Cd2—O7—C5	63.0 (4)	O7—C5—C6—O9	0.5 (5)
O9—Cd2—O7—C5	0.7 (3)	O8—C5—C6—C7	57.8 (5)
O15 ^v —Cd2—O7—C5	83.9 (3)	O7—C5—C6—C7	-121.1 (4)
O12 ^{iv} —Cd2—O9—C6	114.1 (3)	Cd3—O10—C7—C6	136.4 (2)
O7—Cd2—O9—C6	-0.4 (3)	Cd3—O10—C7—C8	13.1 (4)
O14 ^v —Cd2—O9—C6	-153.1 (3)	O9—C6—C7—O10	-56.7 (4)
O15 ^v —Cd2—O9—C6	-85.8 (3)	C5—C6—C7—O10	67.0 (4)
O13—Cd3—O10—C7	91.5 (2)	O9—C6—C7—C8	67.8 (4)
O11—Cd3—O10—C7	-11.8 (2)	C5—C6—C7—C8	-168.5 (3)
O8 ^{vi} —Cd3—O10—C7	-61.3 (3)	Cd3—O11—C8—O12	174.4 (4)
O17 ^v —Cd3—O10—C7	173.6 (2)	Cd3—O11—C8—C7	-5.6 (6)
O16 ^v —Cd3—O10—C7	-116.0 (2)	Cd2 ^{xii} —O12—C8—O11	2.3 (9)
O13—Cd3—O11—C8	-80.2 (4)	Cd2 ^{xii} —O12—C8—C7	-177.7 (3)
O8 ^{vi} —Cd3—O11—C8	160.6 (4)	O10—C7—C8—O11	-6.3 (6)
O17 ^v —Cd3—O11—C8	20.2 (5)	C6—C7—C8—O11	-130.6 (4)
O16 ^v —Cd3—O11—C8	81.3 (4)	O10—C7—C8—O12	173.7 (4)
O10—Cd3—O11—C8	9.5 (4)	C6—C7—C8—O12	49.5 (5)
O11—Cd3—O13—C9	41.6 (4)	Cd2 ⁱ —O14—C9—O13	167.6 (3)
O8 ^{vi} —Cd3—O13—C9	131.4 (3)	Cd2 ⁱ —O14—C9—C10	-12.1 (5)

supplementary materials

O17 ^v —Cd3—O13—C9	-108.7 (3)	Cd3—O13—C9—O14	-21.2 (6)
O16 ^v —Cd3—O13—C9	-97.5 (4)	Cd3—O13—C9—C10	158.5 (2)
O10—Cd3—O13—C9	-29.0 (3)	Cd2 ⁱ —O15—C10—C11	132.7 (2)
O6 ^{vii} —Cd4—O18—C12	-25.4 (3)	Cd2 ⁱ —O15—C10—C9	12.4 (3)
O6 ^{viii} —Cd4—O18—C12	130.2 (3)	O14—C9—C10—O15	-1.9 (5)
O18 ^{ix} —Cd4—O18—C12	-89.6 (3)	O13—C9—C10—O15	178.3 (3)
O4 ^{viii} —Cd4—O18—C12	61.9 (3)	O14—C9—C10—C11	-121.7 (4)
O4 ^{vii} —Cd4—O18—C12	-179.1 (3)	O13—C9—C10—C11	58.6 (4)
Cd1—O1—C1—O2	172.9 (3)	Cd3 ⁱ —O16—C11—C12	5.5 (4)
Cd1—O1—C1—C2	-10.0 (5)	Cd3 ⁱ —O16—C11—C10	126.9 (3)
Cd2—O2—C1—O1	14.3 (6)	O15—C10—C11—O16	-63.9 (4)
Cd2—O2—C1—C2	-162.9 (2)	C9—C10—C11—O16	57.6 (4)
Cd1—O3—C2—C3	130.8 (3)	O15—C10—C11—C12	57.6 (4)
Cd1—O3—C2—C1	11.1 (4)	C9—C10—C11—C12	179.0 (3)
O1—C1—C2—O3	-1.4 (5)	Cd4—O18—C12—O17	1.1 (4)
O2—C1—C2—O3	176.0 (3)	Cd4—O18—C12—C11	-177.6 (3)
O1—C1—C2—C3	-122.5 (4)	Cd3 ⁱ —O17—C12—O18	-161.8 (3)
O2—C1—C2—C3	54.9 (5)	Cd3 ⁱ —O17—C12—C11	16.8 (5)
Cd4 ^x —O4—C3—C2	118.3 (3)	O16—C11—C12—O18	164.7 (3)
Cd4 ^x —O4—C3—C4	-2.1 (5)	C10—C11—C12—O18	44.1 (5)
O3—C2—C3—O4	-69.3 (4)	O16—C11—C12—O17	-14.0 (5)
C1—C2—C3—O4	51.5 (5)	C10—C11—C12—O17	-134.7 (4)
O3—C2—C3—C4	52.7 (5)		

Symmetry codes: (i) $x-1/2, y-1/2, z$; (ii) $-x+5/2, y-1/2, -z+3/2$; (iii) $-x+2, y, -z+3/2$; (iv) $x+1/2, -y+3/2, -z+1$; (v) $x+1/2, y+1/2, z$; (vi) $x-1/2, y+1/2, z$; (vii) $-x+1, y-1, -z+3/2$; (viii) $x-1, y-1, z$; (ix) $-x, y, -z+3/2$; (x) $x+1, y+1, z$; (xi) $x+1/2, y-1/2, z$; (xii) $x-1/2, -y+3/2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 ⁱⁱⁱ —O18 ^{xiii}	0.82	1.96	2.740 (4)	159
O4—H4 ^v —O10 ^v	0.82	2.50	3.236 (6)	149
O9—H9 ^{iv} —O11 ^{iv}	0.82	2.17	2.797 (5)	134
O9—H9 ^w —O1w	0.82	2.12	2.68 (2)	125
O10—H10 ^v —O15 ^v	0.82	2.15	2.938 (4)	160
O15—H15 ⁱ —O1 ⁱ	0.82	2.13	2.717 (4)	128
O16—H16 ^{xiv} —O7 ^{xiv}	0.82	1.84	2.609 (4)	155
O1W—H1W1 ^{iv} —O14 ^{iv}	0.82	2.26	3.034 (19)	157

Symmetry codes: (xiii) $-x+3/2, y+1/2, -z+3/2$; (v) $x+1/2, y+1/2, z$; (iv) $x+1/2, -y+3/2, -z+1$; (i) $x-1/2, y-1/2, z$; (xiv) $x-1, y, z$.

