

[(1*R*,3*S*)-3-(1,3-Dithian-2-yl)-2,2-dimethylcyclopropyl]diphenylmethanol

Risong Na and Min Wang*

Department of Applied Chemistry, China Agricultural University, 100193 Beijing, People's Republic of China

Correspondence e-mail: wangmincau@yahoo.com.cn


Received 30 March 2009; accepted 29 April 2009

 Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.038; wR factor = 0.084; data-to-parameter ratio = 18.8.

In the title compound, $\text{C}_{22}\text{H}_{26}\text{OS}_2$, prepared from (–)-1*R*-cis-caronaldehyde, the 1,3-dithiane ring adopts a chair conformation. An intramolecular $\text{O}-\text{H}\cdots\text{S}$ hydrogen bond influences the molecular conformation. In the crystal, weak intermolecular $\text{C}-\text{H}\cdots\text{S}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains propagating along [010].

Related literature

For the details of preparation of the analogous compound, (1*R*,3*S*)-methyl-3-(1,3-dithian-2-yl)-2,2-dimethylcyclopropane carboxylate, see: Mazzanti *et al.* (1997); Veyrat *et al.* (1997); Perollier *et al.* (1997).



Experimental

Crystal data

 $\text{C}_{22}\text{H}_{26}\text{OS}_2$
 $M_r = 370.55$

 Monoclinic, $P2_1$
 $a = 9.5578$ (19) Å

 $b = 11.199$ (2) Å

 $c = 9.6512$ (19) Å

 $\beta = 101.14$ (3)°

 $V = 1013.6$ (4) Å³
 $Z = 2$

 Mo $K\alpha$ radiation

 $\mu = 0.27$ mm⁻¹
 $T = 123$ K

 $0.40 \times 0.40 \times 0.30$ mm

Data collection

Rigaku R-AXIS RAPID IP

diffractometer

Absorption correction: multi-scan

(ABSCOR; Higashi, 1995)

 $T_{\min} = 0.900$, $T_{\max} = 0.924$

4303 measured reflections

4303 independent reflections

 2920 reflections with $I > 2\sigma(I)$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.084$
 $S = 0.84$

4303 reflections

229 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.30$ e Å⁻³

Absolute structure: Flack (1983),

1878 Friedel pairs

Flack parameter: 0.05 (7)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H2}\cdots\text{S1}$	0.84	2.58	3.330 (2)	149
$\text{C7}-\text{H7A}\cdots\text{S2}^{\text{i}}$	0.99	2.89	3.736 (3)	144
$\text{C9}-\text{H9A}\cdots\text{O1}^{\text{ii}}$	0.99	2.60	3.236 (3)	122

 Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 2000); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku, 2000); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Financial support from the National Natural Science Foundation of China (grant Nos. 20472111 and 20742004) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2542).

References

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Mazzanti, M., Marchon, J. C., Shang, M. Y., Scheidt, W. R., Jia, S. L. & Shelnutt, J. A. (1997). *J. Am. Chem. Soc.* **119**, 12400–12401.
 Perollier, C., Pecaut, J., Ramasseul, R. & Marchon, J. C. (1997). *Bull. Soc. Chim. Fr.* **134**, 517–523.
 Rigaku (2000). *RAPID-AUTO* and *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Veyrat, M., Fantin, L., Desmoulin, S., Petitjean, A., Mazzanti, M., Ramasseul, R., Marchon, J. C. & Bau, R. (1997). *Bull. Soc. Chim. Fr.* **134**, 703–711.

supplementary materials

Acta Cryst. (2009). E65, o1230 [doi:10.1107/S1600536809016079]

[(1*R*,3*S*)-3-(1,3-Dithian-2-yl)-2,2-dimethylcyclopropyl]diphenylmethanol

R. Na and M. Wang

Comment

We designed and prepared a novel type of chiral dithiane alcohol based with chiral *cis*-cyclopropane from (-)-1*R*-*cis*-Caronaldehyde. Details of preparation of the analogue compound were discussed in the literature (Mazzanti *et al.*, 1997; Veyrat *et al.*, 1997; Perollier *et al.*, 1997).

In this paper, the crystal structure of the title compound, (I), is reported. In (I) (Fig. 1), the 1,3-dithiane ring adopts a chair conformation. Intramolecular O—H \cdots S hydrogen bond (Table 1) influences the molecular conformation. In the crystal, weak intermolecular C—H \cdots O and C—H \cdots S hydrogen bonds (Table 1) link the molecules into chains propagated in direction [010].

Experimental

Magnesium (0.4 g, 15.6 mmol) was added to 15 ml of anhydrous THF. A solution of bromobenzene (2.0 g, 12.5 mmol in 5 ml of THF) was added dropwise into the above mixture. Once the reaction began, the rest of the bromobenzene solution was added at a rate that maintained a gentle reflux. When the addition of the bromobenzene solution was complete, the mixture was refluxed for 20 min, and was then cooled to 273 K. (1*R*,3*S*)-Methyl-3-(1,3-dithian-2-yl)-2,2-dimethylcyclopropane carboxylate (5 mmol) (Mazzanti *et al.*, 1997; Veyrat *et al.*, 1997; Perollier *et al.*, 1997) was dissolved in 5 ml of anhydrous THF and added to the prepared Grignard mixture. After the solution of carboxylate had been added, the resulting mixture was stirred at room temperature for an additional 24 h. The reaction was quenched with saturated NH₄Cl (aq), and the mixture was extracted several times with Et₂O. The organic phases were combined, dried over MgSO₄ and concentrated under reduced pressure. The residual yellow solid was purified by recrystallization in Et₂O to yield compound (I) as colourless crystals. Colourless solid, m.p. 427 K; [α]₂₀^D = +19.93 (*c* 0.03, CHCl₃).

Refinement

All H atoms were positioned geometrically and treated as riding on their parent atoms with C—H = 0.95 (C_{aromatic}), 0.98 (C_{methyl}), 0.99 (CH₂) and 1.00 (CH) Å and O—H = 0.84 Å, and with U_{iso} (H) = 1.5 U_{eq} (C_{methyl}, O) and 1.2 U_{eq} (other C).

Figures




Fig. 1. The molecular structure of (I) showing the atomic numbering and 30% probability displacement ellipsoids for non-H atoms.

[(1*R*,3*S*)-3-(1,3-Dithian-2-yl)-2,2-dimethylcyclopropyl]diphenylmethanol

Crystal data

$C_{22}H_{26}OS_2$	$F_{000} = 396$
$M_r = 370.55$	$D_x = 1.214 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
Hall symbol: P 2yb	$\lambda = 0.71073 \text{ \AA}$
$a = 9.5578 (19) \text{ \AA}$	Cell parameters from 7560 reflections
$b = 11.199 (2) \text{ \AA}$	$\theta = 2.2\text{--}27.5^\circ$
$c = 9.6512 (19) \text{ \AA}$	$\mu = 0.27 \text{ mm}^{-1}$
$\beta = 101.14 (3)^\circ$	$T = 123 \text{ K}$
$V = 1013.6 (4) \text{ \AA}^3$	Block, colourless
$Z = 2$	$0.40 \times 0.40 \times 0.30 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID IP diffractometer	4303 independent reflections
Radiation source: fine-focus sealed tube	2920 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.0000$
Detector resolution: $10.00 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.5^\circ$
$T = 123 \text{ K}$	$\theta_{\text{min}} = 2.2^\circ$
ω scans	$h = -12 \rightarrow 12$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$k = -14 \rightarrow 14$
$T_{\text{min}} = 0.900$, $T_{\text{max}} = 0.924$	$l = -12 \rightarrow 12$
4303 measured reflections	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.038$	$w = 1/[\sigma^2(F_o^2) + (0.0379P)^2]$
$wR(F^2) = 0.084$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.84$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4303 reflections	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
229 parameters	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.046 (2)
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1878 Friedel pairs
	Flack parameter: 0.05 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6857 (3)	0.2287 (2)	0.2178 (3)	0.0254 (6)
H1	0.6517	0.2825	0.1355	0.030*
C2	0.6466 (3)	0.2775 (2)	0.3516 (3)	0.0287 (7)
C3	0.8003 (3)	0.2893 (2)	0.3280 (3)	0.0238 (6)
H3	0.8246	0.3717	0.3005	0.029*
C4	0.5595 (3)	0.3923 (3)	0.3379 (3)	0.0462 (9)
H4A	0.4577	0.3727	0.3176	0.069*
H4B	0.5827	0.4407	0.2609	0.069*
H4C	0.5822	0.4373	0.4265	0.069*
C5	0.6172 (3)	0.1950 (3)	0.4668 (3)	0.0369 (8)
H5A	0.5156	0.1745	0.4492	0.055*
H5B	0.6432	0.2350	0.5585	0.055*
H5C	0.6739	0.1220	0.4674	0.055*
C6	0.9233 (2)	0.2264 (2)	0.4217 (3)	0.0236 (6)
H6	0.8867	0.1549	0.4656	0.028*
C7	1.1962 (3)	0.1241 (3)	0.4492 (3)	0.0351 (7)
H7A	1.1632	0.0529	0.4944	0.042*
H7B	1.2751	0.0987	0.4028	0.042*
C8	1.2536 (3)	0.2153 (2)	0.5636 (3)	0.0354 (8)
H8A	1.3404	0.1827	0.6245	0.042*
H8B	1.2811	0.2887	0.5184	0.042*
C9	1.1470 (3)	0.2472 (3)	0.6542 (3)	0.0354 (7)
H9A	1.1958	0.2940	0.7366	0.042*
H9B	1.1109	0.1729	0.6902	0.042*
C10	0.6776 (3)	0.0972 (2)	0.1726 (3)	0.0235 (6)
C11	0.7437 (3)	0.0754 (2)	0.0414 (3)	0.0258 (6)
C12	0.7434 (3)	-0.0429 (3)	-0.0102 (3)	0.0389 (8)
H12	0.6995	-0.1044	0.0341	0.047*
C13	0.8053 (3)	-0.0705 (3)	-0.1232 (3)	0.0479 (9)
H13	0.8059	-0.1507	-0.1551	0.057*
C14	0.8676 (3)	0.0196 (4)	-0.1912 (3)	0.0522 (10)
H14	0.9091	0.0014	-0.2705	0.063*
C15	0.8679 (4)	0.1346 (3)	-0.1422 (4)	0.0537 (10)

supplementary materials

H15	0.9100	0.1958	-0.1885	0.064*
C16	0.8079 (3)	0.1635 (3)	-0.0258 (3)	0.0404 (8)
H16	0.8108	0.2435	0.0074	0.048*
C17	0.5212 (3)	0.0589 (2)	0.1427 (3)	0.0259 (6)
C18	0.4713 (3)	-0.0319 (3)	0.2171 (3)	0.0345 (7)
H18	0.5356	-0.0737	0.2880	0.041*
C19	0.3262 (3)	-0.0628 (3)	0.1886 (3)	0.0412 (8)
H19	0.2933	-0.1268	0.2385	0.049*
C20	0.2309 (3)	-0.0007 (3)	0.0882 (3)	0.0392 (8)
H20	0.1323	-0.0203	0.0713	0.047*
C21	0.2790 (3)	0.0895 (3)	0.0129 (3)	0.0410 (8)
H21	0.2141	0.1313	-0.0575	0.049*
C22	0.4234 (3)	0.1196 (3)	0.0402 (3)	0.0370 (8)
H22	0.4559	0.1825	-0.0117	0.044*
O1	0.75071 (18)	0.02068 (15)	0.28301 (18)	0.0294 (5)
H2	0.8381	0.0324	0.2863	0.035*
S1	1.05026 (7)	0.17887 (6)	0.31479 (7)	0.02959 (18)
S2	0.99674 (7)	0.33324 (6)	0.55984 (8)	0.0331 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0253 (14)	0.0246 (14)	0.0234 (16)	0.0024 (12)	-0.0025 (12)	0.0004 (11)
C2	0.0194 (14)	0.0317 (15)	0.0309 (17)	0.0023 (12)	-0.0056 (12)	-0.0089 (13)
C3	0.0195 (13)	0.0240 (14)	0.0248 (15)	0.0010 (11)	-0.0030 (11)	-0.0005 (11)
C4	0.0302 (17)	0.0495 (19)	0.053 (2)	0.0138 (15)	-0.0065 (15)	-0.0172 (16)
C5	0.0221 (14)	0.058 (2)	0.0314 (16)	-0.0069 (14)	0.0060 (12)	-0.0125 (16)
C6	0.0179 (13)	0.0270 (14)	0.0247 (15)	0.0028 (11)	0.0008 (11)	-0.0022 (11)
C7	0.0208 (15)	0.0375 (17)	0.045 (2)	-0.0002 (12)	0.0011 (13)	-0.0007 (14)
C8	0.0220 (15)	0.0344 (18)	0.046 (2)	-0.0034 (13)	-0.0026 (14)	0.0076 (14)
C9	0.0328 (16)	0.0335 (16)	0.0338 (18)	-0.0009 (14)	-0.0089 (14)	-0.0005 (13)
C10	0.0230 (14)	0.0218 (14)	0.0230 (15)	0.0046 (12)	-0.0021 (11)	0.0027 (11)
C11	0.0221 (14)	0.0318 (15)	0.0216 (14)	0.0074 (13)	-0.0008 (11)	0.0014 (12)
C12	0.0363 (18)	0.0429 (18)	0.0362 (18)	0.0034 (15)	0.0036 (14)	-0.0071 (15)
C13	0.0389 (19)	0.068 (3)	0.036 (2)	0.0096 (18)	0.0057 (16)	-0.0185 (19)
C14	0.0364 (19)	0.093 (3)	0.0271 (18)	0.021 (2)	0.0066 (15)	-0.013 (2)
C15	0.052 (2)	0.069 (3)	0.047 (2)	0.0117 (18)	0.0268 (18)	0.0166 (18)
C16	0.0431 (18)	0.0439 (19)	0.0360 (18)	0.0079 (17)	0.0121 (14)	0.0093 (16)
C17	0.0306 (15)	0.0253 (15)	0.0207 (14)	0.0019 (12)	0.0027 (12)	-0.0046 (12)
C18	0.0322 (16)	0.0349 (17)	0.0321 (17)	-0.0011 (14)	-0.0044 (13)	0.0055 (13)
C19	0.0405 (19)	0.0399 (18)	0.0406 (19)	-0.0124 (15)	0.0013 (15)	0.0031 (15)
C20	0.0277 (16)	0.050 (2)	0.0367 (19)	-0.0058 (15)	-0.0003 (14)	-0.0071 (15)
C21	0.0308 (17)	0.051 (2)	0.0378 (19)	0.0016 (16)	-0.0032 (14)	0.0019 (15)
C22	0.0334 (17)	0.0415 (18)	0.0338 (18)	-0.0031 (14)	0.0005 (14)	0.0093 (14)
O1	0.0285 (11)	0.0267 (10)	0.0287 (11)	0.0028 (8)	-0.0053 (9)	0.0036 (9)
S1	0.0226 (3)	0.0370 (4)	0.0294 (4)	0.0013 (3)	0.0058 (3)	0.0009 (3)
S2	0.0289 (4)	0.0306 (4)	0.0356 (4)	0.0023 (3)	-0.0043 (3)	-0.0068 (4)

Geometric parameters (Å, °)

C1—C2	1.515 (4)	C9—H9B	0.9900
C1—C3	1.531 (3)	C10—O1	1.439 (3)
C1—C10	1.533 (4)	C10—C17	1.529 (4)
C1—H1	1.0000	C10—C11	1.540 (4)
C2—C5	1.513 (4)	C11—C16	1.387 (4)
C2—C4	1.523 (4)	C11—C12	1.415 (4)
C2—C3	1.536 (4)	C12—C13	1.372 (4)
C3—C6	1.511 (3)	C12—H12	0.9500
C3—H3	1.0000	C13—C14	1.398 (5)
C4—H4A	0.9800	C13—H13	0.9500
C4—H4B	0.9800	C14—C15	1.371 (5)
C4—H4C	0.9800	C14—H14	0.9500
C5—H5A	0.9800	C15—C16	1.395 (4)
C5—H5B	0.9800	C15—H15	0.9500
C5—H5C	0.9800	C16—H16	0.9500
C6—S1	1.818 (3)	C17—C18	1.382 (4)
C6—S2	1.827 (3)	C17—C22	1.399 (4)
C6—H6	1.0000	C18—C19	1.404 (4)
C7—C8	1.526 (4)	C18—H18	0.9500
C7—S1	1.818 (3)	C19—C20	1.382 (4)
C7—H7A	0.9900	C19—H19	0.9500
C7—H7B	0.9900	C20—C21	1.375 (4)
C8—C9	1.508 (4)	C20—H20	0.9500
C8—H8A	0.9900	C21—C22	1.396 (4)
C8—H8B	0.9900	C21—H21	0.9500
C9—S2	1.821 (3)	C22—H22	0.9500
C9—H9A	0.9900	O1—H2	0.8400
C2—C1—C3	60.55 (17)	S2—C9—H9A	109.0
C2—C1—C10	125.6 (2)	C8—C9—H9B	109.0
C3—C1—C10	127.8 (2)	S2—C9—H9B	109.0
C2—C1—H1	111.3	H9A—C9—H9B	107.8
C3—C1—H1	111.3	O1—C10—C17	106.8 (2)
C10—C1—H1	111.3	O1—C10—C1	111.8 (2)
C5—C2—C1	121.2 (2)	C17—C10—C1	108.5 (2)
C5—C2—C4	113.8 (3)	O1—C10—C11	107.24 (19)
C1—C2—C4	116.8 (2)	C17—C10—C11	110.0 (2)
C5—C2—C3	118.7 (2)	C1—C10—C11	112.4 (2)
C1—C2—C3	60.24 (18)	C16—C11—C12	118.2 (3)
C4—C2—C3	115.9 (2)	C16—C11—C10	124.1 (2)
C6—C3—C1	125.3 (2)	C12—C11—C10	117.7 (2)
C6—C3—C2	121.8 (2)	C13—C12—C11	121.2 (3)
C1—C3—C2	59.21 (17)	C13—C12—H12	119.4
C6—C3—H3	113.3	C11—C12—H12	119.4
C1—C3—H3	113.3	C12—C13—C14	119.9 (3)
C2—C3—H3	113.3	C12—C13—H13	120.0
C2—C4—H4A	109.5	C14—C13—H13	120.0

supplementary materials

C2—C4—H4B	109.5	C15—C14—C13	119.2 (3)
H4A—C4—H4B	109.5	C15—C14—H14	120.4
C2—C4—H4C	109.5	C13—C14—H14	120.4
H4A—C4—H4C	109.5	C14—C15—C16	121.5 (3)
H4B—C4—H4C	109.5	C14—C15—H15	119.3
C2—C5—H5A	109.5	C16—C15—H15	119.3
C2—C5—H5B	109.5	C11—C16—C15	120.0 (3)
H5A—C5—H5B	109.5	C11—C16—H16	120.0
C2—C5—H5C	109.5	C15—C16—H16	120.0
H5A—C5—H5C	109.5	C18—C17—C22	118.3 (3)
H5B—C5—H5C	109.5	C18—C17—C10	122.1 (2)
C3—C6—S1	108.90 (19)	C22—C17—C10	119.5 (2)
C3—C6—S2	106.04 (18)	C17—C18—C19	120.4 (3)
S1—C6—S2	113.65 (13)	C17—C18—H18	119.8
C3—C6—H6	109.4	C19—C18—H18	119.8
S1—C6—H6	109.4	C20—C19—C18	120.4 (3)
S2—C6—H6	109.4	C20—C19—H19	119.8
C8—C7—S1	114.18 (19)	C18—C19—H19	119.8
C8—C7—H7A	108.7	C21—C20—C19	119.9 (3)
S1—C7—H7A	108.7	C21—C20—H20	120.1
C8—C7—H7B	108.7	C19—C20—H20	120.1
S1—C7—H7B	108.7	C20—C21—C22	119.8 (3)
H7A—C7—H7B	107.6	C20—C21—H21	120.1
C9—C8—C7	112.9 (2)	C22—C21—H21	120.1
C9—C8—H8A	109.0	C21—C22—C17	121.2 (3)
C7—C8—H8A	109.0	C21—C22—H22	119.4
C9—C8—H8B	109.0	C17—C22—H22	119.4
C7—C8—H8B	109.0	C10—O1—H2	105.8
H8A—C8—H8B	107.8	C7—S1—C6	101.43 (13)
C8—C9—S2	113.0 (2)	C9—S2—C6	99.98 (13)
C8—C9—H9A	109.0		
C3—C1—C2—C5	-107.5 (3)	C1—C10—C11—C12	179.7 (2)
C10—C1—C2—C5	10.0 (4)	C16—C11—C12—C13	0.4 (4)
C3—C1—C2—C4	106.0 (3)	C10—C11—C12—C13	-177.4 (3)
C10—C1—C2—C4	-136.5 (3)	C11—C12—C13—C14	-1.4 (4)
C10—C1—C2—C3	117.5 (3)	C12—C13—C14—C15	1.1 (5)
C2—C1—C3—C6	109.2 (3)	C13—C14—C15—C16	0.1 (5)
C10—C1—C3—C6	-4.9 (4)	C12—C11—C16—C15	0.9 (4)
C10—C1—C3—C2	-114.1 (3)	C10—C11—C16—C15	178.5 (3)
C5—C2—C3—C6	-3.5 (4)	C14—C15—C16—C11	-1.1 (5)
C1—C2—C3—C6	-115.0 (3)	O1—C10—C17—C18	1.3 (3)
C4—C2—C3—C6	137.5 (3)	C1—C10—C17—C18	-119.4 (3)
C5—C2—C3—C1	111.5 (3)	C11—C10—C17—C18	117.4 (3)
C4—C2—C3—C1	-107.5 (3)	O1—C10—C17—C22	179.1 (2)
C1—C3—C6—S1	71.5 (3)	C1—C10—C17—C22	58.5 (3)
C2—C3—C6—S1	144.1 (2)	C11—C10—C17—C22	-64.8 (3)
C1—C3—C6—S2	-165.9 (2)	C22—C17—C18—C19	0.8 (4)
C2—C3—C6—S2	-93.3 (3)	C10—C17—C18—C19	178.6 (3)
S1—C7—C8—C9	66.4 (3)	C17—C18—C19—C20	-1.7 (5)

C7—C8—C9—S2	-69.4 (3)	C18—C19—C20—C21	1.9 (5)
C2—C1—C10—O1	-51.7 (3)	C19—C20—C21—C22	-1.3 (5)
C3—C1—C10—O1	26.2 (4)	C20—C21—C22—C17	0.4 (5)
C2—C1—C10—C17	65.9 (3)	C18—C17—C22—C21	-0.2 (4)
C3—C1—C10—C17	143.8 (3)	C10—C17—C22—C21	-178.1 (3)
C2—C1—C10—C11	-172.3 (2)	C8—C7—S1—C6	-55.9 (2)
C3—C1—C10—C11	-94.4 (3)	C3—C6—S1—C7	173.11 (18)
O1—C10—C11—C16	-121.1 (3)	S2—C6—S1—C7	55.16 (17)
C17—C10—C11—C16	123.0 (3)	C8—C9—S2—C6	61.1 (2)
C1—C10—C11—C16	2.1 (3)	C3—C6—S2—C9	-176.75 (19)
O1—C10—C11—C12	56.5 (3)	S1—C6—S2—C9	-57.17 (18)
C17—C10—C11—C12	-59.4 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H2 \cdots S1	0.84	2.58	3.330 (2)	149
C7—H7A \cdots S2 ⁱ	0.99	2.89	3.736 (3)	144
C9—H9A \cdots O1 ⁱⁱ	0.99	2.60	3.236 (3)	122

Symmetry codes: (i) $-x+2, y-1/2, -z+1$; (ii) $-x+2, y+1/2, -z+1$.

Fig. 1

