

2-Chloro-5-nitroaniline

Aamer Saeed,^{a,*} Zaman Ashraf,^b Mahira Batool^a and Michael Bolte^c

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan,
^bRiphah Institute of Pharmaceutical Sciences, Islamabad, Pakistan, and ^cInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany
Correspondence e-mail: aamersaeed@yahoo.com


Received 19 May 2009; accepted 22 May 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.023; wR factor = 0.061; data-to-parameter ratio = 11.8.

The molecule of the title compound, $\text{C}_6\text{H}_5\text{ClN}_2\text{O}_2$, is close to being planar (rms deviation = 0.032 Å for all non-H atoms), with a maximum deviation of $-0.107(3)\text{ \AA}$ for an O atom. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ interactions link the molecules into a three-dimensional network.

Related literature

For applications of substituted nitrobenzene and aniline derivatives, see: Heinisch *et al.* (1997); Wang *et al.* (2000); Yosuke *et al.* (2003); Zou *et al.* (1997). For a related structure, see: Zhang *et al.* (2004). For bond-length data, see: Allen *et al.* (1987). For synthesis, see: Suwanprasop *et al.* (2003).



Experimental

Crystal data



$M_r = 172.57$

Monoclinic, $P2_1/n$
 $a = 13.6233(10)\text{ \AA}$
 $b = 3.7445(3)\text{ \AA}$
 $c = 13.6420(9)\text{ \AA}$
 $\beta = 91.768(5)^{\circ}$
 $V = 695.58(9)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.49\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.35 \times 0.34 \times 0.29\text{ mm}$

Data collection

Stoe IPDSII two-circle diffractometer
Absorption correction: multi-scan (*MULABS*; Blessing, 1995)
 $T_{\min} = 0.847$, $T_{\max} = 0.871$

5108 measured reflections
1300 independent reflections
1266 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.061$
 $S = 1.06$
1300 reflections
110 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A···O1 ⁱ	0.85 (2)	2.33 (2)	3.1521 (18)	163.2 (19)
N2—H2B···N2 ⁱⁱ	0.88 (3)	2.44 (2)	3.1452 (19)	137.4 (18)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-RED* (Stoe & Cie, 2001); data reduction: *X-RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2694).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Heinisch, G., Huber, E., Matusczak, B., Maurer, A. & Prillinge, U. (1997). *Arch. Pharm.* **330**, 29–34.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Stoe & Cie (2001). *X-AREA* and *X-RED*. Stoe & Cie, Darmstadt, Germany.
- Suwanprasop, S., Suksorn, S., Nhujak, T., Roengsumran, S. & Petsom, A. (2003). *Ind. Eng. Chem. Res.* **42**, 5054–5059.
- Wang, Y., Wu, Y. & Tian, H. (2000). *Dyes Pigm.* **44**, 93–100.
- Yosuke, M., Takako, U. & Tetsuo, M. (2003). Book of Abstracts, Congress of Heterocyclic Chemistry, Tokyo, Japan, Vol. 33, pp. 114–115.
- Zhang, Y., Feng, T.-Y. & Li, S.-Y. (2004). *Acta Cryst. E* **60**, o2501–o2502.
- Zou, R., Drach, J. C. & Townsend, L. B. (1997). *J. Med. Chem.* **40**, 811–818.

supporting information

Acta Cryst. (2009). E65, o1417 [doi:10.1107/S160053680901945X]

2-Chloro-5-nitroaniline

Aamer Saeed, Zaman Ashraf, Mahira Batool and Michael Bolte

S1. Comment

Substituted nitrobenzene and aniline derivatives are valuable as intermediates towards a variety of dye and pigments, heterocycles, pesticides, rubber chemicals and agricultural products, and are useful as textile printing agents, nickel stripping agents and polymerization catalysts. Thus, then title compound is an important intermediate or starting point in the syntheses of alkyl derivatives of 2-aminobenzenethiols substituted by chloro and nitro groups (Wang *et al.*, 2000), donor-bridge-acceptor' triad compounds containing the aromatic sulfur bridges (Yosuke *et al.*, 2003), pyridazinobenzodiazepin-5-ones as non-nucleoside HIV Reverse Transcriptase Inhibitors (Heinisch *et al.*, 1997) and 2-chloro-5,6-dihalo-*D*-ribofuranosylbenzimidazoles as potential agents for human cytomegalovirus infections (Zou *et al.*, 1997). We report herein the crystal structure of the title compound, as a key starting point towards many heterocycles.

In the molecule of the title compound (Fig 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges, and may be compared with the corresponding values in N-tert-butyl-4-chloro-5-methyl-2-nitroaniline (Zhang *et al.*, 2004). Ring A (C1–C6) is, of course, planar. Atoms C11, O1, O2, N1 and N2 are 0.019 (3), 0.104 (3), -0.107 (3), 0.003 (3) and -0.066 (3) Å away from the ring plane, respectively. So, the molecule is nearly planar.


In the crystal structure, intermolecular N—H···O and N—H···N hydrogen bonds (Table 1) link the molecules into a network, in which they may be effective in the stabilization of the structure.

S2. Experimental

The title compound was prepared by nitration and selective reduction of 4-nitroaniline according to the literature method (Suwanprasop *et al.*, 2003). Recrystallization from methanol afforded the title compound. Anal. calcd. for $C_6H_5ClN_2O_2$: C, 41.76; H, 2.92; N, 16.23%; found: C, 41.71; H, 2.97; N, 16.16%

S3. Refinement

H atoms (for NH_2) were located in a difference synthesis and refined isotropically. The remaining H atoms were positioned geometrically, with $C—H = 0.95$ Å for aromatic H and constrained to ride on their parent atoms, with $U_{iso}(H) = 1.2U_{eq}(C)$.

**Figure 1**

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

2-Chloro-5-nitroaniline

Crystal data

$C_6H_5ClN_2O_2$
 $M_r = 172.57$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 13.6233 (10)$ Å
 $b = 3.7445 (3)$ Å
 $c = 13.6420 (9)$ Å
 $\beta = 91.768 (5)^\circ$
 $V = 695.58 (9)$ Å³
 $Z = 4$

$F(000) = 352$
 $D_x = 1.648$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 7189 reflections
 $\theta = 3.4\text{--}25.9^\circ$
 $\mu = 0.49$ mm⁻¹
 $T = 173$ K
Block, orange
 $0.35 \times 0.34 \times 0.29$ mm

Data collection

Stoe IPDSII two-circle diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan (*MULABS*; Blessing, 1995)
 $T_{\min} = 0.847$, $T_{\max} = 0.871$

5108 measured reflections
1300 independent reflections
1266 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 3.4^\circ$
 $h = -16 \rightarrow 14$
 $k = -4 \rightarrow 4$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.061$
 $S = 1.06$
1300 reflections
110 parameters
0 restraints
Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0448P)^2 + 0.0466P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.21$ e Å⁻³
Extinction correction: *SHELXL97* (Sheldrick, 2008), $Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.034 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.91913 (3)	0.59906 (8)	0.62642 (3)	0.02963 (14)
O1	0.50548 (9)	0.8247 (4)	0.37207 (9)	0.0385 (3)
O2	0.60383 (10)	1.0787 (4)	0.27322 (9)	0.0475 (4)
N1	0.58750 (10)	0.9201 (3)	0.34951 (9)	0.0258 (3)
N2	0.70761 (10)	0.4832 (4)	0.66629 (9)	0.0255 (3)
H2B	0.7582 (18)	0.367 (5)	0.6920 (18)	0.043 (6)*
H2A	0.6514 (17)	0.384 (4)	0.6685 (15)	0.029 (5)*
C1	0.82203 (10)	0.6968 (3)	0.54548 (10)	0.0215 (3)
C2	0.72634 (11)	0.6173 (3)	0.57428 (10)	0.0203 (3)
C3	0.64960 (10)	0.6935 (3)	0.50720 (10)	0.0206 (3)
H3	0.5836	0.6427	0.5230	0.025*
C4	0.67060 (10)	0.8434 (3)	0.41772 (10)	0.0206 (3)
C5	0.76477 (11)	0.9244 (3)	0.38893 (11)	0.0231 (3)
H5	0.7764	1.0285	0.3269	0.028*
C6	0.84118 (11)	0.8461 (4)	0.45509 (11)	0.0248 (3)
H6	0.9070	0.8952	0.4383	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0223 (2)	0.03330 (19)	0.0329 (2)	0.00210 (13)	-0.00586 (12)	-0.00076 (14)
O1	0.0221 (6)	0.0617 (7)	0.0315 (6)	-0.0054 (5)	-0.0011 (4)	0.0122 (5)
O2	0.0374 (7)	0.0750 (10)	0.0298 (6)	-0.0099 (6)	-0.0040 (5)	0.0271 (6)
N1	0.0252 (7)	0.0302 (6)	0.0218 (6)	-0.0012 (5)	0.0000 (5)	0.0029 (5)
N2	0.0266 (7)	0.0291 (6)	0.0206 (6)	-0.0039 (6)	-0.0011 (5)	0.0041 (5)
C1	0.0215 (7)	0.0184 (5)	0.0245 (7)	0.0006 (5)	-0.0014 (5)	-0.0047 (5)
C2	0.0246 (7)	0.0181 (6)	0.0182 (7)	-0.0004 (5)	0.0014 (5)	-0.0026 (4)
C3	0.0204 (6)	0.0216 (6)	0.0201 (6)	-0.0019 (5)	0.0034 (5)	-0.0021 (5)
C4	0.0222 (7)	0.0196 (6)	0.0198 (6)	0.0001 (5)	0.0006 (5)	-0.0015 (5)
C5	0.0257 (7)	0.0235 (6)	0.0203 (7)	-0.0021 (5)	0.0060 (6)	-0.0007 (5)
C6	0.0199 (7)	0.0247 (6)	0.0301 (7)	-0.0009 (5)	0.0054 (5)	-0.0027 (5)

Geometric parameters (\AA , $^\circ$)

N1—O1	1.2216 (19)	C2—C3	1.397 (2)
N1—O2	1.2245 (18)	C3—C4	1.382 (2)

N2—H2B	0.88 (3)	C3—H3	0.9500
N2—H2A	0.85 (2)	C4—C5	1.387 (2)
C1—C6	1.386 (2)	C4—N1	1.4713 (18)
C1—C2	1.405 (2)	C5—C6	1.388 (2)
C1—Cl1	1.7357 (14)	C5—H5	0.9500
C2—N2	1.3830 (18)	C6—H6	0.9500
O1—N1—O2	123.19 (14)	C4—C3—C2	119.37 (12)
O1—N1—C4	118.40 (12)	C4—C3—H3	120.3
O2—N1—C4	118.41 (13)	C2—C3—H3	120.3
C2—N2—H2B	112.2 (15)	C3—C4—C5	123.90 (13)
C2—N2—H2A	112.4 (14)	C3—C4—N1	117.51 (12)
H2B—N2—H2A	117.7 (18)	C5—C4—N1	118.59 (13)
C6—C1—C2	122.42 (13)	C4—C5—C6	116.84 (13)
C6—C1—Cl1	119.36 (11)	C4—C5—H5	121.6
C2—C1—Cl1	118.22 (11)	C6—C5—H5	121.6
N2—C2—C3	120.88 (13)	C1—C6—C5	120.38 (13)
N2—C2—C1	121.98 (13)	C1—C6—H6	119.8
C3—C2—C1	117.09 (13)	C5—C6—H6	119.8
C6—C1—C2—N2	-177.05 (13)	N1—C4—C5—C6	-179.64 (12)
Cl1—C1—C2—N2	3.57 (17)	C2—C1—C6—C5	0.2 (2)
C6—C1—C2—C3	0.33 (19)	Cl1—C1—C6—C5	179.57 (10)
Cl1—C1—C2—C3	-179.04 (9)	C4—C5—C6—C1	-0.5 (2)
N2—C2—C3—C4	176.87 (12)	C3—C4—N1—O1	-5.21 (19)
C1—C2—C3—C4	-0.55 (19)	C5—C4—N1—O1	174.73 (13)
C2—C3—C4—C5	0.2 (2)	C3—C4—N1—O2	174.29 (13)
C2—C3—C4—N1	-179.82 (12)	C5—C4—N1—O2	-5.8 (2)
C3—C4—C5—C6	0.3 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2A···O1 ⁱ	0.85 (2)	2.33 (2)	3.1521 (18)	163.2 (19)
N2—H2B···N2 ⁱⁱ	0.88 (3)	2.44 (2)	3.1452 (19)	137.4 (18)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+3/2, y-1/2, -z+3/2$.