

# A neutral cubane with a $Zn_4O_4$ core: tetrabenzoatotetrakis( $\mu_3$ -hydroxydi-2-pyridylmethanolato)tetrzinc(II)–acetone–methanol (1/2/1)

Dong Hoon Shin,<sup>a</sup> Sim-Hee Han,<sup>b</sup> Pan-Gi Kim,<sup>c</sup> Cheal Kim<sup>a\*</sup> and Youngmee Kim<sup>d\*</sup>

<sup>a</sup>Department of Fine Chemistry, and Eco-Product and Materials Education Center, Seoul National University of Technology, Seoul 139-743, Republic of Korea,

<sup>b</sup>Division of Forest Genetic Resources, Korea Forest Research Institute, Suwon, Gyeonggi-Do 441-350, Republic of Korea, <sup>c</sup>Department of Forest & Environment Resources, Kyungpook National University, Sangju 742-711, Republic of Korea, and

<sup>d</sup>Department of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Republic of Korea

Correspondence e-mail: chealkim@sunt.ac.kr, ymeekim@ewha.ac.kr


Received 28 April 2009; accepted 12 May 2009

Key indicators: single-crystal X-ray study;  $T = 170$  K; mean  $\sigma(C-C) = 0.005$  Å; disorder in solvent or counterion;  $R$  factor = 0.034;  $wR$  factor = 0.110; data-to-parameter ratio = 18.9.

In the title compound,  $[Zn_4(C_{11}H_9N_2O_2)_4(C_7H_5O_2)_4] \cdot 2(CH_3)_2CO \cdot CH_3OH$ , the tetranuclear molecule lies on a fourfold inversion axis.  $Zn^{II}$  ions and  $\mu_3$ -O atoms in the cubane core occupy alternating vertices, forming two interpenetrating tetrahedra. Each  $Zn^{II}$  ion is further coordinated by two N atoms from two different  $(py)_2C(OH)O$  ligands ( $py$  is pyridyl) and one O atom from a monodentate benzoate ligand, forming a distorted octahedral environment. The  $(py)_2C(OH)O$  ligand acts in an  $\eta^1:\eta^3:\eta^1:\mu_3$  manner, forming two five-membered  $ZnNCCO$  chelating rings with two different  $Zn^{II}$  atoms sharing a common C–O bond, and an alkoxide-type bond to a third  $Zn^{II}$  ion. There are four symmetry-related intramolecular  $O-H \cdots O$  hydrogen bonds between the two types of ligands. In the asymmetric unit, there is a half-occupancy acetone solvent molecule and a half-occupancy methanol solvent molecule that lies on a twofold rotation axis.

## Related literature

For background to transition metal ions as the major cationic contributors to the inorganic composition of natural water and biological fluids, see: Daniele *et al.* (2008); For related crystal structures, see: Lee *et al.* (2008); Park *et al.* (2008); Yu *et al.* (2008); Stoumpos *et al.* (2008); Papaefstathiou & Perlepes (2002); Papatriantafyllopoulou *et al.* (2007).



## Experimental

### Crystal data

$[Zn_4(C_{11}H_9N_2O_2)_4(C_7H_5O_2)_4] \cdot 2C_3H_6O \cdot CH_4O$   
 $M_r = 6795.70$   
 Tetragonal,  $I\bar{4}2d$   
 $a = 14.3201(4)$  Å  
 $c = 37.730(2)$  Å

$V = 7737.1(5)$  Å<sup>3</sup>  
 $Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.30$  mm<sup>-1</sup>  
 $T = 170$  K  
 $0.10 \times 0.08 \times 0.05$  mm

### Data collection

Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 1997)  
 $T_{min} = 0.883$ ,  $T_{max} = 0.937$

22787 measured reflections  
 4628 independent reflections  
 4279 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.029$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.110$   
 $S = 1.09$   
 4628 reflections  
 245 parameters  
 5 restraints

H-atom parameters constrained  
 $\Delta\rho_{max} = 1.37$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.42$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983),  
 2045 Friedel pairs  
 Flack parameter: 0.002 (13)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O2-H2O \cdots O4$	0.86	1.81	2.664 (3)	172

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the Korea Ministry of the Environment 'ET-Human resource development Project' and the Cooperative Research Program for Agricultural Science & Technology Development (20070301-036-019-02) is gratefully acknowledged.

---

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2815).

---

## References

- Bruker (1997). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Daniele, P. G., Foti, C., Gianguzza, A., Prenesti, E. & Sammartano, S. (2008). *Coord. Chem. Rev.* **252**, 1093–1107.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Lee, E. Y., Park, B. K., Kim, C., Kim, S.-J. & Kim, Y. (2008). *Acta Cryst.* **E64**, m286.
- Papaefstathiou, G. S. & Perlepes, S. P. (2002). *Comments Inorg. Chem.* **23**, 249–274.
- Papatriantafyllopoulou, C., Efthymiou, C. G., Raptopoulou, C. P., Vicente, R., Manessi-Zoupa, E., Psycharis, V., Escuer, A. & Perlepes, S. P. (2007). *J. Mol. Struct.* **829**, 176–188.
- Park, B. K., Jang, K.-H., Kim, P.-G., Kim, C. & Kim, Y. (2008). *Acta Cryst.* **E64**, m1141.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoumpos, C. C., Gass, I. A., Milios, C. J., Kefalloniti, E., Raptopoulou, C. P., Terzis, A., Lalioti, N., Brechin, E. K. & Perlepes, S. P. (2008). *Inorg. Chem. Commun.* **11**, 196–202.
- Yu, S. M., Park, C.-H., Kim, P.-G., Kim, C. & Kim, Y. (2008). *Acta Cryst.* **E64**, m881–m882.

## supporting information

*Acta Cryst.* (2009). E65, m658–m659 [doi:10.1107/S1600536809017772]

## A neutral cubane with a $\text{Zn}^{\text{II}}_4\text{O}_4$ core: tetrabenzoatotetrakis( $\mu_3$ -hydroxydi-2-pyridylmethanolato)tetrzinc(II)–acetone–methanol (1/2/1)

Dong Hoon Shin, Sim-Hee Han, Pan-Gi Kim, Cheal Kim and Youngmee Kim

### S1. Comment

Transition metal ions have, recently, received attention as the major cationic contributors to the inorganic composition of natural water and biological fluids (Daniele, *et al.*, 2008). While the main interest is focused on the interaction of transition metal ions with biologically active molecules such as amino acids, proteins, sugars and nucleotides, the study on the interaction of the transition metal ions with fulvic acids and humic acids, mainly found in soil, is in incipient stages. As models to examine the interaction, therefore, we have previously used copper(II) benzoate as a building block and reported the structures of copper(II) benzoates with quinoxaline, 6-methylquinoline, and 3-methylquinoline (Lee, *et al.*, 2008; Yu, *et al.*, 2008; Park, *et al.*, 2008). In this work, we have employed zinc(II) benzoate as a building block and di-2-pyridyl ketone as a ligand. We report herein the structure of the product of zinc(II) benzoate with di-2-pyridyl ketone.


The crystal structure contains tetranuclear  $[\text{Zn}_4(\text{O}_2\text{CPh})_4\{(\text{py})_2\text{C}(\text{OH})\text{O}\}_4]$  molecules (Fig. 1), similar to the corresponding  $\text{Mn}_4$  cubane compound (Stoumpos, *et al.*, 2008). The tetramolecular molecule lies on a fourfold inversion center and hence the asymmetric unit contains a quarter of a molecule.  $\text{Zn}^{\text{II}}$  ions and  $\mu_3$ -O atoms in the cubane  $[\text{Zn}^{\text{II}}_4(\mu_3\text{-OR})_4]^{4+}$  core occupy alternate vertices. Thus, the molecule consists of two interpenetrating tetrahedra: one contains four  $\mu_3$ -O atoms originating from the  $(\text{py})_2\text{C}(\text{OH})\text{O}$  ligands, and the other contains four  $\text{Zn}^{\text{II}}$  atoms. Each  $\text{Zn}^{\text{II}}$  center is coordinated by two N atoms from two different  $(\text{py})_2\text{C}(\text{OH})\text{O}$  ligands and one O atom from a monodentate  $\text{PhCO}_2$  ligand to form a distorted octahedral geometry. The  $(\text{py})_2\text{C}(\text{OH})\text{O}$  ligands acts as  $\eta^1:\eta^3:\eta^1:\mu_3$  to form two five-membered  $\text{ZnNCCO}$  chelating rings with two different  $\text{Zn}^{\text{II}}$  ions sharing a common C—O edge and an alkoxide-type bond to a third  $\text{Zn}^{\text{II}}$  ion. This ligation mode is common for the hydrated di-2-pyridyl ketone,  $(\text{py})_2\text{C}(\text{OH})\text{O}^-$  (Papaefstathiou & Perlepes, 2002; Papatriantafyllopoulou, *et al.*, 2007). There are intramolecular hydrogen bonds interactions between the protonated O atom of the  $(\text{py})_2\text{C}(\text{OH})\text{O}$  ligand and the uncoordinated O atom of the monodentate  $\text{PhCO}_2$  group.

### S2. Experimental

38.0 mg (0.125 mmol) of  $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  and 35.5 mg (0.25 mmol) of  $\text{C}_6\text{H}_5\text{COONH}_4$  were dissolved in 4 ml water and carefully layered by 4 ml solution of a mixture of acetone, methanol and ethanol (2/2/2) of di-2-pyridyl ketone ligand (46.1 mg, 0.25 mmol). Crystals of the title compound suitable for X-ray analysis were obtained in a few weeks.

### S3. Refinement

H atoms were placed in calculated positions with C—H distances of 0.93–0.98 Å and O—H = 0.82 Å. They were included in the refinement in a riding-motion approximation with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}} \text{ and O})$ .

**Figure 1**

Partially labeled molecular structure of the title complex. Displacement ellipsoids are shown at the 30% probability level. The green dotted lines represent hydrogen bonds. Solvent molecules are not shown.

### Tetrabenzoatotetrakis( $\mu_3$ -hydroxydi-2-pyridylmethanolato)tetrazinc(II)- acetone-methanol (1/2/1)

#### Crystal data

$[\text{Zn}_4(\text{C}_{11}\text{H}_9\text{N}_2\text{O}_2)_4(\text{C}_7\text{H}_5\text{O}_2)_4] \cdot 2\text{C}_3\text{H}_6\text{O} \cdot \text{CH}_4\text{O}$

$M_r = 6795.70$

Tetragonal,  $I\bar{4}2d$

Hall symbol: I -4 2bw

$a = 14.3201(4) \text{ \AA}$

$c = 37.730(2) \text{ \AA}$

$V = 7737.1(5) \text{ \AA}^3$

$Z = 1$

$F(000) = 3496$

$D_x = 1.458 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 8208 reflections

$\theta = 2.8\text{--}27.0^\circ$

$\mu = 1.30 \text{ mm}^{-1}$

$T = 170 \text{ K}$

Polyhedron, colorless

$0.10 \times 0.08 \times 0.05 \text{ mm}$

#### Data collection

Bruker SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 1997)

$T_{\min} = 0.883$ ,  $T_{\max} = 0.937$

22787 measured reflections

4628 independent reflections

4279 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$   
 $\theta_{\text{max}} = 28.3^\circ$ ,  $\theta_{\text{min}} = 1.5^\circ$   
 $h = -18 \rightarrow 18$

$k = -9 \rightarrow 18$   
 $l = -48 \rightarrow 48$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.110$   
 $S = 1.09$   
 4628 reflections  
 245 parameters  
 5 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0787P)^2 + 0.4409P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 1.37 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.42 \text{ e } \text{\AA}^{-3}$   
 Absolute structure: Flack (1983), 2045 Friedel  
 pairs  
 Absolute structure parameter: 0.002 (13)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.61621 (2)	0.51226 (2)	0.470190 (8)	0.01677 (10)	
O1	0.50301 (15)	0.59983 (13)	0.47540 (5)	0.0177 (4)	
O2	0.53176 (14)	0.75354 (15)	0.45808 (6)	0.0237 (5)	
H2O	0.5908	0.7429	0.4575	0.036*	
O3	0.73924 (15)	0.58089 (15)	0.47172 (6)	0.0257 (4)	
O4	0.71640 (16)	0.73233 (17)	0.46138 (9)	0.0390 (6)	
N1	0.58789 (18)	0.55925 (18)	0.41603 (6)	0.0197 (5)	
N2	0.67260 (17)	0.38313 (19)	0.45377 (6)	0.0204 (5)	
C1	0.6300 (2)	0.5287 (2)	0.38641 (8)	0.0262 (6)	
H1	0.6712	0.4768	0.3881	0.031*	
C2	0.6162 (3)	0.5691 (3)	0.35381 (9)	0.0366 (8)	
H2	0.6465	0.5455	0.3333	0.044*	
C3	0.5566 (3)	0.6455 (3)	0.35178 (10)	0.0404 (10)	
H3	0.5448	0.6748	0.3296	0.049*	
C4	0.5141 (3)	0.6789 (3)	0.38253 (8)	0.0331 (8)	
H4	0.4741	0.7318	0.3818	0.040*	
C5	0.5320 (2)	0.6328 (2)	0.41431 (8)	0.0210 (6)	
C6	0.4901 (2)	0.6676 (2)	0.44992 (7)	0.0187 (5)	
C7	0.6154 (2)	0.3128 (2)	0.44535 (7)	0.0192 (5)	
C8	0.6484 (2)	0.2278 (2)	0.43286 (9)	0.0256 (6)	

H8	0.6064	0.1783	0.4277	0.031*	
C9	0.7445 (3)	0.2165 (2)	0.42796 (10)	0.0325 (8)	
H9	0.7685	0.1604	0.4181	0.039*	
C10	0.8032 (2)	0.2871 (3)	0.43751 (10)	0.0330 (8)	
H10	0.8689	0.2797	0.4353	0.040*	
C11	0.7658 (2)	0.3707 (2)	0.45060 (9)	0.0280 (7)	
H11	0.8069	0.4197	0.4574	0.034*	
C12	0.7640 (2)	0.6651 (2)	0.47105 (9)	0.0227 (6)	
C13	0.8626 (2)	0.6836 (2)	0.48381 (8)	0.0237 (6)	
C14	0.9023 (2)	0.7726 (2)	0.48046 (11)	0.0341 (8)	
H14	0.8676	0.8218	0.4699	0.041*	
C15	0.9924 (3)	0.7888 (3)	0.49264 (12)	0.0461 (10)	
H15	1.0196	0.8489	0.4900	0.055*	
C16	1.0423 (3)	0.7181 (3)	0.50846 (13)	0.0474 (11)	
H16	1.1039	0.7295	0.5168	0.057*	
C17	1.0032 (3)	0.6313 (3)	0.51216 (11)	0.0398 (8)	
H17	1.0374	0.5829	0.5234	0.048*	
C18	0.9141 (2)	0.6138 (3)	0.49960 (9)	0.0306 (7)	
H18	0.8882	0.5530	0.5019	0.037*	
O1S	0.9292 (16)	1.1086 (15)	0.3972 (6)	0.218 (6)*	0.50
C1S	0.9466 (13)	1.0280 (17)	0.3923 (6)	0.218 (6)*	0.50
C2S	0.965 (2)	0.972 (2)	0.4275 (8)	0.218 (6)*	0.50
H2S1	1.0259	0.9886	0.4372	0.328*	0.50
H2S2	0.9634	0.9045	0.4224	0.328*	0.50
H2S3	0.9160	0.9865	0.4448	0.328*	0.50
C3S	0.9503 (19)	0.975 (2)	0.3559 (7)	0.218 (6)*	0.50
H3S1	0.8956	0.9340	0.3537	0.328*	0.50
H3S2	1.0073	0.9370	0.3547	0.328*	0.50
H3S3	0.9504	1.0201	0.3364	0.328*	0.50
O2S	0.7500	1.0096 (11)	0.3750	0.110 (5)*	0.50
H2S	0.7197	1.0292	0.3574	0.164*	0.25
C21S	0.7500	0.9049 (11)	0.3750	0.139 (10)*	0.50
H21A	0.7999	0.8821	0.3595	0.208*	0.25
H21B	0.6896	0.8821	0.3664	0.208*	0.25
H21C	0.7605	0.8821	0.3992	0.208*	0.25

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.01658 (17)	0.01739 (17)	0.01634 (15)	0.00064 (12)	0.00086 (12)	0.00068 (12)
O1	0.0189 (9)	0.0175 (9)	0.0167 (9)	0.0029 (8)	0.0001 (8)	0.0034 (7)
O2	0.0206 (10)	0.0193 (10)	0.0311 (11)	-0.0009 (9)	0.0009 (8)	-0.0005 (9)
O3	0.0216 (11)	0.0258 (11)	0.0296 (11)	-0.0035 (9)	0.0013 (9)	-0.0002 (10)
O4	0.0214 (11)	0.0262 (13)	0.0693 (19)	-0.0015 (9)	-0.0062 (11)	0.0019 (12)
N1	0.0201 (12)	0.0213 (12)	0.0176 (11)	0.0003 (9)	0.0009 (9)	0.0007 (9)
N2	0.0182 (12)	0.0232 (12)	0.0199 (11)	0.0025 (11)	0.0015 (9)	0.0025 (10)
C1	0.0244 (15)	0.0299 (16)	0.0244 (14)	0.0035 (13)	0.0059 (12)	0.0000 (12)
C2	0.044 (2)	0.044 (2)	0.0218 (15)	0.0112 (18)	0.0101 (15)	0.0016 (14)

C3	0.050 (2)	0.048 (2)	0.0227 (16)	0.0158 (19)	0.0113 (16)	0.0114 (15)
C4	0.0378 (19)	0.0376 (18)	0.0238 (15)	0.0120 (16)	0.0058 (14)	0.0085 (13)
C5	0.0206 (14)	0.0214 (14)	0.0211 (14)	0.0008 (12)	0.0015 (10)	0.0043 (11)
C6	0.0199 (14)	0.0168 (13)	0.0194 (12)	0.0026 (12)	0.0006 (11)	0.0040 (10)
C7	0.0206 (14)	0.0207 (13)	0.0163 (12)	0.0010 (12)	0.0005 (11)	0.0006 (10)
C8	0.0267 (15)	0.0255 (16)	0.0245 (15)	0.0056 (13)	-0.0002 (12)	-0.0017 (12)
C9	0.0317 (18)	0.0298 (17)	0.0361 (18)	0.0124 (15)	0.0060 (15)	-0.0024 (14)
C10	0.0231 (16)	0.0371 (19)	0.0388 (19)	0.0086 (14)	0.0068 (14)	0.0018 (15)
C11	0.0214 (15)	0.0312 (17)	0.0315 (17)	-0.0005 (13)	0.0033 (12)	0.0037 (14)
C12	0.0192 (14)	0.0267 (15)	0.0221 (14)	-0.0017 (11)	0.0049 (12)	-0.0030 (13)
C13	0.0204 (15)	0.0263 (16)	0.0246 (14)	-0.0009 (12)	0.0040 (12)	-0.0069 (12)
C14	0.0275 (18)	0.0260 (17)	0.049 (2)	-0.0028 (14)	0.0040 (14)	-0.0056 (14)
C15	0.0283 (18)	0.0291 (18)	0.081 (3)	-0.0061 (16)	-0.001 (2)	-0.0103 (19)
C16	0.0218 (17)	0.048 (2)	0.072 (3)	-0.0035 (16)	-0.0083 (18)	-0.020 (2)
C17	0.0292 (17)	0.038 (2)	0.052 (2)	0.0050 (16)	-0.0075 (17)	-0.0062 (17)
C18	0.0270 (16)	0.0290 (16)	0.0357 (17)	-0.0039 (14)	-0.0012 (13)	-0.0028 (14)

*Geometric parameters (Å, °)*

Zn1—O3	2.018 (2)	C9—C10	1.365 (5)
Zn1—O1	2.059 (2)	C9—H9	0.9500
Zn1—O1 <sup>i</sup>	2.0776 (19)	C10—C11	1.401 (5)
Zn1—N2	2.111 (3)	C10—H10	0.9500
Zn1—N1	2.189 (2)	C11—H11	0.9500
Zn1—O1 <sup>ii</sup>	2.351 (2)	C12—C13	1.515 (4)
O1—C6	1.378 (3)	C13—C18	1.378 (5)
O1—Zn1 <sup>iii</sup>	2.0777 (19)	C13—C14	1.401 (5)
O1—Zn1 <sup>ii</sup>	2.352 (2)	C14—C15	1.389 (5)
O2—C6	1.402 (4)	C14—H14	0.9500
O2—H2O	0.8587	C15—C16	1.376 (6)
O3—C12	1.257 (4)	C15—H15	0.9500
O4—C12	1.235 (4)	C16—C17	1.371 (6)
N1—C5	1.325 (4)	C16—H16	0.9500
N1—C1	1.343 (4)	C17—C18	1.384 (5)
N2—C7	1.336 (4)	C17—H17	0.9500
N2—C11	1.352 (4)	C18—H18	0.9500
C1—C2	1.374 (5)	O1S—C1S	1.195 (10)
C1—H1	0.9500	C1S—C2S	1.57 (2)
C2—C3	1.391 (5)	C1S—C3S	1.57 (2)
C2—H2	0.9500	C2S—H2S1	0.9800
C3—C4	1.394 (5)	C2S—H2S2	0.9800
C3—H3	0.9500	C2S—H2S3	0.9800
C4—C5	1.392 (4)	C3S—H3S1	0.9800
C4—H4	0.9500	C3S—H3S2	0.9800
C5—C6	1.553 (4)	C3S—H3S3	0.9800
C6—C7 <sup>ii</sup>	1.546 (4)	O2S—C21S	1.499 (2)
C7—C8	1.388 (4)	O2S—H2S	0.8400
C7—C6 <sup>i</sup>	1.546 (4)	C21S—H21A	0.9800

C8—C9	1.398 (5)	C21S—H21B	0.9800
C8—H8	0.9500	C21S—H21C	0.9800
O3—Zn1—O1	112.83 (9)	C7—C8—H8	120.6
O3—Zn1—O1 <sup>i</sup>	96.98 (9)	C9—C8—H8	120.6
O1—Zn1—O1 <sup>i</sup>	83.16 (8)	C10—C9—C8	119.1 (3)
O3—Zn1—N2	95.80 (9)	C10—C9—H9	120.5
O1—Zn1—N2	149.64 (9)	C8—C9—H9	120.5
O1 <sup>i</sup> —Zn1—N2	103.94 (9)	C9—C10—C11	119.4 (3)
O3—Zn1—N1	92.22 (9)	C9—C10—H10	120.3
O1—Zn1—N1	75.88 (9)	C11—C10—H10	120.3
O1 <sup>i</sup> —Zn1—N1	159.01 (9)	N2—C11—C10	121.4 (3)
N2—Zn1—N1	93.79 (10)	N2—C11—H11	119.3
O3—Zn1—O1 <sup>ii</sup>	164.53 (8)	C10—C11—H11	119.3
O1—Zn1—O1 <sup>ii</sup>	80.57 (8)	O4—C12—O3	126.7 (3)
O1 <sup>i</sup> —Zn1—O1 <sup>ii</sup>	76.33 (8)	O4—C12—C13	118.1 (3)
N2—Zn1—O1 <sup>ii</sup>	72.80 (8)	O3—C12—C13	115.2 (3)
N1—Zn1—O1 <sup>ii</sup>	98.82 (8)	C18—C13—C14	118.8 (3)
C6—O1—Zn1	117.87 (17)	C18—C13—C12	120.6 (3)
C6—O1—Zn1 <sup>iii</sup>	126.55 (17)	C14—C13—C12	120.6 (3)
Zn1—O1—Zn1 <sup>iii</sup>	104.24 (9)	C15—C14—C13	119.9 (4)
C6—O1—Zn1 <sup>ii</sup>	108.97 (17)	C15—C14—H14	120.0
Zn1—O1—Zn1 <sup>ii</sup>	98.51 (8)	C13—C14—H14	120.1
Zn1 <sup>iii</sup> —O1—Zn1 <sup>ii</sup>	94.78 (7)	C16—C15—C14	120.2 (4)
C6—O2—H2O	104.9	C16—C15—H15	119.9
C12—O3—Zn1	135.5 (2)	C14—C15—H15	119.9
C5—N1—C1	119.3 (3)	C17—C16—C15	119.9 (4)
C5—N1—Zn1	113.71 (19)	C17—C16—H16	120.0
C1—N1—Zn1	126.4 (2)	C15—C16—H16	120.0
C7—N2—C11	119.1 (3)	C16—C17—C18	120.5 (4)
C7—N2—Zn1	119.66 (19)	C16—C17—H17	119.8
C11—N2—Zn1	121.3 (2)	C18—C17—H17	119.8
N1—C1—C2	122.9 (3)	C13—C18—C17	120.6 (4)
N1—C1—H1	118.6	C13—C18—H18	119.7
C2—C1—H1	118.5	C17—C18—H18	119.7
C1—C2—C3	118.0 (3)	O1S—C1S—C2S	114 (2)
C1—C2—H2	121.0	O1S—C1S—C3S	128 (2)
C3—C2—H2	121.0	C2S—C1S—C3S	119 (2)
C2—C3—C4	119.4 (3)	C1S—C2S—H2S1	109.5
C2—C3—H3	120.3	C1S—C2S—H2S2	109.4
C4—C3—H3	120.3	H2S1—C2S—H2S2	109.5
C5—C4—C3	118.3 (3)	C1S—C2S—H2S3	109.5
C5—C4—H4	120.9	H2S1—C2S—H2S3	109.5
C3—C4—H4	120.9	H2S2—C2S—H2S3	109.5
N1—C5—C4	122.0 (3)	C1S—C3S—H3S1	109.5
N1—C5—C6	116.5 (2)	C1S—C3S—H3S2	109.4
C4—C5—C6	121.4 (3)	H3S1—C3S—H3S2	109.5
O1—C6—O2	114.1 (2)	C1S—C3S—H3S3	109.5



O1—C6—C7 <sup>ii</sup>	109.6 (2)	H3S1—C3S—H3S3	109.5
O2—C6—C7 <sup>ii</sup>	106.3 (2)	H3S2—C3S—H3S3	109.5
O1—C6—C5	109.0 (2)	C21S—O2S—H2S	109.5
O2—C6—C5	107.9 (2)	O2S—C21S—H21A	109.00
C7 <sup>ii</sup> —C6—C5	109.8 (2)	O2S—C21S—H21B	109.00
N2—C7—C8	122.1 (3)	H21A—C21S—H21B	110.00
N2—C7—C6 <sup>ii</sup>	115.9 (2)	O2S—C21S—H21C	109.00
C8—C7—C6 <sup>ii</sup>	122.0 (3)	H21A—C21S—H21C	109.00
C7—C8—C9	118.8 (3)	H21B—C21S—H21C	109.00

Symmetry codes: (i)  $y, -x+1, -z+1$ ; (ii)  $-x+1, -y+1, z$ ; (iii)  $-y+1, x, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O2—H2O...O4	0.86	1.81	2.664 (3)	172