

## ***N,N'-(Phenyliminodimethylene)diprop-2-enamide hemihydrate***

Dhanapal Tamilvendan,<sup>a</sup> Ganesan Venkatesa Prabhu,<sup>a</sup> Frank R. Fronczeck<sup>b</sup> and Nagarajan Vembu<sup>c\*</sup>

<sup>a</sup>Department of Chemistry, National Institute of Technology, Tiruchirappalli 620 015, India,

<sup>b</sup>Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803-1804, USA, and <sup>c</sup>Department of Chemistry, Urumu Dhanalakshmi College, Tiruchirappalli 620 019, India

Correspondence e-mail: vembu57@yahoo.com


Received 3 May 2009; accepted 12 May 2009

Key indicators: single-crystal X-ray study;  $T = 90$  K; mean  $\sigma(C-C) = 0.002$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.120; data-to-parameter ratio = 20.3.

In the title compound,  $C_{14}H_{17}N_3O_2 \cdot 0.5H_2O$ , the asymmetric unit consists of an *N,N'*-(phenyliminodimethylene)diprop-2-enamide molecule and one half-molecule of water, with the O atom of the latter having 2 site symmetry. The supramolecular architecture is framed by the interplay of two-dimensional networks of both O—H···O and N—H···O interactions supported by C—H···O and edge-to-face C—H···π interactions.

### Related literature

For a detailed description of Mannich bases and their applications, see: Friedrich *et al.* (1991); Bohme & Mannich (1955); Afsah *et al.* (2008); Terzioglu *et al.* (2006); Ravichandran *et al.* (2007); Pandeya *et al.* (2000). For hydrogen bonds, see: Desiraju & Steiner (1999); Jeffrey (1997). For hydrogen-bond motifs, see: Bernstein *et al.* (1995); Etter (1990).



### Experimental

#### Crystal data



$M_r = 268.31$

Orthorhombic,  $Pbcn$

$a = 17.074$  (2) Å

$b = 9.8366$  (15) Å

$c = 16.316$  (2) Å

$V = 2740.3$  (6) Å<sup>3</sup>

$Z = 8$

Mo  $K\alpha$  radiation

$\mu = 0.09$  mm<sup>-1</sup>

$T = 90$  K

$0.30 \times 0.23 \times 0.12$  mm

#### Data collection

Nonius KappaCCD diffractometer  
with an Oxford Cryosystems  
Cryostream cooler  
Absorption correction: none

9467 measured reflections  
5065 independent reflections  
3885 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.120$   
 $S = 1.02$   
5065 reflections

249 parameters  
All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.39$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.28$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N5—H5···O11	0.876 (15)	2.318 (15)	3.0476 (11)	140.8 (12)
C4—H4A···O7	0.984 (13)	2.366 (13)	2.8089 (12)	106.5 (9)
C15—H15···O7 <sup>i</sup>	0.953 (14)	2.563 (14)	3.4922 (14)	165.1 (10)
OW—HW···O7 <sup>i</sup>	0.845 (17)	1.990 (17)	2.8193 (9)	166.7 (16)
N1—H1···O1 <sup>ii</sup>	0.891 (15)	2.089 (15)	2.9651 (11)	167.4 (14)
C2—H2B···Cg1 <sup>iii</sup>	0.966 (13)	3.178	3.874	130.40
C8—H8···Cg1 <sup>iv</sup>	0.966 (16)	2.571 (15)	3.4444 (12)	150.6 (13)

Symmetry codes: (i)  $-x + 2, y, -z - \frac{1}{2}$ ; (ii)  $-x + \frac{3}{2}, y + \frac{1}{2}, z$ ; (iii)  $-x + 2, -y + 2, -z$ ; (iv)  $-x + \frac{1}{2}, y - \frac{3}{2}, z$ . Cg1 is the centroid of the C14—C19 ring.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The purchase of the diffractometer was made possible by grant No. LEQSF (1999–2000)-ENH-TR-13, administered by the Louisiana Board of Regents.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2820).

### References

- Afsah, E. M., Hammouda, M., Khalifa, M. M. & Al-shahaby, E. H. (2008). *Z. Naturforsch. Teil B*, **63**, 577–584.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. (1995). *Angew. Chem. Int. Ed. Engl.*, **34**, 1555–1573.
- Bohme, H. & Mannich, C. (1955). *Chem. Ber.*, **88**, I–XXVI.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology*. New York: Oxford University Press Inc.
- Etter, M. C. (1990). *Acc. Chem. Res.*, **23**, 120–126.
- Friedrich, C., Dallman, A., Mannich, C. & Pharmazie, D. (1991). *Pharm. Ztg.*, **136**, 691–701.
- Jeffrey, G. A. (1997). *An Introduction to Hydrogen Bonding*. New York: Oxford University Press Inc.
- Nonius, B. V. (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pandeya, S. N., Sriram, D., De Clercq, E. & Nath, G. (2000). *Arzneim. Forsch.*, **35**, 249–255.
- Ravichandran, V., Mohan, S. & Suresh Kumar, K. (2007). *Arkivoc Newslett.*, **14**, 51–57.
- Sheldrick, G. M. (2008). *Acta Cryst. A*, **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*, **65**, 148–155.
- Terzioglu, N., Karali, N., Gursoy, A., Pannecouque, C., Leysen, P., Paeshuyse, J., De Neyts, J. & Clercq, E. (2006). *Arkivoc*, pp. 109–118.

# supporting information

*Acta Cryst.* (2009). E65, o1343 [doi:10.1107/S1600536809017784]

## ***N,N'-(Phenyliminodimethylene)diprop-2-enamide hemihydrate***

**Dhanapal Tamilvendan, Ganesan Venkatesa Prabhu, Frank R. Fronczek and Nagarajan Vembu**

### **S1. Comment**

The Mannich reaction is a three-component condensation in which a compound containing an active H atom (substrate) is allowed to react with an aldehyde or ketone and a primary or secondary amine with concomitant release of water to produce a new base known as a Mannich base, in which the active hydrogen is replaced by an aminomethyl group. The formation of both carbon-carbon and carbon-nitrogen bond in this aminomethylation process makes the Mannich reaction an extremely useful synthetic transformation. Mannich bases have wide application in the areas of pharmaceuticals (Friedrich *et al.*, 1991) and macromolecular chemistry (Bohme & Mannich, 1955; Afsah *et al.*, 2008). Some Mannich bases have antimalarial, antiviral (Terzioglu *et al.*, 2006) properties while some other act as antihistamines, anti-inflammatories (Ravichandran *et al.*, 2007) and antimicrobials (Pandeya *et al.*, 2000). The present investigation is aimed at the elucidation of the molecular and crystal structure of the title compound which was obtained by the Mannich condensation of aniline, formaldehyde and acrylamide.

The asymmetric unit of (I) consists of *N,N*-[(phenylimino)dimethanediyl]bisprop-2-enamide and half a water molecule (Fig. 1).


The crystal structure of (I) is stabilized by O—H···O, N—H···O and C—H···O interactions. The range of H···O distances (Table 1) found in (I) agrees with those found for O—H···O & N—H···O (Jeffrey, 1997) and C—H···O hydrogen bonds (Desiraju & Steiner, 1999). Each of N5—H5···O11 and C15—H15···O7 interactions generate a ring motif of graph set (Bernstein *et al.*, 1995; Etter, 1990), S(8). An S(5) motif is formed by C4—H4A···O7. The Ow—Hw···O7<sup>i</sup>, N1—H1···O11<sup>ii</sup> and N5—H5···O11 interactions together generate an extended two dimensional network along the base vectors, [0 1 0] & [1 0 0] and through the plane (0 0 - 1). The supramolecular architecture is completed by the interplay of two edge to face C—H···π interactions (Table 1).

### **S2. Experimental**


7.1 g (0.1 mol) acrylamide and 10 ml (0.1M) of formaldehyde were dissolved in minimum amount of distilled water and the contents were mixed well to get a clear solution. 10 ml (0.1M) of aniline was added to the mixture in small installments with stirring. After 48 hr colorless solid was obtained which was washed with ethanol and dried at 343 K. The resulting organic compound was recrystallized from hot ethanol to yield the diffraction quality crystals of the title compound.

### **S3. Refinement**

All H-atoms were located in difference maps and their positions and isotropic displacement parameters were freely refined.

**Figure 1**


The asymmetric unit of (I) with the atoms labelled and displacement ellipsoids depicted at the 50% probability level for all non-H atoms. H-atoms are drawn as spheres of arbitrary radius.

**Figure 2**

The molecular packing viewed along the  $b$ -axis. Dashed lines represent the O—H $\cdots$ O, N—H $\cdots$ O and C—H $\cdots$ O interactions within the lattice.

### *N,N'-(Phenyliminodimethylene)diprop-2-enamide hemihydrate*

#### Crystal data



$M_r = 268.31$

Orthorhombic,  $Pbcn$

Hall symbol: -P 2n 2ab

$a = 17.074 (2) \text{ \AA}$

$b = 9.8366 (15) \text{ \AA}$

$c = 16.316 (2) \text{ \AA}$

$V = 2740.3 (6) \text{ \AA}^3$

$Z = 8$

$F(000) = 1144$

$D_x = 1.301 \text{ Mg m}^{-3}$

Melting point: 398 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 34434 reflections

$\theta = 2.5\text{--}33.0^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 90\text{ K}$   
Fragment, colorless

$0.30 \times 0.23 \times 0.12\text{ mm}$

#### Data collection

Nonius KappaCCD  
diffractometer with an Oxford Cryosystems  
Cryostream cooler  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  and  $\varphi$  scans  
9467 measured reflections

5065 independent reflections  
3885 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$   
 $\theta_{\text{max}} = 33.0^\circ, \theta_{\text{min}} = 2.7^\circ$   
 $h = -26 \rightarrow 26$   
 $k = -15 \rightarrow 15$   
 $l = -24 \rightarrow 24$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.120$   
 $S = 1.02$   
5065 reflections  
249 parameters  
0 restraints  
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
Hydrogen site location: difference Fourier map  
All H-atom parameters refined  
 $w = 1/[\sigma^2(F_o^2) + (0.0597P)^2 + 0.7128P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.39\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.28\text{ e \AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.80007 (5)	0.89440 (8)	0.13232 (5)	0.01594 (16)
C2	0.88056 (6)	0.84769 (10)	0.12071 (6)	0.01569 (17)
N3	0.89921 (5)	0.80690 (8)	0.03825 (5)	0.01545 (16)
C4	0.91579 (6)	0.66584 (10)	0.02225 (6)	0.01665 (18)
N5	0.85082 (5)	0.59623 (8)	-0.01804 (5)	0.01671 (16)
C6	0.85921 (6)	0.52348 (9)	-0.08750 (6)	0.01635 (18)
O7	0.92364 (4)	0.50903 (8)	-0.12139 (5)	0.02232 (17)
C8	0.78566 (6)	0.46251 (11)	-0.11941 (6)	0.0210 (2)
C9	0.78569 (7)	0.37886 (13)	-0.18250 (7)	0.0267 (2)
C10	0.73876 (5)	0.80842 (9)	0.13486 (5)	0.01421 (17)
O11	0.74626 (4)	0.68455 (7)	0.12367 (4)	0.01761 (15)
C12	0.66234 (6)	0.87457 (10)	0.15303 (6)	0.01837 (19)
C13	0.59676 (6)	0.80534 (12)	0.16321 (7)	0.0247 (2)
C14	0.90493 (5)	0.90267 (10)	-0.02424 (6)	0.01502 (17)
C15	0.91823 (5)	0.86297 (11)	-0.10565 (6)	0.01785 (19)

C16	0.92620 (6)	0.96063 (12)	-0.16649 (6)	0.0226 (2)
C17	0.92139 (6)	1.09817 (12)	-0.14964 (7)	0.0249 (2)
C18	0.90668 (6)	1.13762 (11)	-0.06939 (7)	0.0229 (2)
C19	0.89752 (6)	1.04206 (10)	-0.00757 (7)	0.01868 (19)
OW	1.0000	0.37093 (12)	-0.2500	0.0356 (3)
H1	0.7899 (9)	0.9829 (15)	0.1371 (9)	0.028 (4)*
H2A	0.8901 (7)	0.7705 (14)	0.1559 (8)	0.018 (3)*
H2B	0.9140 (7)	0.9206 (14)	0.1393 (8)	0.015 (3)*
H4A	0.9616 (8)	0.6566 (13)	-0.0140 (8)	0.018 (3)*
H4B	0.9274 (7)	0.6225 (13)	0.0743 (8)	0.017 (3)*
H5	0.8046 (9)	0.6058 (14)	0.0045 (9)	0.027 (3)*
H8	0.7378 (9)	0.4864 (15)	-0.0914 (10)	0.035 (4)*
H9A	0.7377 (8)	0.3357 (15)	-0.2005 (9)	0.029 (4)*
H9B	0.8348 (9)	0.3542 (15)	-0.2119 (10)	0.036 (4)*
H12	0.6636 (9)	0.9714 (16)	0.1589 (10)	0.034 (4)*
H13A	0.5471 (8)	0.8504 (15)	0.1751 (8)	0.027 (3)*
H13B	0.5952 (10)	0.7040 (19)	0.1577 (11)	0.045 (5)*
H15	0.9212 (7)	0.7696 (15)	-0.1208 (8)	0.018 (3)*
H16	0.9353 (9)	0.9338 (15)	-0.2219 (9)	0.031 (4)*
H17	0.9294 (9)	1.1688 (16)	-0.1936 (10)	0.038 (4)*
H18	0.9027 (8)	1.2309 (16)	-0.0557 (9)	0.029 (4)*
H19	0.8871 (8)	1.0736 (14)	0.0458 (9)	0.022 (3)*
HW	1.0232 (10)	0.4232 (17)	-0.2832 (11)	0.048 (5)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0168 (4)	0.0121 (3)	0.0189 (4)	-0.0001 (3)	0.0009 (3)	-0.0017 (3)
C2	0.0163 (4)	0.0169 (4)	0.0138 (4)	0.0001 (3)	-0.0014 (3)	-0.0017 (3)
N3	0.0195 (4)	0.0131 (3)	0.0137 (3)	0.0013 (3)	0.0007 (3)	-0.0011 (3)
C4	0.0184 (4)	0.0141 (4)	0.0175 (4)	0.0024 (3)	-0.0011 (3)	-0.0010 (3)
N5	0.0166 (4)	0.0167 (4)	0.0168 (4)	-0.0006 (3)	0.0038 (3)	-0.0035 (3)
C6	0.0181 (4)	0.0153 (4)	0.0157 (4)	0.0014 (3)	0.0026 (3)	-0.0010 (3)
O7	0.0176 (3)	0.0259 (4)	0.0235 (4)	0.0004 (3)	0.0054 (3)	-0.0080 (3)
C8	0.0177 (4)	0.0243 (5)	0.0210 (5)	-0.0002 (4)	0.0027 (3)	-0.0048 (4)
C9	0.0219 (5)	0.0343 (6)	0.0241 (5)	0.0014 (4)	-0.0020 (4)	-0.0094 (4)
C10	0.0172 (4)	0.0133 (4)	0.0122 (4)	-0.0002 (3)	-0.0010 (3)	0.0001 (3)
O11	0.0213 (3)	0.0118 (3)	0.0197 (3)	0.0002 (2)	0.0004 (3)	-0.0009 (2)
C12	0.0191 (4)	0.0148 (4)	0.0212 (5)	0.0011 (3)	0.0008 (3)	-0.0012 (3)
C13	0.0204 (5)	0.0241 (5)	0.0294 (5)	-0.0018 (4)	0.0037 (4)	-0.0072 (4)
C14	0.0119 (4)	0.0163 (4)	0.0169 (4)	-0.0010 (3)	-0.0011 (3)	0.0011 (3)
C15	0.0154 (4)	0.0199 (5)	0.0182 (4)	0.0000 (3)	0.0031 (3)	0.0010 (4)
C16	0.0184 (4)	0.0305 (6)	0.0189 (5)	-0.0014 (4)	0.0038 (4)	0.0055 (4)
C17	0.0208 (5)	0.0264 (5)	0.0274 (5)	-0.0039 (4)	0.0000 (4)	0.0116 (4)
C18	0.0206 (5)	0.0177 (5)	0.0305 (6)	-0.0034 (3)	-0.0044 (4)	0.0054 (4)
C19	0.0180 (4)	0.0164 (4)	0.0216 (5)	-0.0016 (3)	-0.0032 (3)	-0.0002 (3)
OW	0.0567 (8)	0.0170 (5)	0.0330 (7)	0.000	0.0289 (6)	0.000

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

N1—C10	1.3465 (12)	C10—O11	1.2387 (11)
N1—C2	1.4613 (13)	C10—C12	1.4878 (14)
N1—H1	0.891 (15)	C12—C13	1.3210 (15)
C2—N3	1.4397 (12)	C12—H12	0.957 (16)
C2—H2A	0.966 (13)	C13—H13A	0.976 (14)
C2—H2B	0.966 (13)	C13—H13B	1.001 (19)
N3—C14	1.3915 (12)	C14—C15	1.4030 (14)
N3—C4	1.4400 (12)	C14—C19	1.4036 (14)
C4—N5	1.4599 (12)	C15—C16	1.3880 (14)
C4—H4A	0.984 (13)	C15—H15	0.953 (14)
C4—H4B	0.970 (13)	C16—C17	1.3830 (17)
N5—C6	1.3480 (12)	C16—H16	0.954 (15)
N5—H5	0.876 (15)	C17—C18	1.3886 (17)
C6—O7	1.2394 (12)	C17—H17	1.008 (16)
C6—C8	1.4859 (14)	C18—C19	1.3876 (15)
C8—C9	1.3178 (15)	C18—H18	0.947 (16)
C8—H8	0.966 (16)	C19—H19	0.942 (14)
C9—H9A	0.969 (14)	OW—HW	0.845 (17)
C9—H9B	0.996 (16)		
C10—N1—C2	122.52 (8)	H9A—C9—H9B	117.4 (12)
C10—N1—H1	117.3 (10)	O11—C10—N1	122.20 (9)
C2—N1—H1	120.2 (10)	O11—C10—C12	123.38 (9)
N3—C2—N1	114.64 (8)	N1—C10—C12	114.41 (8)
N3—C2—H2A	107.4 (8)	C13—C12—C10	122.88 (9)
N1—C2—H2A	109.2 (8)	C13—C12—H12	121.3 (9)
N3—C2—H2B	111.7 (8)	C10—C12—H12	115.8 (9)
N1—C2—H2B	106.4 (8)	C12—C13—H13A	121.8 (9)
H2A—C2—H2B	107.3 (11)	C12—C13—H13B	121.6 (10)
C14—N3—C2	120.77 (8)	H13A—C13—H13B	116.6 (13)
C14—N3—C4	120.38 (8)	N3—C14—C15	121.10 (9)
C2—N3—C4	118.78 (8)	N3—C14—C19	120.87 (9)
N3—C4—N5	112.59 (8)	C15—C14—C19	118.03 (9)
N3—C4—H4A	110.7 (7)	C16—C15—C14	120.02 (10)
N5—C4—H4A	106.8 (7)	C16—C15—H15	118.5 (8)
N3—C4—H4B	107.8 (8)	C14—C15—H15	121.5 (8)
N5—C4—H4B	110.0 (8)	C17—C16—C15	121.95 (10)
H4A—C4—H4B	108.8 (11)	C17—C16—H16	118.0 (9)
C6—N5—C4	123.15 (8)	C15—C16—H16	120.1 (9)
C6—N5—H5	120.4 (10)	C16—C17—C18	118.13 (10)
C4—N5—H5	116.4 (9)	C16—C17—H17	121.7 (9)
O7—C6—N5	122.03 (9)	C18—C17—H17	120.2 (9)
O7—C6—C8	123.19 (9)	C19—C18—C17	121.10 (10)
N5—C6—C8	114.78 (8)	C19—C18—H18	118.4 (9)
C9—C8—C6	121.70 (9)	C17—C18—H18	120.5 (9)
C9—C8—H8	121.4 (9)	C18—C19—C14	120.72 (10)

C6—C8—H8	116.9 (9)	C18—C19—H19	118.1 (8)
C8—C9—H9A	120.7 (9)	C14—C19—H19	121.2 (8)
C8—C9—H9B	121.9 (9)		
C10—N1—C2—N3	75.00 (12)	N1—C10—C12—C13	-175.46 (10)
N1—C2—N3—C14	70.15 (11)	C2—N3—C14—C15	-175.96 (8)
N1—C2—N3—C4	-112.95 (9)	C4—N3—C14—C15	7.19 (13)
C14—N3—C4—N5	-78.90 (10)	C2—N3—C14—C19	4.02 (13)
C2—N3—C4—N5	104.19 (10)	C4—N3—C14—C19	-172.83 (8)
N3—C4—N5—C6	127.89 (10)	N3—C14—C15—C16	-177.94 (9)
C4—N5—C6—O7	1.15 (15)	C19—C14—C15—C16	2.09 (13)
C4—N5—C6—C8	-179.03 (9)	C14—C15—C16—C17	-0.07 (15)
O7—C6—C8—C9	5.99 (17)	C15—C16—C17—C18	-1.19 (16)
N5—C6—C8—C9	-173.83 (11)	C16—C17—C18—C19	0.39 (15)
C2—N1—C10—O11	-3.45 (14)	C17—C18—C19—C14	1.67 (15)
C2—N1—C10—C12	175.89 (8)	N3—C14—C19—C18	177.14 (9)
O11—C10—C12—C13	3.88 (16)	C15—C14—C19—C18	-2.88 (14)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N5—H5···O11	0.876 (15)	2.318 (15)	3.0476 (11)	140.8 (12)
C4—H4A···O7	0.984 (13)	2.366 (13)	2.8089 (12)	106.5 (9)
C15—H15···O7	0.953 (14)	2.563 (14)	3.4922 (14)	165.1 (10)
OW—HW···O7 <sup>i</sup>	0.845 (17)	1.990 (17)	2.8193 (9)	166.7 (16)
N1—H1···O11 <sup>ii</sup>	0.891 (15)	2.089 (15)	2.9651 (11)	167.4 (14)
C2—H2B···Cg1 <sup>iii</sup>	0.966 (13)	3.178	3.874	130.40
C8—H8···Cg1 <sup>iv</sup>	0.966 (16)	2.571	3.444	150.57

Symmetry codes: (i)  $-x+2, y, -z-1/2$ ; (ii)  $-x+3/2, y+1/2, z$ ; (iii)  $-x+2, -y+2, -z$ ; (iv)  $-x+1/2, y-3/2, z$ .