

## 4,4'-(Anthracene-9,10-diyl)dibenzoic acid dimethylformamide disolvate

Hong Li,<sup>a\*</sup> Zhi-Qiang Wang,<sup>b</sup> Liu-Zhi Yang,<sup>a</sup> Yan-Qi Liu<sup>a</sup> and Duo-Bin Mao<sup>a</sup>

<sup>a</sup>School of Food and Biological Engineering, Zhengzhou University of Light Industry, Zhengzhou 450002, People's Republic of China, and <sup>b</sup>College of Chemistry and Chemical Engineering, Luoyang Normal University, Luoyang 471022, People's Republic of China

Correspondence e-mail: lihong@zzuli.edu.cn


Received 26 March 2009; accepted 22 April 2009

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in solvent or counterion;  $R$  factor = 0.053;  $wR$  factor = 0.162; data-to-parameter ratio = 13.4.

In the title compound,  $\text{C}_{28}\text{H}_{18}\text{O}_4 \cdot 2\text{C}_3\text{H}_7\text{NO}$ , the dihedral angle between the benzene rings and the anthracene system is  $74.05(12)^\circ$ . A crystallographic inversion centre is located in the middle of the anthracene unit. The dimethylformamide solvent molecules are partially disordered over two positions of approximately equal occupancy [0.529(6):0.471(6)]. Inter-molecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds with the major occupancy formamide O atom as acceptor result in the formation of 2:1 solvate-complex aggregates, which are alternately linked to shorter solvate units *via* weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  contacts generated from the rotational disorder of the formamide O atom (minor occupancy component). Weak  $\text{C}-\text{H}\cdots\pi$  interactions between the solvent molecules as the donor and the outer anthracene rings support these contacts in the crystal structure for both disorder components.

### Related literature

For the structure of 4-(2,5-dihexyloxyphenyl)benzoic acid and the syntheses of related compounds, see: Li *et al.* (2008). For palladium-catalysed Suzuki coupling reactions, see: Xu *et al.* (2006, 2008); Li *et al.* (2006) and literature cited therein.



### Experimental

#### Crystal data

$\text{C}_{28}\text{H}_{18}\text{O}_4 \cdot 2\text{C}_3\text{H}_7\text{NO}$   
 $M_r = 564.62$   
 Triclinic,  $P\bar{1}$   
 $a = 7.3692(15)$  Å  
 $b = 8.9981(18)$  Å  
 $c = 12.124(2)$  Å  
 $\alpha = 71.157(3)^\circ$   
 $\beta = 77.640(3)^\circ$

$\gamma = 79.754(3)^\circ$   
 $V = 738.0(3)$  Å<sup>3</sup>  
 $Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.23 \times 0.16 \times 0.06$  mm

#### Data collection

Bruker SMART APEX CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.980$ ,  $T_{\max} = 0.994$

5691 measured reflections  
 2721 independent reflections  
 1467 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$   
 $wR(F^2) = 0.162$   
 $S = 1.02$   
 2721 reflections

203 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.16$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.20$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O2}-\text{H2D}\cdots\text{O3}^i$	0.82	1.79	2.603 (4)	170
$\text{C5}-\text{H5}\cdots\text{O3}^i$	0.93	2.63	3.478 (5)	152
$\text{C16}-\text{H16A}\cdots\text{Cg1}$	0.96	2.91	3.485 (3)	120

Symmetry code: (i)  $-x, -y, -z + 1$ . Cg1 is the centroid of the anthracene ring C8, C9, C10, C12A-C14A.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

This work was supported by the Doctoral Foundation of Zhengzhou University of Light Industry (000420).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2166).

### References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Li, H., Wu, Y. J. & Yan, W. B. (2006). *J. Organomet. Chem.* **691**, 5688–5696.  
 Li, H., Zhang, L., Liu, Y.-Q., Mao, D.-B. & Zhang, W.-Y. (2008). *Acta Cryst. E* **64**, o2094.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.  
 Xu, C., Gong, J. F., Guo, T., Zhang, Y. H. & Wu, Y. J. (2008). *J. Mol. Catal. A Chem.* **279**, 69–76.  
 Xu, C., Gong, J. F., Yue, S. F., Zhu, Y. & Wu, Y. J. (2006). *Dalton Trans.* pp. 4730–4739.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1223 [ doi:10.1107/S1600536809014858 ]

## 4,4'-(Anthracene-9,10-diyl)dibenzoic acid dimethylformamide disolvate

H. Li, Z.-Q. Wang, L.-Z. Yang, Y.-Q. Liu and D.-B. Mao

### Comment

Cyclopalladated ferrocenylimine complexes with monophosphino ligands were successfully used as catalysts for Suzuki reactions (Xu *et al.* 2006; Li *et al.*, 2006; Xu *et al.* 2008). We have recently reported that the structure of 4-(2,5-dihexyloxyphenyl)benzoic acid was obtained from the Suzuki coupling reaction (Li *et al.*, 2008). The title compound was derived from the Suzuki reaction of 9,10-dibromoanthracene and 4-carboxyphenylboronic acid.

In the title compound (Fig.1), the dihedral angle between benzene rings and anthracene rings is  $74.05(12)^\circ$ . A crystallographic inversion centre is in the middle of the anthracene unit, and an approximate two-fold pseudo rotation axis is running along the plane of the anthracene unit. The dimethylformamide solvent molecules are partially disordered over two positions, O3 and O3', of approximately equal occupancy, (0.529 (6) and 0.471 (6), respectively). The different intermolecular hydrogen bonding contacts are shown in Fig. 1 (O3' is the acceptor) and Fig. 2 (with O3 as acceptor). The intermolecular O—H $\cdots$ O hydrogen bonds result in the formation of long 2:1 solvate:complex aggregates, (Table 1) which are alternately linked *via* weak intermolecular C—H $\cdots$ O contacts generated from the rotational disorder of the formamide oxygen atom (0.471 (6) site occupancy). C—H $\cdots$  $\pi$  interactions support these contacts in the crystal structure forming a one-dimensional supramolecular architecture (Fig. 1 and Fig. 2).

### Experimental

The title compound was obtained from the Suzuki coupling reaction of 9,10-dibromoanthracene and 4-carboxyphenylboronic acid as described in the literature (Li *et al.*, 2008) and recrystallized from dimethylformamide at room temperature to give the desired crystals suitable for single-crystal X-ray diffraction.

### Refinement

H atoms attached to C atoms of the title compound were placed in geometrically idealized positions and treated as riding with C—H distances constrained to 0.93 (aromatic CH), or 0.96 Å (methyl CH<sub>3</sub>), and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  (1.5 $U_{\text{eq}}$  for methyl H). The atom-site occupancies for the rotational disordered formamide oxygen atoms O3 and O3' refined to a ratio of 0.53/0.47.

Alert levels A and B for short intermolecular O1 $\cdots$ O3' and H2D $\cdots$ H15' contacts with distances of 2.50 Å and 2.01 Å may be explained by the difficulties to split the whole solvent molecule due to the pseudo two-fold rotation of the methyl groups around the N1—C15 axis. BUMP instruction or splitting of the whole solvent molecule resulted in unstable refinements. Introduction of shift-limiting restraints (DAMP instruction) resulted in larger R-values without improving the geometries. Therefore the partial disorder refinement (O3, O3', H15, H15') was preferred as a compromise.

## Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids at the 30% probability level (suffix A denotes the symmetry code:  $-x + 1, -y + 1, -z$ ). Weak C—H $\cdots$  $\pi$  and C—H $\cdots$ O hydrogen bonding contacts are indicated with dashed lines. Cg1 is the centroid of the anthracene ring C8, C9, C10, C12A, C13A, C14A.




Fig. 2. Partial view of the crystal packing showing the intermolecular O—H $\cdots$ O hydrogen bonds and weak C—H $\cdots$  $\pi$  interactions. Cg1 is the centroid of the anthracene ring (C8, C9, C10, C12A, C13A, C14A).

## 4,4'-(Anthracene-9,10-diyl)dibenzoic acid dimethylformamide disolvate

### Crystal data

$C_{28}H_{18}O_4 \cdot 2C_3H_7NO$

$M_r = 564.62$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 7.3692\ (15)\ \text{\AA}$

$b = 8.9981\ (18)\ \text{\AA}$

$c = 12.124\ (2)\ \text{\AA}$

$\alpha = 71.157\ (3)^\circ$

$\beta = 77.640\ (3)^\circ$

$\gamma = 79.754\ (3)^\circ$

$V = 738.0\ (3)\ \text{\AA}^3$

$Z = 1$

$F_{000} = 298$

$D_x = 1.270\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 879 reflections

$\theta = 2.9\text{--}22.0^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 295\ \text{K}$

Block, colourless

$0.23 \times 0.16 \times 0.06\ \text{mm}$

### Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295\ \text{K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.980, T_{\max} = 0.994$

5691 measured reflections

2721 independent reflections

1467 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 25.5^\circ$

$\theta_{\min} = 2.4^\circ$

$h = -8 \rightarrow 8$

$k = -10 \rightarrow 9$

$l = -14 \rightarrow 14$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.053$$

$$wR(F^2) = 0.162$$

$$S = 1.02$$

2721 reflections

203 parameters

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0732P)^2 + 0.04P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C15	0.5707 (5)	0.2790 (4)	0.4860 (3)	0.0742 (9)	0.529 (6)
H15	0.4771	0.2595	0.4536	0.089*	0.529 (6)
O3	0.5840 (5)	0.2039 (5)	0.5938 (4)	0.0918 (18)	0.529 (6)
C15'	0.5707 (5)	0.2790 (4)	0.4860 (3)	0.0742 (9)	0.471 (6)
H15'	0.5944	0.2141	0.5598	0.089*	0.471 (6)
O3'	0.4309 (7)	0.2700 (6)	0.4479 (4)	0.096 (2)	0.471 (6)
N1	0.6843 (3)	0.3826 (3)	0.41928 (19)	0.0637 (6)	
C16	0.6620 (5)	0.4705 (4)	0.2994 (3)	0.0999 (12)	
H16A	0.5498	0.4476	0.2831	0.150*	
H16B	0.7677	0.4407	0.2459	0.150*	
H16C	0.6535	0.5816	0.2895	0.150*	
C17	0.8377 (4)	0.4104 (5)	0.4627 (3)	0.0945 (11)	
H17A	0.8375	0.3450	0.5429	0.142*	
H17B	0.8249	0.5196	0.4598	0.142*	
H17C	0.9532	0.3852	0.4146	0.142*	
O1	-0.1924 (3)	-0.0937 (3)	0.42504 (19)	0.1024 (9)	
O2	-0.3377 (3)	-0.0181 (3)	0.2730 (2)	0.1090 (9)	
H2D	-0.4140	-0.0728	0.3213	0.163*	
C1	-0.0616 (3)	0.0915 (3)	0.2520 (2)	0.0496 (6)	
C2	-0.0651 (4)	0.1757 (3)	0.1348 (2)	0.0638 (8)	
H2	-0.1589	0.1662	0.0978	0.077*	
C3	0.0713 (4)	0.2746 (3)	0.0719 (2)	0.0594 (7)	
H3	0.0669	0.3313	-0.0068	0.071*	
C4	0.2133 (3)	0.2901 (3)	0.1243 (2)	0.0445 (6)	

## supplementary materials

C5	0.2169 (3)	0.2037 (3)	0.2414 (2)	0.0525 (7)
H5	0.3117	0.2117	0.2783	0.063*
C6	0.0805 (3)	0.1053 (3)	0.3045 (2)	0.0537 (7)
H6	0.0852	0.0480	0.3831	0.064*
C7	-0.2053 (4)	-0.0146 (3)	0.3219 (3)	0.0643 (8)
C8	0.6573 (4)	0.1754 (3)	-0.1369 (2)	0.0640 (8)
H8	0.6611	0.0821	-0.1552	0.077*
C9	0.5159 (4)	0.2148 (3)	-0.0568 (2)	0.0549 (7)
H9	0.4239	0.1477	-0.0204	0.066*
C10	0.5040 (3)	0.3575 (3)	-0.02629 (19)	0.0432 (6)
C11	0.3592 (3)	0.3985 (3)	0.05850 (19)	0.0419 (6)
C12	0.3533 (3)	0.5398 (3)	0.08512 (19)	0.0419 (6)
C13	0.2061 (3)	0.5896 (3)	0.1681 (2)	0.0514 (7)
H13	0.1110	0.5259	0.2059	0.062*
C14	0.2008 (4)	0.7263 (3)	0.1932 (2)	0.0620 (8)
H14	0.1031	0.7554	0.2478	0.074*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C15	0.066 (2)	0.089 (2)	0.066 (2)	-0.0197 (18)	-0.0072 (17)	-0.0164 (19)
O3	0.082 (3)	0.114 (4)	0.071 (3)	-0.054 (3)	-0.012 (2)	0.006 (3)
C15'	0.066 (2)	0.089 (2)	0.066 (2)	-0.0197 (18)	-0.0072 (17)	-0.0164 (19)
O3'	0.087 (4)	0.120 (4)	0.079 (3)	-0.060 (3)	-0.015 (3)	0.000 (3)
N1	0.0574 (14)	0.0753 (17)	0.0547 (14)	-0.0202 (12)	-0.0040 (11)	-0.0113 (12)
C16	0.088 (2)	0.119 (3)	0.071 (2)	-0.022 (2)	-0.0099 (18)	0.004 (2)
C17	0.079 (2)	0.125 (3)	0.086 (2)	-0.045 (2)	-0.0063 (19)	-0.028 (2)
O1	0.0933 (16)	0.128 (2)	0.0722 (15)	-0.0639 (15)	-0.0176 (13)	0.0188 (14)
O2	0.0826 (16)	0.137 (2)	0.0938 (17)	-0.0672 (15)	-0.0245 (14)	0.0188 (15)
C1	0.0454 (14)	0.0470 (15)	0.0548 (16)	-0.0134 (12)	-0.0041 (12)	-0.0114 (12)
C2	0.0535 (16)	0.0701 (19)	0.0652 (18)	-0.0227 (14)	-0.0159 (14)	-0.0045 (15)
C3	0.0605 (16)	0.0624 (18)	0.0512 (15)	-0.0233 (14)	-0.0133 (13)	-0.0001 (13)
C4	0.0460 (14)	0.0408 (14)	0.0468 (14)	-0.0098 (11)	-0.0061 (11)	-0.0115 (12)
C5	0.0541 (15)	0.0553 (16)	0.0474 (15)	-0.0189 (13)	-0.0085 (12)	-0.0077 (13)
C6	0.0593 (16)	0.0519 (16)	0.0453 (14)	-0.0170 (13)	-0.0041 (12)	-0.0058 (12)
C7	0.0506 (17)	0.0634 (19)	0.074 (2)	-0.0191 (14)	-0.0030 (15)	-0.0121 (16)
C8	0.085 (2)	0.0442 (16)	0.0605 (17)	-0.0144 (14)	0.0043 (15)	-0.0192 (14)
C9	0.0667 (17)	0.0400 (15)	0.0549 (16)	-0.0186 (12)	0.0012 (13)	-0.0107 (12)
C10	0.0494 (14)	0.0359 (14)	0.0416 (13)	-0.0110 (11)	-0.0061 (11)	-0.0054 (11)
C11	0.0453 (13)	0.0386 (14)	0.0391 (13)	-0.0119 (10)	-0.0083 (11)	-0.0034 (11)
C12	0.0426 (13)	0.0404 (14)	0.0404 (13)	-0.0080 (10)	-0.0063 (10)	-0.0074 (11)
C13	0.0485 (14)	0.0494 (16)	0.0499 (15)	-0.0125 (12)	0.0041 (12)	-0.0104 (12)
C14	0.0724 (18)	0.0492 (17)	0.0573 (17)	-0.0080 (14)	0.0079 (14)	-0.0178 (14)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C15—O3	1.279 (4)	C3—H3	0.9300
C15—N1	1.314 (4)	C4—C5	1.385 (3)
C15—H15	0.9300	C4—C11	1.499 (3)

N1—C17	1.434 (4)	C5—C6	1.389 (3)
N1—C16	1.441 (3)	C5—H5	0.9300
C16—H16A	0.9600	C6—H6	0.9300
C16—H16B	0.9600	C8—C9	1.346 (3)
C16—H16C	0.9600	C8—C14 <sup>i</sup>	1.406 (4)
C17—H17A	0.9600	C8—H8	0.9300
C17—H17B	0.9600	C9—C10	1.430 (3)
C17—H17C	0.9600	C9—H9	0.9300
O1—C7	1.238 (3)	C10—C11	1.403 (3)
O2—C7	1.255 (3)	C10—C12 <sup>i</sup>	1.438 (3)
O2—H2D	0.8200	C11—C12	1.401 (3)
C1—C6	1.378 (3)	C12—C13	1.428 (3)
C1—C2	1.381 (3)	C12—C10 <sup>i</sup>	1.438 (3)
C1—C7	1.485 (3)	C13—C14	1.353 (3)
C2—C3	1.392 (3)	C13—H13	0.9300
C2—H2	0.9300	C14—C8 <sup>i</sup>	1.406 (4)
C3—C4	1.382 (3)	C14—H14	0.9300
O3—C15—N1	122.5 (4)	C4—C5—C6	120.7 (2)
O3—C15—H15	118.8	C4—C5—H5	119.6
N1—C15—H15	118.8	C6—C5—H5	119.6
C15—N1—C17	121.0 (3)	C1—C6—C5	120.8 (2)
C15—N1—C16	121.5 (3)	C1—C6—H6	119.6
C17—N1—C16	117.5 (2)	C5—C6—H6	119.6
N1—C16—H16A	109.5	O1—C7—O2	122.7 (3)
N1—C16—H16B	109.5	O1—C7—C1	119.7 (3)
H16A—C16—H16B	109.5	O2—C7—C1	117.6 (3)
N1—C16—H16C	109.5	C9—C8—C14 <sup>i</sup>	120.8 (3)
H16A—C16—H16C	109.5	C9—C8—H8	119.6
H16B—C16—H16C	109.5	C14 <sup>i</sup> —C8—H8	119.6
N1—C17—H17A	109.5	C8—C9—C10	121.5 (2)
N1—C17—H17B	109.5	C8—C9—H9	119.2
H17A—C17—H17B	109.5	C10—C9—H9	119.2
N1—C17—H17C	109.5	C11—C10—C9	122.0 (2)
H17A—C17—H17C	109.5	C11—C10—C12 <sup>i</sup>	119.9 (2)
H17B—C17—H17C	109.5	C9—C10—C12 <sup>i</sup>	118.1 (2)
C7—O2—H2D	109.5	C12—C11—C10	119.9 (2)
C6—C1—C2	118.9 (2)	C12—C11—C4	119.3 (2)
C6—C1—C7	119.4 (2)	C10—C11—C4	120.8 (2)
C2—C1—C7	121.6 (2)	C11—C12—C13	122.3 (2)
C1—C2—C3	120.2 (2)	C11—C12—C10 <sup>i</sup>	120.2 (2)
C1—C2—H2	119.9	C13—C12—C10 <sup>i</sup>	117.5 (2)
C3—C2—H2	119.9	C14—C13—C12	122.0 (2)
C4—C3—C2	121.2 (2)	C14—C13—H13	119.0
C4—C3—H3	119.4	C12—C13—H13	119.0
C2—C3—H3	119.4	C13—C14—C8 <sup>i</sup>	120.0 (2)
C3—C4—C5	118.1 (2)	C13—C14—H14	120.0

## supplementary materials

---

C3—C4—C11	121.9 (2)	C8 <sup>i</sup> —C14—H14	120.0
C5—C4—C11	119.9 (2)		
O3—C15—N1—C17	3.6 (5)	C8—C9—C10—C11	-178.9 (2)
O3—C15—N1—C16	-177.4 (4)	C8—C9—C10—C12 <sup>i</sup>	0.9 (4)
C6—C1—C2—C3	-1.1 (4)	C9—C10—C11—C12	-179.8 (2)
C7—C1—C2—C3	179.7 (2)	C12 <sup>i</sup> —C10—C11—C12	0.4 (4)
C1—C2—C3—C4	0.5 (4)	C9—C10—C11—C4	2.2 (3)
C2—C3—C4—C5	0.3 (4)	C12 <sup>i</sup> —C10—C11—C4	-177.6 (2)
C2—C3—C4—C11	-178.8 (2)	C3—C4—C11—C12	106.8 (3)
C3—C4—C5—C6	-0.5 (4)	C5—C4—C11—C12	-72.3 (3)
C11—C4—C5—C6	178.6 (2)	C3—C4—C11—C10	-75.2 (3)
C2—C1—C6—C5	0.9 (4)	C5—C4—C11—C10	105.7 (3)
C7—C1—C6—C5	-179.9 (2)	C10—C11—C12—C13	178.1 (2)
C4—C5—C6—C1	-0.1 (4)	C4—C11—C12—C13	-3.9 (3)
C6—C1—C7—O1	-2.6 (4)	C10—C11—C12—C10 <sup>i</sup>	-0.4 (4)
C2—C1—C7—O1	176.6 (3)	C4—C11—C12—C10 <sup>i</sup>	177.6 (2)
C6—C1—C7—O2	176.5 (3)	C11—C12—C13—C14	-179.8 (2)
C2—C1—C7—O2	-4.3 (4)	C10 <sup>i</sup> —C12—C13—C14	-1.3 (4)
C14 <sup>i</sup> —C8—C9—C10	0.3 (4)	C12—C13—C14—C8 <sup>i</sup>	0.1 (4)

Symmetry codes: (i)  $-x+1, -y+1, -z$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2D <sup>ii</sup> —O3 <sup>ii</sup>	0.82	1.79	2.603 (4)	170
C5—H5 <sup>iii</sup> —O3'	0.93	2.63	3.478 (5)	152
C16—H16A <sup>iii</sup> —Cg1	0.96	2.91	3.485 (3)	120

Symmetry codes: (ii)  $-x, -y, -z+1$ .

Fig. 1




Fig. 2

