

Bis[μ -1,3-bis[(benzimidazol-1-yl)methyl]benzene- $\kappa^2N^3:N^3'$]bis[dichloridozinc(II)] dimethylformamide disolvate

Li-Zhen Zhao,^a Ping Li,^a Bao-Liang Cao^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, Jining Normal College, Wulanchabu, Inner Mongolia 012000, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 27 April 2009; accepted 28 April 2009

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.037; wR factor = 0.097; data-to-parameter ratio = 18.5.

In the title compound, $[\text{Zn}_2\text{Cl}_4(\text{C}_{22}\text{H}_{18}\text{N}_4)_2] \cdot 2\text{C}_3\text{H}_7\text{NO}$, the 1,3-bis[(benzimidazol-1-yl)methyl]benzene ligand bridges two ZnCl_2 units, forming a centrosymmetric dinuclear molecule. The Zn^{II} atom shows a distorted tetrahedral coordination within a Cl_2N_2 donor set.

Related literature

For the crystal structure of 1,3-bis((benzimidazol-1-yl)methyl)benzene, which was isolated as the malonic acid co-crystal, see: Aakeröy *et al.* (2005). For related metal complexes, see: Fan *et al.* (2006); Raehm *et al.* (2003).



Experimental

Crystal data

$[\text{Zn}_2\text{Cl}_4(\text{C}_{22}\text{H}_{18}\text{N}_4)_2] \cdot 2\text{C}_3\text{H}_7\text{NO}$
 $M_r = 1095.54$
Monoclinic, $C2/c$
 $a = 24.0069$ (5) Å
 $b = 9.8217$ (2) Å
 $c = 23.9723$ (5) Å
 $\beta = 117.695$ (1)°

$V = 5004.8$ (2) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.22$ mm⁻¹
 $T = 120$ K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan *SADABS* (Sheldrick, 1996)
 $T_{\text{min}} = 0.711$, $T_{\text{max}} = 0.888$

17166 measured reflections
5730 independent reflections
4449 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.097$
 $S = 1.04$
5730 reflections

309 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.92$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.57$ e Å⁻³

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2009).

We acknowledge support from the Scientific Research Projects of Higher Education of Inner Mongolia (NJzy08217) and the University of Malaya.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2441).

References

- Aakeröy, C. B., Desper, J., Elisabeth, E., Helfrich, B. A., Levin, B. & Urbina, J. F. (2005). *Z. Kristallogr.* **200**, 325–332.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Fan, J., Yee, G. T., Wang, G. & Hanson, B. E. (2006). *Inorg. Chem.* **45**, 599–608.
Raehm, L., Mimassi, L., Guyard-Duhayon, C. & Amouri, H. (2003). *Inorg. Chem.* **42**, 5654–5659.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2009). *pubCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m613 [doi:10.1107/S1600536809015943]

Bis{ μ -1,3-bis[(benzimidazol-1-yl)methyl]benzene- $\kappa^2N^3:N^{3'}$ }bis[dichloridozinc(II)] dimethylformamide disolvate

L.-Z. Zhao, P. Li, B.-L. Cao and S. W. Ng

Comment

(type here to add)

Experimental

The compound was prepared from a mixture of boric acid (0.17 g), zinc chloride (0.27 g), 1,3-bis((benzimidazol-1-yl)methyl)benzene (0.45 g) in DMF (3.6 ml) and water (0.2 ml). The mixture was sealed in 25-ml Teflon-lined stainless-steel vessel, which was heated at 423 K for 5 days. The vessel was then cooled to room temperature slowly. Crystals were picked out manually.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with $U(H)$ fixed at $1.2U_{eq}(C)$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $[ZnCl_2(C_{22}H_{18}N_4)]_2 \cdot 2DMF$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Only one DMF molecule is shown. Unlabelled non-H atoms are related by a centre of inversion.

Bis{ μ -1,3-bis[(benzimidazol-1-yl)methyl]benzene- $\kappa^2N^3:N^{3'}$ }bis[dichloridozinc(II)] dimethylformamide disolvate

Crystal data

$[Zn_2Cl_4(C_{22}H_{18}N_4)_2] \cdot 2C_3H_7NO$

$M_r = 1095.54$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 24.0069$ (5) Å

$F_{000} = 2256$

$D_x = 1.454$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 4135 reflections

$\theta = 2.3$ – 27.2°

supplementary materials

$b = 9.8217(2) \text{ \AA}$
 $c = 23.9723(5) \text{ \AA}$
 $\beta = 117.695(1)^\circ$
 $V = 5004.8(2) \text{ \AA}^3$
 $Z = 4$

$\mu = 1.22 \text{ mm}^{-1}$
 $T = 120 \text{ K}$
Prism, colorless
 $0.30 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 120 \text{ K}$
 ω scans
Absorption correction: Multi-scan SADABS (Sheldrick, 1996)
 $T_{\min} = 0.711$, $T_{\max} = 0.888$
17166 measured reflections

5730 independent reflections
4449 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\text{max}} = 27.5^\circ$
 $\theta_{\text{min}} = 1.9^\circ$
 $h = -30 \rightarrow 31$
 $k = -12 \rightarrow 12$
 $l = -31 \rightarrow 28$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.097$
 $S = 1.04$
5730 reflections
309 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0452P)^2 + 5.1375P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.92 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.57 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.499493 (12)	0.24704 (3)	0.615569 (12)	0.01976 (9)
Cl1	0.54858 (3)	0.40106 (6)	0.69289 (3)	0.02422 (14)
Cl2	0.54751 (3)	0.05513 (6)	0.61286 (3)	0.03540 (17)
O1	0.28873 (14)	0.6931 (3)	0.61655 (11)	0.0738 (8)
N1	0.46559 (9)	0.32960 (19)	0.52860 (9)	0.0194 (4)
N2	0.42352 (9)	0.33960 (19)	0.42428 (9)	0.0183 (4)
N3	0.16359 (9)	0.2139 (2)	0.36682 (9)	0.0223 (4)
N4	0.08242 (9)	0.2789 (2)	0.38075 (9)	0.0224 (4)
N5	0.37081 (12)	0.8246 (2)	0.67983 (10)	0.0369 (6)
C1	0.43461 (10)	0.4540 (2)	0.50863 (10)	0.0185 (5)
C2	0.42823 (11)	0.5612 (2)	0.54354 (11)	0.0224 (5)
H2	0.4464	0.5574	0.5882	0.027*

C3	0.39418 (11)	0.6732 (2)	0.51009 (12)	0.0263 (5)
H3	0.3887	0.7479	0.5322	0.032*
C4	0.36741 (11)	0.6791 (2)	0.44394 (12)	0.0260 (5)
H4	0.3444	0.7579	0.4227	0.031*
C5	0.37366 (11)	0.5743 (2)	0.40925 (11)	0.0232 (5)
H5	0.3556	0.5785	0.3646	0.028*
C6	0.40796 (10)	0.4615 (2)	0.44313 (11)	0.0199 (5)
C7	0.45751 (11)	0.2659 (2)	0.47682 (11)	0.0202 (5)
H7	0.4738	0.1779	0.4766	0.024*
C8	0.40337 (11)	0.2955 (2)	0.35930 (11)	0.0206 (5)
H8A	0.4220	0.2054	0.3595	0.025*
H8B	0.4187	0.3611	0.3382	0.025*
C9	0.33254 (11)	0.2861 (2)	0.32330 (11)	0.0209 (5)
C10	0.29914 (11)	0.2108 (2)	0.34747 (11)	0.0208 (5)
H10	0.3215	0.1614	0.3856	0.025*
C11	0.23407 (11)	0.2072 (2)	0.31670 (11)	0.0222 (5)
C12	0.20141 (12)	0.2768 (3)	0.25988 (12)	0.0285 (6)
H12	0.1567	0.2742	0.2382	0.034*
C13	0.23399 (13)	0.3496 (3)	0.23511 (12)	0.0338 (6)
H13	0.2117	0.3959	0.1961	0.041*
C14	0.29951 (12)	0.3554 (3)	0.26713 (11)	0.0291 (6)
H14	0.3216	0.4073	0.2503	0.035*
C15	0.19950 (12)	0.1274 (2)	0.34467 (12)	0.0257 (5)
H15A	0.1702	0.0633	0.3126	0.031*
H15B	0.2303	0.0730	0.3805	0.031*
C16	0.10264 (11)	0.1968 (2)	0.35076 (11)	0.0223 (5)
H16	0.0766	0.1318	0.3206	0.027*
C17	0.13518 (11)	0.3561 (2)	0.42061 (11)	0.0221 (5)
C18	0.14072 (12)	0.4591 (2)	0.46287 (11)	0.0251 (5)
H18	0.1058	0.4870	0.4686	0.030*
C19	0.19923 (12)	0.5187 (3)	0.49612 (12)	0.0285 (6)
H19	0.2047	0.5892	0.5254	0.034*
C20	0.25048 (12)	0.4777 (3)	0.48768 (12)	0.0300 (6)
H20	0.2899	0.5214	0.5114	0.036*
C21	0.24572 (12)	0.3754 (3)	0.44588 (12)	0.0283 (6)
H21	0.2808	0.3474	0.4405	0.034*
C22	0.18653 (11)	0.3158 (2)	0.41212 (11)	0.0227 (5)
C23	0.31081 (17)	0.7967 (4)	0.64606 (15)	0.0501 (9)
H23	0.2819	0.8643	0.6446	0.060*
C24	0.4155 (2)	0.7252 (4)	0.6803 (2)	0.0749 (12)
H24A	0.3929	0.6512	0.6510	0.112*
H24B	0.4396	0.6883	0.7229	0.112*
H24C	0.4443	0.7688	0.6672	0.112*
C25	0.3954 (2)	0.9448 (4)	0.71704 (16)	0.0667 (12)
H25A	0.3605	1.0025	0.7133	0.100*
H25B	0.4211	0.9949	0.7020	0.100*
H25C	0.4213	0.9192	0.7612	0.100*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.01891 (15)	0.01906 (14)	0.01959 (14)	0.00064 (11)	0.00751 (11)	0.00179 (10)
C11	0.0220 (3)	0.0268 (3)	0.0201 (3)	-0.0046 (2)	0.0067 (2)	-0.0007 (2)
C12	0.0442 (4)	0.0243 (3)	0.0351 (3)	0.0130 (3)	0.0162 (3)	0.0053 (3)
O1	0.093 (2)	0.0474 (15)	0.0492 (14)	-0.0216 (15)	0.0061 (14)	-0.0035 (12)
N1	0.0187 (10)	0.0179 (10)	0.0211 (9)	0.0017 (8)	0.0088 (8)	0.0031 (8)
N2	0.0178 (10)	0.0180 (9)	0.0194 (9)	0.0005 (8)	0.0089 (8)	0.0005 (7)
N3	0.0229 (11)	0.0223 (10)	0.0239 (10)	-0.0048 (8)	0.0126 (9)	-0.0058 (8)
N4	0.0228 (10)	0.0218 (10)	0.0236 (10)	-0.0057 (8)	0.0116 (9)	-0.0045 (8)
N5	0.0455 (15)	0.0325 (13)	0.0262 (11)	-0.0073 (11)	0.0112 (11)	-0.0024 (10)
C1	0.0144 (11)	0.0178 (11)	0.0231 (11)	-0.0012 (8)	0.0085 (9)	0.0003 (9)
C2	0.0172 (12)	0.0240 (12)	0.0237 (12)	-0.0005 (9)	0.0078 (10)	-0.0013 (9)
C3	0.0247 (13)	0.0203 (12)	0.0344 (13)	0.0034 (10)	0.0143 (11)	-0.0034 (10)
C4	0.0224 (13)	0.0211 (12)	0.0328 (13)	0.0068 (10)	0.0116 (11)	0.0061 (10)
C5	0.0207 (12)	0.0224 (12)	0.0249 (12)	0.0035 (9)	0.0092 (10)	0.0037 (9)
C6	0.0160 (11)	0.0202 (12)	0.0238 (12)	-0.0021 (9)	0.0096 (9)	0.0001 (9)
C7	0.0168 (11)	0.0186 (11)	0.0248 (12)	0.0005 (9)	0.0094 (9)	0.0031 (9)
C8	0.0229 (12)	0.0219 (11)	0.0199 (11)	0.0008 (9)	0.0123 (10)	0.0003 (9)
C9	0.0247 (13)	0.0202 (11)	0.0203 (11)	-0.0013 (9)	0.0128 (10)	-0.0025 (9)
C10	0.0243 (12)	0.0180 (11)	0.0198 (11)	0.0005 (9)	0.0099 (10)	-0.0002 (9)
C11	0.0266 (13)	0.0183 (11)	0.0236 (12)	-0.0034 (9)	0.0132 (10)	-0.0067 (9)
C12	0.0236 (13)	0.0315 (14)	0.0244 (12)	-0.0035 (10)	0.0062 (10)	-0.0048 (10)
C13	0.0312 (15)	0.0420 (16)	0.0202 (12)	-0.0028 (12)	0.0051 (11)	0.0071 (11)
C14	0.0290 (14)	0.0343 (14)	0.0240 (12)	-0.0048 (11)	0.0124 (11)	0.0049 (11)
C15	0.0268 (13)	0.0235 (13)	0.0307 (13)	-0.0033 (10)	0.0166 (11)	-0.0057 (10)
C16	0.0218 (12)	0.0229 (12)	0.0227 (12)	-0.0063 (9)	0.0106 (10)	-0.0034 (9)
C17	0.0220 (12)	0.0212 (12)	0.0210 (11)	-0.0034 (9)	0.0082 (10)	0.0000 (9)
C18	0.0274 (13)	0.0227 (12)	0.0264 (12)	-0.0009 (10)	0.0135 (11)	-0.0037 (10)
C19	0.0316 (14)	0.0248 (13)	0.0251 (13)	-0.0028 (11)	0.0100 (11)	-0.0058 (10)
C20	0.0250 (13)	0.0295 (14)	0.0311 (14)	-0.0088 (11)	0.0093 (11)	-0.0077 (11)
C21	0.0235 (13)	0.0293 (14)	0.0323 (13)	-0.0060 (10)	0.0130 (11)	-0.0050 (11)
C22	0.0256 (13)	0.0206 (12)	0.0225 (12)	-0.0053 (10)	0.0117 (10)	-0.0033 (9)
C23	0.053 (2)	0.0425 (18)	0.0376 (17)	-0.0103 (16)	0.0071 (15)	0.0080 (14)
C24	0.075 (3)	0.078 (3)	0.083 (3)	0.007 (2)	0.047 (3)	-0.009 (2)
C25	0.094 (3)	0.043 (2)	0.0422 (19)	-0.0191 (19)	0.014 (2)	-0.0115 (15)

Geometric parameters (\AA , $^\circ$)

Zn1—N1	2.022 (2)	C8—H8B	0.9900
Zn1—N4 ⁱ	2.025 (2)	C9—C14	1.383 (3)
Zn1—C11	2.2544 (6)	C9—C10	1.399 (3)
Zn1—C12	2.2262 (7)	C10—C11	1.384 (3)
O1—C23	1.211 (4)	C10—H10	0.9500
N1—C7	1.321 (3)	C11—C12	1.394 (4)
N1—C1	1.394 (3)	C11—C15	1.506 (3)

N2—C7	1.349 (3)	C12—C13	1.381 (4)
N2—C6	1.391 (3)	C12—H12	0.9500
N2—C8	1.465 (3)	C13—C14	1.394 (4)
N3—C16	1.339 (3)	C13—H13	0.9500
N3—C22	1.389 (3)	C14—H14	0.9500
N3—C15	1.474 (3)	C15—H15A	0.9900
N4—C16	1.314 (3)	C15—H15B	0.9900
N4—C17	1.403 (3)	C16—H16	0.9500
N4—Zn1 ⁱ	2.025 (2)	C17—C18	1.393 (3)
N5—C23	1.312 (4)	C17—C22	1.397 (3)
N5—C25	1.431 (4)	C18—C19	1.382 (3)
N5—C24	1.447 (5)	C18—H18	0.9500
C1—C6	1.395 (3)	C19—C20	1.396 (4)
C1—C2	1.397 (3)	C19—H19	0.9500
C2—C3	1.382 (3)	C20—C21	1.386 (4)
C2—H2	0.9500	C20—H20	0.9500
C3—C4	1.408 (3)	C21—C22	1.396 (3)
C3—H3	0.9500	C21—H21	0.9500
C4—C5	1.375 (3)	C23—H23	0.9500
C4—H4	0.9500	C24—H24A	0.9800
C5—C6	1.394 (3)	C24—H24B	0.9800
C5—H5	0.9500	C24—H24C	0.9800
C7—H7	0.9500	C25—H25A	0.9800
C8—C9	1.510 (3)	C25—H25B	0.9800
C8—H8A	0.9900	C25—H25C	0.9800
N1—Zn1—N4 ⁱ	99.30 (8)	C10—C11—C12	119.4 (2)
N1—Zn1—C11	112.62 (6)	C10—C11—C15	119.7 (2)
N1—Zn1—C12	105.97 (6)	C12—C11—C15	120.9 (2)
N4 ⁱ —Zn1—C11	101.28 (6)	C13—C12—C11	120.0 (2)
N4 ⁱ —Zn1—C12	114.92 (6)	C13—C12—H12	120.0
C11—Zn1—C12	120.81 (3)	C11—C12—H12	120.0
C7—N1—C1	105.68 (18)	C12—C13—C14	120.2 (2)
C7—N1—Zn1	126.28 (15)	C12—C13—H13	119.9
C1—N1—Zn1	126.89 (15)	C14—C13—H13	119.9
C7—N2—C6	107.21 (19)	C9—C14—C13	120.4 (2)
C7—N2—C8	126.25 (19)	C9—C14—H14	119.8
C6—N2—C8	126.44 (19)	C13—C14—H14	119.8
C16—N3—C22	107.3 (2)	N3—C15—C11	113.3 (2)
C16—N3—C15	124.6 (2)	N3—C15—H15A	108.9
C22—N3—C15	127.7 (2)	C11—C15—H15A	108.9
C16—N4—C17	105.0 (2)	N3—C15—H15B	108.9
C16—N4—Zn1 ⁱ	124.06 (16)	C11—C15—H15B	108.9
C17—N4—Zn1 ⁱ	128.75 (16)	H15A—C15—H15B	107.7
C23—N5—C25	124.9 (3)	N4—C16—N3	113.5 (2)
C23—N5—C24	117.5 (3)	N4—C16—H16	123.3
C25—N5—C24	117.6 (3)	N3—C16—H16	123.3
N1—C1—C6	109.0 (2)	C18—C17—C22	121.3 (2)

supplementary materials

N1—C1—C2	130.1 (2)	C18—C17—N4	129.7 (2)
C6—C1—C2	120.9 (2)	C22—C17—N4	109.0 (2)
C3—C2—C1	116.9 (2)	C19—C18—C17	117.1 (2)
C3—C2—H2	121.6	C19—C18—H18	121.5
C1—C2—H2	121.6	C17—C18—H18	121.5
C2—C3—C4	121.6 (2)	C18—C19—C20	121.5 (2)
C2—C3—H3	119.2	C18—C19—H19	119.3
C4—C3—H3	119.2	C20—C19—H19	119.3
C5—C4—C3	121.9 (2)	C21—C20—C19	122.1 (2)
C5—C4—H4	119.0	C21—C20—H20	118.9
C3—C4—H4	119.0	C19—C20—H20	118.9
C4—C5—C6	116.3 (2)	C20—C21—C22	116.3 (2)
C4—C5—H5	121.8	C20—C21—H21	121.9
C6—C5—H5	121.8	C22—C21—H21	121.9
N2—C6—C5	132.1 (2)	N3—C22—C21	133.0 (2)
N2—C6—C1	105.54 (19)	N3—C22—C17	105.2 (2)
C5—C6—C1	122.4 (2)	C21—C22—C17	121.8 (2)
N1—C7—N2	112.6 (2)	O1—C23—N5	126.3 (4)
N1—C7—H7	123.7	O1—C23—H23	116.8
N2—C7—H7	123.7	N5—C23—H23	116.8
N2—C8—C9	110.68 (19)	N5—C24—H24A	109.5
N2—C8—H8A	109.5	N5—C24—H24B	109.5
C9—C8—H8A	109.5	H24A—C24—H24B	109.5
N2—C8—H8B	109.5	N5—C24—H24C	109.5
C9—C8—H8B	109.5	H24A—C24—H24C	109.5
H8A—C8—H8B	108.1	H24B—C24—H24C	109.5
C14—C9—C10	118.9 (2)	N5—C25—H25A	109.5
C14—C9—C8	120.7 (2)	N5—C25—H25B	109.5
C10—C9—C8	120.4 (2)	H25A—C25—H25B	109.5
C11—C10—C9	121.0 (2)	N5—C25—H25C	109.5
C11—C10—H10	119.5	H25A—C25—H25C	109.5
C9—C10—H10	119.5	H25B—C25—H25C	109.5
N4 ⁱ —Zn1—N1—C7	-101.91 (19)	C9—C10—C11—C15	-178.8 (2)
C12—Zn1—N1—C7	17.5 (2)	C10—C11—C12—C13	-0.6 (4)
C11—Zn1—N1—C7	151.65 (17)	C15—C11—C12—C13	179.9 (2)
N4 ⁱ —Zn1—N1—C1	64.00 (19)	C11—C12—C13—C14	-0.9 (4)
C12—Zn1—N1—C1	-176.59 (17)	C10—C9—C14—C13	-0.2 (4)
C11—Zn1—N1—C1	-42.44 (19)	C8—C9—C14—C13	-178.1 (2)
C7—N1—C1—C6	-0.1 (2)	C12—C13—C14—C9	1.3 (4)
Zn1—N1—C1—C6	-168.31 (15)	C16—N3—C15—C11	128.1 (2)
C7—N1—C1—C2	-179.8 (2)	C22—N3—C15—C11	-59.4 (3)
Zn1—N1—C1—C2	12.0 (3)	C10—C11—C15—N3	112.2 (2)
N1—C1—C2—C3	-179.9 (2)	C12—C11—C15—N3	-68.3 (3)
C6—C1—C2—C3	0.5 (3)	C17—N4—C16—N3	-0.5 (3)
C1—C2—C3—C4	-0.3 (4)	Zn1 ⁱ —N4—C16—N3	-164.84 (16)
C2—C3—C4—C5	0.0 (4)	C22—N3—C16—N4	0.6 (3)
C3—C4—C5—C6	0.0 (4)	C15—N3—C16—N4	174.3 (2)
C7—N2—C6—C5	-179.8 (2)	C16—N4—C17—C18	179.7 (2)

C8—N2—C6—C5	-3.4 (4)	Zn1 ⁱ —N4—C17—C18	-17.1 (4)
C7—N2—C6—C1	0.0 (2)	C16—N4—C17—C22	0.3 (3)
C8—N2—C6—C1	176.4 (2)	Zn1 ⁱ —N4—C17—C22	163.61 (17)
C4—C5—C6—N2	180.0 (2)	C22—C17—C18—C19	-0.3 (4)
C4—C5—C6—C1	0.1 (3)	N4—C17—C18—C19	-179.6 (2)
N1—C1—C6—N2	0.0 (2)	C17—C18—C19—C20	0.0 (4)
C2—C1—C6—N2	179.7 (2)	C18—C19—C20—C21	-0.1 (4)
N1—C1—C6—C5	179.9 (2)	C19—C20—C21—C22	0.4 (4)
C2—C1—C6—C5	-0.4 (4)	C16—N3—C22—C21	179.6 (3)
C1—N1—C7—N2	0.1 (3)	C15—N3—C22—C21	6.1 (4)
Zn1—N1—C7—N2	168.44 (15)	C16—N3—C22—C17	-0.3 (3)
C6—N2—C7—N1	-0.1 (3)	C15—N3—C22—C17	-173.8 (2)
C8—N2—C7—N1	-176.5 (2)	C20—C21—C22—N3	179.4 (3)
C7—N2—C8—C9	115.4 (2)	C20—C21—C22—C17	-0.7 (4)
C6—N2—C8—C9	-60.3 (3)	C18—C17—C22—N3	-179.4 (2)
N2—C8—C9—C14	125.4 (2)	N4—C17—C22—N3	0.0 (3)
N2—C8—C9—C10	-52.4 (3)	C18—C17—C22—C21	0.7 (4)
C14—C9—C10—C11	-1.3 (4)	N4—C17—C22—C21	-179.9 (2)
C8—C9—C10—C11	176.5 (2)	C25—N5—C23—O1	-176.3 (3)
C9—C10—C11—C12	1.7 (3)	C24—N5—C23—O1	2.7 (5)

Symmetry codes: (i) $-x+1/2, -y+1/2, -z+1$.

Fig. 1

