

## 5-Chloro-8-hydroxyquinolinium nitrate

Seik Weng Ng

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 25 May 2009; accepted 26 May 2009

Key indicators: single-crystal X-ray study;  $T = 123$  K; mean  $\sigma(C-C) = 0.005$  Å;  
R factor = 0.062; wR factor = 0.192; data-to-parameter ratio = 14.4.

The 5-chloro-8-hydroxyquinolinium cation in the title ion pair,  $C_9H_7ClNO^+ \cdot NO_3^-$ , is approximately coplanar with the nitrate anion [dihedral angle = 16.1 (1)°]. Two ion pairs are hydrogen bonded ( $2 \times O-H \cdots O$  and  $2 \times N-H \cdots O$ ) about a center of inversion, generating an  $R_4^4(14)$  ring.

### Related literature

The 8-hydroxyquinolinium cation has been isolated as a number of salts; for the 8-hydroxyquinolinium chloride hydrate, see: Skakle *et al.* (2006). For the crystal structure of 5-chloro-8-hydroxyquinoline, see: Banerjee & Saha (1986).



### Experimental

#### Crystal data



$M_r = 242.62$

### Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{min} = 0.925$ ,  $T_{max} = 0.980$

6472 measured reflections  
2196 independent reflections  
1574 reflections with  $I > 2\sigma$   
 $R_{int} = 0.049$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.062$   
 $wR(F^2) = 0.192$   
 $S = 1.07$   
2196 reflections  
153 parameters  
2 restraints

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\text{max}} = 1.20$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.34$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1—H1o···O2 <sup>i</sup>	0.84 (1)	1.87 (1)	2.695 (3)	169 (4)
N1—H1n···O2	0.88 (1)	1.95 (1)	2.816 (3)	167 (4)

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

I thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2465).

### References

- Banerjee, T. & Saha, N. N. (1986). *Acta Cryst.* C42, 1408–1411.
- Barbour, L. J. (2001). *J. Supramol. Chem.* 1, 189–191.
- Bruker (2008). APEX2 and SAINT Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.
- Skakle, J. M. S., Wardell, J. L. & Wardell, S. M. S. V. (2006). *Acta Cryst.* C62, o312–o314.
- Westrip, S. P. (2009). publCIF. In preparation.

# supporting information

*Acta Cryst.* (2009). E65, o1450 [doi:10.1107/S160053680901993X]

## 5-Chloro-8-hydroxyquinolinium nitrate

Seik Weng Ng


### S1. Experimental

Zinc acetate (0.19 g, 1 mmol) and 5-chloro-8-hydroxyquinoline (0.36 g, 2 mmol) were loaded into a convection tube; the tube was filled with dry methanol and kept at 333 K. Yellow crystals were collected from the side-arm after several days.

### S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 Å) and were included in the refinement in the riding model approximation with  $U(H)$  fixed at 1.2 $U(C)$ . The ammonium and hydroxy H-atoms were located in a difference Fourier map, and were refined with distance restraints of N–H = 0.88±0.01 Å and O–H = 0.84±0.01 Å; their isotropic temperature factors were refined.

The final difference Fourier map had a large peak at about 1 Å from the C11 atom.



**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of a pair of hydrogen-bonded  $[C_9H_7ClNO][NO_3]$  ion pairs drawn at the 70% probability level. Hydrogen atoms are drawn spheres of arbitrary radius and dashed lines denote hydrogen bonds.

## 5-Chloro-8-hydroxyquinolinium nitrate

### Crystal data

$C_9H_7ClNO^+ \cdot NO_3^-$   
 $M_r = 242.62$

Monoclinic,  $P2_1/n$   
Hall symbol: -P 2yn

$a = 7.4379 (3)$  Å  
 $b = 11.5518 (6)$  Å  
 $c = 11.2288 (5)$  Å  
 $\beta = 95.831 (3)^\circ$   
 $V = 959.80 (8)$  Å<sup>3</sup>  
 $Z = 4$   
 $F(000) = 496$   
 $D_x = 1.679$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 1079 reflections  
 $\theta = 2.5\text{--}26.4^\circ$   
 $\mu = 0.40$  mm<sup>-1</sup>  
 $T = 123$  K  
Prism, yellow  
 $0.20 \times 0.05 \times 0.05$  mm

*Data collection*

Bruker SMART APEX  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.925$ ,  $T_{\max} = 0.980$

6472 measured reflections  
2196 independent reflections  
1574 reflections with  $I > 2\sigma I$   
 $R_{\text{int}} = 0.049$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.5^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -15 \rightarrow 12$   
 $l = -14 \rightarrow 14$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.062$   
 $wR(F^2) = 0.192$   
 $S = 1.07$   
2196 reflections  
153 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.1164P)^2 + 0.1797P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.20$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.34$  e Å<sup>-3</sup>

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	-0.30656 (11)	0.30338 (8)	0.76985 (8)	0.0321 (3)
O1	0.3429 (3)	0.4706 (2)	0.5632 (2)	0.0279 (6)
H1O	0.350 (6)	0.5423 (11)	0.553 (4)	0.042 (12)*
O2	0.5861 (3)	0.30211 (18)	0.4592 (2)	0.0261 (6)
O3	0.6434 (3)	0.12125 (19)	0.4947 (2)	0.0325 (6)
O4	0.8190 (3)	0.2201 (2)	0.3904 (2)	0.0286 (6)
N1	0.2829 (4)	0.2424 (2)	0.5774 (2)	0.0203 (6)
H1N	0.385 (3)	0.265 (4)	0.552 (3)	0.037 (11)*
N2	0.6849 (4)	0.2125 (2)	0.4472 (2)	0.0220 (6)
C1	0.2635 (4)	0.1291 (3)	0.5806 (3)	0.0250 (7)
H1A	0.3516	0.0806	0.5504	0.030*
C2	0.1155 (5)	0.0794 (3)	0.6278 (3)	0.0277 (7)
H2	0.1012	-0.0023	0.6295	0.033*
C3	-0.0094 (4)	0.1513 (3)	0.6720 (3)	0.0265 (7)
H3	-0.1093	0.1185	0.7063	0.032*
C4	0.0084 (4)	0.2727 (3)	0.6672 (3)	0.0213 (7)
C5	-0.1144 (4)	0.3518 (3)	0.7089 (3)	0.0254 (7)

C6	-0.0849 (5)	0.4687 (3)	0.7003 (3)	0.0325 (8)
H6	-0.1694	0.5213	0.7282	0.039*
C7	0.0666 (5)	0.5118 (3)	0.6513 (3)	0.0297 (8)
H7	0.0832	0.5931	0.6460	0.036*
C8	0.1920 (4)	0.4385 (3)	0.6108 (3)	0.0231 (7)
C9	0.1613 (4)	0.3178 (3)	0.6184 (3)	0.0204 (7)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0219 (5)	0.0405 (6)	0.0359 (5)	-0.0012 (3)	0.0128 (3)	-0.0003 (4)
O1	0.0275 (13)	0.0173 (12)	0.0413 (14)	0.0001 (9)	0.0147 (10)	0.0025 (10)
O2	0.0276 (13)	0.0138 (11)	0.0388 (13)	0.0032 (9)	0.0124 (10)	-0.0010 (9)
O3	0.0337 (14)	0.0145 (11)	0.0513 (16)	-0.0003 (9)	0.0135 (12)	0.0044 (10)
O4	0.0270 (13)	0.0265 (13)	0.0342 (13)	0.0027 (9)	0.0121 (10)	-0.0009 (10)
N1	0.0173 (14)	0.0205 (13)	0.0235 (14)	0.0010 (10)	0.0046 (11)	0.0003 (10)
N2	0.0214 (14)	0.0175 (13)	0.0274 (14)	0.0005 (10)	0.0038 (11)	-0.0025 (10)
C1	0.0241 (17)	0.0199 (16)	0.0320 (18)	0.0020 (12)	0.0073 (13)	-0.0003 (13)
C2	0.0278 (17)	0.0205 (16)	0.0356 (18)	-0.0003 (13)	0.0066 (14)	0.0058 (14)
C3	0.0227 (17)	0.0272 (17)	0.0300 (17)	-0.0079 (13)	0.0043 (13)	0.0027 (13)
C4	0.0208 (16)	0.0253 (16)	0.0180 (15)	-0.0034 (12)	0.0034 (12)	0.0000 (12)
C5	0.0182 (16)	0.0337 (19)	0.0252 (16)	-0.0003 (13)	0.0074 (12)	0.0025 (13)
C6	0.0271 (18)	0.0279 (18)	0.045 (2)	0.0074 (14)	0.0137 (15)	-0.0051 (15)
C7	0.0291 (19)	0.0210 (17)	0.0405 (19)	0.0021 (13)	0.0111 (15)	-0.0024 (14)
C8	0.0240 (16)	0.0201 (16)	0.0258 (16)	-0.0012 (12)	0.0058 (12)	0.0006 (12)
C9	0.0239 (17)	0.0171 (15)	0.0203 (15)	0.0021 (12)	0.0021 (12)	0.0009 (11)

*Geometric parameters ( $\text{\AA}$ ,  $\text{^\circ}$ )*

C11—C5	1.739 (3)	C2—H2	0.9500
O1—C8	1.343 (4)	C3—C4	1.410 (5)
O1—H1O	0.84 (1)	C3—H3	0.9500
O2—N2	1.285 (3)	C4—C5	1.405 (4)
O3—N2	1.234 (3)	C4—C9	1.411 (4)
O4—N2	1.240 (4)	C5—C6	1.374 (5)
N1—C1	1.318 (4)	C6—C7	1.395 (5)
N1—C9	1.368 (4)	C6—H6	0.9500
N1—H1N	0.88 (1)	C7—C8	1.371 (4)
C1—C2	1.393 (5)	C7—H7	0.9500
C1—H1A	0.9500	C8—C9	1.417 (4)
C2—C3	1.375 (5)		
C8—O1—H1O	113 (3)	C5—C4—C3	124.6 (3)
C1—N1—C9	123.0 (3)	C9—C4—C3	117.7 (3)
C1—N1—H1N	114 (3)	C6—C5—C4	120.1 (3)
C9—N1—H1N	123 (3)	C6—C5—Cl1	119.2 (3)
O3—N2—O4	122.1 (3)	C4—C5—Cl1	120.7 (3)
O3—N2—O2	118.2 (3)	C5—C6—C7	121.3 (3)

O4—N2—O2	119.7 (3)	C5—C6—H6	119.3
N1—C1—C2	120.8 (3)	C7—C6—H6	119.3
N1—C1—H1A	119.6	C8—C7—C6	120.9 (3)
C2—C1—H1A	119.6	C8—C7—H7	119.5
C3—C2—C1	118.5 (3)	C6—C7—H7	119.5
C3—C2—H2	120.7	O1—C8—C7	125.8 (3)
C1—C2—H2	120.7	O1—C8—C9	116.3 (3)
C2—C3—C4	121.2 (3)	C7—C8—C9	118.0 (3)
C2—C3—H3	119.4	N1—C9—C4	118.8 (3)
C4—C3—H3	119.4	N1—C9—C8	119.3 (3)
C5—C4—C9	117.8 (3)	C4—C9—C8	121.9 (3)
C9—N1—C1—C2	-0.5 (5)	C6—C7—C8—O1	-179.4 (3)
N1—C1—C2—C3	-0.5 (5)	C6—C7—C8—C9	1.0 (5)
C1—C2—C3—C4	1.6 (5)	C1—N1—C9—C4	0.5 (4)
C2—C3—C4—C5	179.2 (3)	C1—N1—C9—C8	-179.6 (3)
C2—C3—C4—C9	-1.7 (5)	C5—C4—C9—N1	179.8 (3)
C9—C4—C5—C6	0.8 (5)	C3—C4—C9—N1	0.6 (4)
C3—C4—C5—C6	179.9 (3)	C5—C4—C9—C8	-0.2 (4)
C9—C4—C5—Cl1	179.8 (2)	C3—C4—C9—C8	-179.4 (3)
C3—C4—C5—Cl1	-1.1 (5)	O1—C8—C9—N1	-0.4 (4)
C4—C5—C6—C7	-0.6 (5)	C7—C8—C9—N1	179.3 (3)
Cl1—C5—C6—C7	-179.5 (3)	O1—C8—C9—C4	179.6 (3)
C5—C6—C7—C8	-0.4 (6)	C7—C8—C9—C4	-0.7 (5)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1o···O2 <sup>i</sup>	0.84 (1)	1.87 (1)	2.695 (3)	169 (4)
N1—H1n···O2	0.88 (1)	1.95 (1)	2.816 (3)	167 (4)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .