

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**(Dimethylformamide- $\kappa$ O)[4,4',6,6'-tetrabromo-2,2'-[*o*-phenylenebis(nitrilomethylidene)]diphenolato- $\kappa^4$ O,N,N',O']-copper(II) dimethylformamide solvate**

Yu Wu,\* Bin Xie, Li-Ke Zou, Wei-Ping Wu and Lu Lu

College of Chemistry and Pharmaceutical Engineering, Sichuan University of Science and Engineering, Zigong, Sichuan 643000, People's Republic of China

Correspondence e-mail: [solicitor.wu@126.com](mailto:solicitor.wu@126.com)


Received 30 May 2009; accepted 7 June 2009

Key indicators: single-crystal X-ray study;  $T = 93$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å; disorder in solvent or counterion;  $R$  factor = 0.035;  $wR$  factor = 0.063; data-to-parameter ratio = 16.1.

In the title compound,  $[\text{Cu}(\text{C}_{20}\text{H}_{10}\text{Br}_4\text{N}_2\text{O}_2)(\text{C}_3\text{H}_7\text{NO})] \cdot \text{C}_3\text{H}_7\text{NO}$ , the  $\text{Cu}^{\text{II}}$  ion is coordinated by two N atoms and two O atoms from a tetradentate Schiff base ligand and the O atom of one dimethylformamide ligand in an almost square-pyramidal geometry. The uncoordinated dimethylformamide solvent molecule is disordered over two sets of positions with occupancies of 0.741 (4) and 0.259 (4). The crystal packing is stabilized by  $\text{C}-\text{H} \cdots \text{O}$  interactions.

## Related literature

For the preparation of 3,5-dibromosalicylaldehyde, see: Elzbieta *et al.* (1964). For a related structure, see: Bei *et al.* (2003).



## Experimental

## Crystal data

$[\text{Cu}(\text{C}_{20}\text{H}_{10}\text{Br}_4\text{N}_2\text{O}_2)(\text{C}_3\text{H}_7\text{NO})] \cdot \text{C}_3\text{H}_7\text{NO}$   
 $M_r = 839.67$

Triclinic,  $P\bar{1}$   
 $a = 7.3742$  (10) Å  
 $b = 11.9542$  (19) Å

$c = 17.212$  (2) Å  
 $\alpha = 94.207$  (9)°  
 $\beta = 100.310$  (6)°  
 $\gamma = 104.117$  (5)°  
 $V = 1436.8$  (3) Å<sup>3</sup>

$Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 6.36$  mm<sup>-1</sup>  
 $T = 93$  K  
 $0.25 \times 0.25 \times 0.25$  mm

## Data collection

Rigaku SPIDER diffractometer  
 Absorption correction: multi-scan  
 (*RAPID-AUTO*; Rigaku, 2004)  
 $T_{\text{min}} = 0.299$ ,  $T_{\text{max}} = 0.299$   
 (expected range = 0.204–0.204)

11906 measured reflections  
 6330 independent reflections  
 5305 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.027$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$   
 $wR(F^2) = 0.063$   
 $S = 1.02$   
 6330 reflections  
 392 parameters

8 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.75$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.52$  e Å<sup>-3</sup>

Table 1

Selected bond lengths (Å).

Cu1—O1	1.910 (2)	Cu1—N2	1.958 (2)
Cu1—O2	1.9174 (19)	Cu1—O3	2.501 (2)
Cu1—N1	1.954 (2)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C5}-\text{H5} \cdots \text{O3}^{\text{i}}$	0.95	2.47	3.319 (4)	148
$\text{C7}-\text{H7} \cdots \text{O3}^{\text{i}}$	0.95	2.37	3.243 (4)	153
$\text{C14}-\text{H14} \cdots \text{O4}^{\text{ii}}$	0.95	2.33	3.226 (14)	156
$\text{C22}-\text{H22A} \cdots \text{O1}$	0.98	2.54	3.378 (4)	143

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x + 2, -y + 1, -z + 2$ .

Data collection: *RAPID-AUTO* (Rigaku, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge financial assistance from Sichuan University of Science and Engineering (grant No. 2007 ZR005), the Education Committee of Sichuan Province of China (grant Nos. 2006 A110 and 07ZA161) and the Science and Technology Office of Zigong City, China (grant No. 08X01).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2819).

## References

- Bei, F.-L., Ma, W.-X. & Wang, J. (2003). *Chin. J. Inorg. Chem.* **19**, 609–612.  
 Elzbieta, C., Zygmunt, E. & Romuald, K. (1964). *Diss. Pharm.* **15**, 369–378.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Rigaku (2004). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2009). E65, m758 [ doi:10.1107/S160053680902162X ]

**(Dimethylformamide- $\kappa O$ ){4,4',6,6'-tetrabromo-2,2'-[*o*-phenylenebis(nitrilomethylidyne)]diphenolato- $\kappa^4 O,N,N',O'$ }copper(II) dimethylformamide solvate**

**Y. Wu, B. Xie, L.-K. Zou, W.-P. Wu and L. Lu**

### Comment

The crystal structure and some properties of 1,2-*N,N'*-disalicylidenephenylaminato-nickel(II) was previously reported by Bei *et al.* (2003). We report here the synthesis and crystal structure of the title complex, [Cu(C<sub>14</sub>H<sub>8</sub>N<sub>2</sub>O<sub>3</sub>Br<sub>2</sub>)(C<sub>3</sub>H<sub>7</sub>NO)](C<sub>3</sub>H<sub>7</sub>NO).

The contents of the asymmetric unit are shown in Fig.1. The Cu<sup>II</sup> ion is coordinated by two N atoms and two O atoms from one 4,4',6,6'-tetrabromo-2,2'-[1,2-phenylenebis(nitrilomethylidyne)]diphenolate dianion and one O atom of *N,N*-dimethylformamide ligand in a square-pyramidal geometry (Table 1). The crystal packing is stabilized by C—H $\cdots$ O interactions (Table 2).

### Experimental

The title complex was synthesized in two stages. In the first stage, 3,5- dibromosalicylaldehyde was prepared according to Elzbieta *et al.* (1964). Two mole equivalents of 3,5-dibromosalicylaldehyde in ethanol (50 ml) was slowly added to 1,2-phenylenediamine (6 g) in ethanol (100 ml) with continuous stirring. The Schiff base molecule, *viz.* 4,4',6,6'-tetrabromo-2,2'-[1,2-phenylenebis(nitrilomethylidyne)] diphenol, precipitated immediately. In the second stage, the ligand (1 mmol), Cu(OAc)<sub>2</sub> (1 mmol) and DMF (25 ml) were refluxed for 1 h. The hot solution was filtered and allowed to stand at room temperature undisturbed for about three weeks, resulting in dark green crystals.

### Refinement

The uncoordinated *N,N*-dimethylformamide solvent molecule is disordered over two positions with occupancies of 0.741 (4) and 0.259 (4). The N—C(*sp*<sup>3</sup>), N—C(*sp*<sup>2</sup>) and C—O distances in both disorder components were restrained to 1.460 (3), 1.340 (3) and 1.220 (3) Å, respectively. H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 or 0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. A view of the title complex, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for the sake of clarity. Both disorder components of the uncoordinated solvent molecule are shown.

## (Dimethylformamide- $\kappa$ O){4,4',6,6'-tetrabromo-2,2'-[o-phenylenebis(nitrilomethylidyne)]diphenolato- $\kappa^4$ O,N,N',O'}copper(II) dimethylformamide solvate

### Crystal data

[Cu(C <sub>20</sub> H <sub>10</sub> Br <sub>4</sub> N <sub>2</sub> O <sub>2</sub> )(C <sub>3</sub> H <sub>7</sub> NO)]·C <sub>3</sub> H <sub>7</sub> NO	$Z = 2$
$M_r = 839.67$	$F_{000} = 818$
Triclinic, $P\bar{1}$	$D_x = 1.941 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.3742 (10) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 11.9542 (19) \text{ \AA}$	Cell parameters from 4531 reflections
$c = 17.212 (2) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$\alpha = 94.207 (9)^\circ$	$\mu = 6.36 \text{ mm}^{-1}$
$\beta = 100.310 (6)^\circ$	$T = 93 \text{ K}$
$\gamma = 104.117 (5)^\circ$	Block, dark green
$V = 1436.8 (3) \text{ \AA}^3$	$0.25 \times 0.25 \times 0.25 \text{ mm}$

### Data collection

Rigaku SPIDER diffractometer	6330 independent reflections
Radiation source: Rotating Anode	5305 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 93 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
$\omega$ scans	$\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (RAPID-AUTO; Rigaku, 2004)	$h = -8 \rightarrow 9$
$T_{\text{min}} = 0.299$ , $T_{\text{max}} = 0.299$	$k = -15 \rightarrow 15$
11906 measured reflections	$l = -22 \rightarrow 22$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H-atom parameters constrained
$wR(F^2) = 0.063$	$w = 1/[\sigma^2(F_o^2) + (0.0247P)^2]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
6330 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
392 parameters	$\Delta\rho_{\text{max}} = 0.75 \text{ e \AA}^{-3}$
8 restraints	$\Delta\rho_{\text{min}} = -0.52 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.39299 (5)	0.60816 (3)	0.67052 (2)	0.01305 (9)	
Br1	0.12215 (5)	0.88993 (3)	0.535887 (19)	0.01998 (8)	
Br2	-0.00681 (5)	0.60574 (3)	0.244696 (18)	0.02312 (9)	
Br3	0.74637 (5)	0.75127 (3)	1.105720 (18)	0.02390 (9)	
Br4	0.49378 (5)	0.96915 (3)	0.842406 (19)	0.02212 (9)	
O1	0.2620 (3)	0.69347 (17)	0.60103 (11)	0.0141 (5)	
O2	0.4409 (3)	0.73265 (18)	0.75456 (11)	0.0152 (5)	
N1	0.3449 (3)	0.4761 (2)	0.58884 (14)	0.0124 (5)	
N2	0.4835 (4)	0.5038 (2)	0.74155 (14)	0.0142 (6)	
C1	0.2062 (4)	0.6699 (3)	0.52455 (17)	0.0129 (6)	
C2	0.1319 (4)	0.7497 (3)	0.48000 (17)	0.0137 (6)	
C3	0.0698 (4)	0.7320 (3)	0.39909 (17)	0.0150 (7)	
H3	0.0224	0.7885	0.3718	0.018*	
C4	0.0777 (4)	0.6295 (3)	0.35772 (17)	0.0161 (7)	
C5	0.1450 (4)	0.5472 (3)	0.39550 (17)	0.0157 (7)	
H5	0.1484	0.4777	0.3660	0.019*	
C6	0.2099 (4)	0.5664 (3)	0.47917 (17)	0.0125 (6)	
C7	0.2743 (4)	0.4742 (3)	0.51432 (18)	0.0142 (6)	
H7	0.2642	0.4061	0.4799	0.017*	
C8	0.3973 (4)	0.3788 (3)	0.61884 (18)	0.0134 (6)	
C9	0.3827 (4)	0.2740 (3)	0.57325 (19)	0.0172 (7)	
H9	0.3361	0.2643	0.5174	0.021*	
C10	0.4360 (4)	0.1849 (3)	0.60964 (19)	0.0191 (7)	
H10	0.4272	0.1139	0.5788	0.023*	
C11	0.5026 (5)	0.1987 (3)	0.69131 (19)	0.0217 (7)	
H11	0.5367	0.1361	0.7159	0.026*	
C12	0.5201 (4)	0.3013 (3)	0.73727 (19)	0.0184 (7)	
H12	0.5667	0.3097	0.7931	0.022*	
C13	0.4690 (4)	0.3924 (3)	0.70124 (18)	0.0139 (6)	
C14	0.5440 (4)	0.5275 (3)	0.81769 (17)	0.0149 (7)	
H14	0.5803	0.4682	0.8457	0.018*	
C15	0.5606 (4)	0.6358 (3)	0.86300 (17)	0.0147 (7)	
C16	0.6296 (4)	0.6425 (3)	0.94604 (17)	0.0168 (7)	

## supplementary materials

H16	0.6600	0.5771	0.9680	0.020*	
C17	0.6526 (4)	0.7431 (3)	0.99485 (17)	0.0172 (7)	
C18	0.6111 (4)	0.8407 (3)	0.96363 (18)	0.0174 (7)	
H18	0.6280	0.9104	0.9976	0.021*	
C19	0.5458 (4)	0.8344 (3)	0.88351 (18)	0.0156 (7)	
C20	0.5124 (4)	0.7327 (3)	0.82895 (17)	0.0142 (7)	
O3	0.7162 (3)	0.69177 (19)	0.64181 (13)	0.0218 (5)	
N3	0.8292 (4)	0.8852 (2)	0.68559 (16)	0.0251 (7)	
C21	0.8417 (5)	0.7771 (3)	0.6803 (2)	0.0272 (8)	
H21	0.9556	0.7628	0.7081	0.033*	
C22	0.6627 (5)	0.9150 (3)	0.6398 (2)	0.0278 (8)	
H22A	0.5753	0.8447	0.6082	0.033*	
H22B	0.7052	0.9719	0.6043	0.033*	
H22C	0.5967	0.9483	0.6765	0.033*	
C23	0.9791 (6)	0.9809 (3)	0.7305 (3)	0.0478 (12)	
H23A	1.0806	0.9512	0.7597	0.057*	
H23B	0.9273	1.0233	0.7682	0.057*	
H23C	1.0315	1.0333	0.6941	0.057*	
O4	1.225 (3)	0.6119 (8)	1.0649 (3)	0.0317 (16)	0.741 (4)
C24	1.1892 (7)	0.7023 (4)	1.0473 (2)	0.0281 (12)	0.741 (4)
H24	1.2090	0.7632	1.0891	0.034*	0.741 (4)
N4	1.1242 (16)	0.7216 (4)	0.9731 (2)	0.0216 (12)	0.741 (4)
C25	1.0952 (8)	0.8347 (4)	0.9571 (4)	0.0382 (14)	0.741 (4)
H25A	1.1095	0.8828	1.0075	0.046*	0.741 (4)
H25B	0.9665	0.8243	0.9254	0.046*	0.741 (4)
H25C	1.1900	0.8729	0.9276	0.046*	0.741 (4)
C26	1.0890 (7)	0.6327 (4)	0.9057 (3)	0.0340 (13)	0.741 (4)
H26A	1.1286	0.5649	0.9240	0.041*	0.741 (4)
H26B	1.1622	0.6637	0.8663	0.041*	0.741 (4)
H26C	0.9525	0.6095	0.8815	0.041*	0.741 (4)
O4'	1.229 (8)	0.603 (3)	1.0485 (10)	0.0317 (16)	0.26
C24'	1.1591 (17)	0.6115 (9)	0.9800 (7)	0.018 (3)	0.259 (4)
H24'	1.1272	0.5434	0.9429	0.022*	0.259 (4)
N4'	1.123 (5)	0.7071 (13)	0.9520 (7)	0.0216 (12)	0.26
C25'	1.0462 (17)	0.7099 (12)	0.8680 (6)	0.022 (3)	0.259 (4)
H25D	1.1422	0.7618	0.8453	0.026*	0.259 (4)
H25E	0.9312	0.7383	0.8630	0.026*	0.259 (4)
H25F	1.0138	0.6314	0.8395	0.026*	0.259 (4)
C26'	1.1704 (18)	0.8156 (10)	1.0052 (7)	0.019 (3)	0.259 (4)
H26D	1.0541	0.8407	1.0070	0.023*	0.259 (4)
H26E	1.2613	0.8756	0.9854	0.023*	0.259 (4)
H26F	1.2280	0.8034	1.0587	0.023*	0.259 (4)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.0175 (2)	0.01176 (19)	0.01046 (18)	0.00541 (16)	0.00143 (15)	0.00295 (15)
Br1	0.02559 (19)	0.01350 (16)	0.02113 (17)	0.00819 (14)	0.00104 (14)	0.00254 (13)

Br2	0.02115 (18)	0.0359 (2)	0.01140 (16)	0.00857 (15)	-0.00079 (13)	0.00359 (14)
Br3	0.02659 (19)	0.0333 (2)	0.01011 (16)	0.00644 (15)	0.00017 (14)	0.00550 (14)
Br4	0.0337 (2)	0.01807 (17)	0.01618 (16)	0.01086 (15)	0.00296 (14)	0.00318 (13)
O1	0.0191 (12)	0.0135 (11)	0.0094 (10)	0.0063 (9)	-0.0006 (9)	0.0018 (9)
O2	0.0226 (12)	0.0148 (11)	0.0090 (10)	0.0081 (10)	0.0009 (9)	0.0018 (9)
N1	0.0110 (13)	0.0112 (13)	0.0143 (13)	0.0025 (10)	0.0010 (11)	0.0016 (11)
N2	0.0157 (14)	0.0127 (13)	0.0164 (13)	0.0049 (11)	0.0060 (11)	0.0050 (11)
C1	0.0115 (15)	0.0142 (16)	0.0120 (15)	0.0013 (12)	0.0015 (12)	0.0045 (13)
C2	0.0134 (16)	0.0121 (15)	0.0160 (16)	0.0029 (12)	0.0038 (13)	0.0035 (13)
C3	0.0103 (15)	0.0182 (17)	0.0167 (16)	0.0031 (13)	0.0013 (13)	0.0080 (13)
C4	0.0119 (16)	0.0222 (18)	0.0114 (15)	0.0017 (13)	-0.0014 (13)	0.0032 (14)
C5	0.0166 (17)	0.0175 (17)	0.0127 (15)	0.0043 (13)	0.0039 (13)	-0.0017 (13)
C6	0.0146 (16)	0.0131 (16)	0.0118 (15)	0.0057 (13)	0.0043 (13)	0.0027 (13)
C7	0.0143 (16)	0.0127 (16)	0.0160 (16)	0.0014 (13)	0.0070 (13)	0.0017 (13)
C8	0.0093 (15)	0.0114 (15)	0.0209 (16)	0.0021 (12)	0.0058 (13)	0.0060 (13)
C9	0.0149 (16)	0.0156 (16)	0.0202 (17)	0.0039 (13)	0.0013 (14)	0.0027 (14)
C10	0.0188 (17)	0.0125 (16)	0.0267 (18)	0.0047 (14)	0.0064 (15)	0.0015 (14)
C11	0.0241 (19)	0.0177 (17)	0.0294 (19)	0.0095 (14)	0.0117 (15)	0.0127 (15)
C12	0.0197 (17)	0.0226 (18)	0.0181 (17)	0.0098 (14)	0.0088 (14)	0.0086 (14)
C13	0.0145 (16)	0.0113 (15)	0.0179 (16)	0.0041 (12)	0.0064 (13)	0.0031 (13)
C14	0.0146 (16)	0.0183 (17)	0.0144 (15)	0.0055 (13)	0.0049 (13)	0.0094 (13)
C15	0.0122 (16)	0.0202 (17)	0.0120 (15)	0.0049 (13)	0.0017 (13)	0.0029 (13)
C16	0.0147 (16)	0.0237 (18)	0.0150 (16)	0.0085 (14)	0.0028 (13)	0.0101 (14)
C17	0.0158 (17)	0.0257 (18)	0.0085 (15)	0.0039 (14)	0.0001 (13)	0.0027 (14)
C18	0.0152 (17)	0.0217 (18)	0.0151 (16)	0.0047 (14)	0.0036 (13)	0.0003 (14)
C19	0.0168 (17)	0.0161 (16)	0.0159 (16)	0.0066 (13)	0.0042 (13)	0.0049 (13)
C20	0.0113 (16)	0.0193 (17)	0.0127 (15)	0.0041 (13)	0.0034 (12)	0.0037 (13)
O3	0.0247 (13)	0.0146 (12)	0.0243 (13)	0.0014 (10)	0.0073 (11)	-0.0015 (10)
N3	0.0296 (17)	0.0163 (15)	0.0258 (16)	0.0010 (13)	0.0043 (13)	-0.0002 (13)
C21	0.030 (2)	0.029 (2)	0.0271 (19)	0.0131 (17)	0.0084 (17)	0.0096 (17)
C22	0.036 (2)	0.0224 (19)	0.0259 (19)	0.0092 (17)	0.0055 (17)	0.0013 (16)
C23	0.047 (3)	0.023 (2)	0.057 (3)	-0.0038 (19)	-0.012 (2)	-0.001 (2)
O4	0.0336 (19)	0.031 (2)	0.029 (3)	0.0112 (18)	-0.003 (4)	0.011 (2)
C24	0.021 (3)	0.032 (3)	0.027 (3)	0.004 (2)	-0.002 (2)	0.006 (2)
N4	0.0193 (16)	0.021 (2)	0.024 (3)	0.005 (2)	0.001 (4)	0.007 (2)
C25	0.030 (3)	0.035 (3)	0.056 (4)	0.012 (3)	0.014 (3)	0.023 (3)
C26	0.024 (3)	0.049 (4)	0.032 (3)	0.010 (3)	0.008 (2)	0.012 (3)
O4'	0.0336 (19)	0.031 (2)	0.029 (3)	0.0112 (18)	-0.003 (4)	0.011 (2)
C24'	0.013 (6)	0.013 (6)	0.028 (7)	-0.004 (5)	0.007 (6)	0.011 (6)
N4'	0.0193 (16)	0.021 (2)	0.024 (3)	0.005 (2)	0.001 (4)	0.007 (2)
C25'	0.014 (7)	0.037 (8)	0.014 (6)	0.006 (6)	-0.003 (5)	0.016 (6)
C26'	0.013 (7)	0.021 (8)	0.024 (7)	0.005 (6)	0.004 (6)	-0.002 (6)

*Geometric parameters (Å, °)*

Cu1—O1	1.910 (2)	C16—C17	1.372 (4)
Cu1—O2	1.9174 (19)	C16—H16	0.95
Cu1—N1	1.954 (2)	C17—C18	1.397 (4)
Cu1—N2	1.958 (2)	C18—C19	1.367 (4)

## supplementary materials

---

Cu1—O3	2.501 (2)	C18—H18	0.95
Br1—C2	1.894 (3)	C19—C20	1.423 (4)
Br2—C4	1.911 (3)	O3—C21	1.247 (4)
Br3—C17	1.897 (3)	N3—C21	1.316 (4)
Br4—C19	1.901 (3)	N3—C23	1.441 (4)
O1—C1	1.295 (3)	N3—C22	1.471 (4)
O2—C20	1.294 (3)	C21—H21	0.95
N1—C7	1.291 (4)	C22—H22A	0.98
N1—C8	1.416 (4)	C22—H22B	0.98
N2—C14	1.293 (4)	C22—H22C	0.98
N2—C13	1.426 (4)	C23—H23A	0.98
C1—C2	1.418 (4)	C23—H23B	0.98
C1—C6	1.423 (4)	C23—H23C	0.98
C2—C3	1.369 (4)	O4—C24	1.218 (3)
C3—C4	1.390 (4)	C24—N4	1.340 (3)
C3—H3	0.95	C24—H24	0.95
C4—C5	1.361 (4)	N4—C26	1.456 (3)
C5—C6	1.416 (4)	N4—C25	1.457 (3)
C5—H5	0.95	C25—H25A	0.98
C6—C7	1.434 (4)	C25—H25B	0.98
C7—H7	0.95	C25—H25C	0.98
C8—C9	1.399 (4)	C26—H26A	0.98
C8—C13	1.405 (4)	C26—H26B	0.98
C9—C10	1.377 (4)	C26—H26C	0.98
C9—H9	0.95	O4'—C24'	1.220 (3)
C10—C11	1.387 (4)	C24'—N4'	1.340 (3)
C10—H10	0.95	C24'—H24'	0.95
C11—C12	1.375 (4)	N4'—C25'	1.460 (3)
C11—H11	0.95	N4'—C26'	1.460 (3)
C12—C13	1.389 (4)	C25'—H25D	0.98
C12—H12	0.95	C25'—H25E	0.98
C14—C15	1.428 (4)	C25'—H25F	0.98
C14—H14	0.95	C26'—H26D	0.98
C15—C16	1.418 (4)	C26'—H26E	0.98
C15—C20	1.427 (4)	C26'—H26F	0.98
O1—Cu1—O2	89.11 (8)	C16—C15—C14	115.6 (3)
O1—Cu1—N1	93.40 (9)	C20—C15—C14	123.6 (3)
O2—Cu1—N1	177.19 (10)	C17—C16—C15	120.3 (3)
O1—Cu1—N2	170.18 (10)	C17—C16—H16	119.9
O2—Cu1—N2	93.64 (9)	C15—C16—H16	119.9
N1—Cu1—N2	83.67 (10)	C16—C17—C18	120.7 (3)
O1—Cu1—O3	96.00 (9)	C16—C17—Br3	119.8 (2)
O2—Cu1—O3	89.70 (8)	C18—C17—Br3	119.4 (2)
N1—Cu1—O3	91.29 (9)	C19—C18—C17	119.0 (3)
N2—Cu1—O3	93.44 (9)	C19—C18—H18	120.5
C1—O1—Cu1	126.73 (19)	C17—C18—H18	120.5
C20—O2—Cu1	126.9 (2)	C18—C19—C20	124.0 (3)
C7—N1—C8	121.5 (3)	C18—C19—Br4	118.1 (2)
C7—N1—Cu1	125.3 (2)	C20—C19—Br4	117.9 (2)

C8—N1—Cu1	113.22 (19)	O2—C20—C19	119.9 (3)
C14—N2—C13	121.4 (3)	O2—C20—C15	124.9 (3)
C14—N2—Cu1	125.7 (2)	C19—C20—C15	115.2 (3)
C13—N2—Cu1	112.92 (19)	C21—O3—Cu1	126.8 (2)
O1—C1—C2	119.9 (3)	C21—N3—C23	123.2 (3)
O1—C1—C6	125.0 (3)	C21—N3—C22	120.3 (3)
C2—C1—C6	115.1 (3)	C23—N3—C22	116.3 (3)
C3—C2—C1	123.8 (3)	O3—C21—N3	125.4 (3)
C3—C2—Br1	118.3 (2)	O3—C21—H21	117.3
C1—C2—Br1	117.8 (2)	N3—C21—H21	117.3
C2—C3—C4	118.6 (3)	N3—C22—H22A	109.5
C2—C3—H3	120.7	N3—C22—H22B	109.5
C4—C3—H3	120.7	H22A—C22—H22B	109.5
C5—C4—C3	121.8 (3)	N3—C22—H22C	109.5
C5—C4—Br2	119.8 (2)	H22A—C22—H22C	109.5
C3—C4—Br2	118.5 (2)	H22B—C22—H22C	109.5
C4—C5—C6	119.3 (3)	N3—C23—H23A	109.5
C4—C5—H5	120.3	N3—C23—H23B	109.5
C6—C5—H5	120.3	H23A—C23—H23B	109.5
C5—C6—C1	121.4 (3)	N3—C23—H23C	109.5
C5—C6—C7	115.7 (3)	H23A—C23—H23C	109.5
C1—C6—C7	123.0 (3)	H23B—C23—H23C	109.5
N1—C7—C6	125.4 (3)	O4—C24—N4	124.5 (4)
N1—C7—H7	117.3	O4—C24—H24	117.8
C6—C7—H7	117.3	N4—C24—H24	117.8
C9—C8—C13	119.4 (3)	C24—N4—C26	121.2 (4)
C9—C8—N1	125.2 (3)	C24—N4—C25	120.7 (4)
C13—C8—N1	115.3 (3)	C26—N4—C25	118.0 (4)
C10—C9—C8	119.8 (3)	O4'—C24'—N4'	126.4 (14)
C10—C9—H9	120.1	O4'—C24'—H24'	116.8
C8—C9—H9	120.1	N4'—C24'—H24'	116.8
C9—C10—C11	120.1 (3)	C24'—N4'—C25'	122.5 (11)
C9—C10—H10	119.9	C24'—N4'—C26'	119.9 (11)
C11—C10—H10	119.9	C25'—N4'—C26'	117.5 (10)
C12—C11—C10	121.2 (3)	N4'—C25'—H25D	109.5
C12—C11—H11	119.4	N4'—C25'—H25E	109.5
C10—C11—H11	119.4	H25D—C25'—H25E	109.5
C11—C12—C13	119.3 (3)	N4'—C25'—H25F	109.5
C11—C12—H12	120.3	H25D—C25'—H25F	109.5
C13—C12—H12	120.3	H25E—C25'—H25F	109.5
C12—C13—C8	120.1 (3)	N4'—C26'—H26D	109.5
C12—C13—N2	125.0 (3)	N4'—C26'—H26E	109.5
C8—C13—N2	114.8 (3)	H26D—C26'—H26E	109.5
N2—C14—C15	125.1 (3)	N4'—C26'—H26F	109.5
N2—C14—H14	117.4	H26D—C26'—H26F	109.5
C15—C14—H14	117.4	H26E—C26'—H26F	109.5
C16—C15—C20	120.8 (3)		
O2—Cu1—O1—C1	169.2 (2)	C8—C9—C10—C11	0.5 (5)
N1—Cu1—O1—C1	-12.0 (2)	C9—C10—C11—C12	-1.2 (5)

## supplementary materials

O3—Cu1—O1—C1	79.6 (2)	C10—C11—C12—C13	0.4 (5)
O1—Cu1—O2—C20	171.6 (2)	C11—C12—C13—C8	1.0 (5)
N2—Cu1—O2—C20	1.1 (3)	C11—C12—C13—N2	-178.7 (3)
O3—Cu1—O2—C20	-92.4 (2)	C9—C8—C13—C12	-1.6 (5)
O1—Cu1—N1—C7	9.3 (3)	N1—C8—C13—C12	178.3 (3)
N2—Cu1—N1—C7	179.9 (3)	C9—C8—C13—N2	178.1 (3)
O3—Cu1—N1—C7	-86.8 (3)	N1—C8—C13—N2	-1.9 (4)
O1—Cu1—N1—C8	-171.3 (2)	C14—N2—C13—C12	-1.5 (5)
N2—Cu1—N1—C8	-0.7 (2)	Cu1—N2—C13—C12	-179.0 (3)
O3—Cu1—N1—C8	92.6 (2)	C14—N2—C13—C8	178.7 (3)
O2—Cu1—N2—C14	1.5 (3)	Cu1—N2—C13—C8	1.3 (3)
N1—Cu1—N2—C14	-177.6 (3)	C13—N2—C14—C15	-179.1 (3)
O3—Cu1—N2—C14	91.5 (3)	Cu1—N2—C14—C15	-2.0 (4)
O2—Cu1—N2—C13	178.8 (2)	N2—C14—C15—C16	179.7 (3)
N1—Cu1—N2—C13	-0.3 (2)	N2—C14—C15—C20	-0.4 (5)
O3—Cu1—N2—C13	-91.2 (2)	C20—C15—C16—C17	-0.6 (5)
Cu1—O1—C1—C2	-172.1 (2)	C14—C15—C16—C17	179.3 (3)
Cu1—O1—C1—C6	8.9 (4)	C15—C16—C17—C18	-0.9 (5)
O1—C1—C2—C3	179.9 (3)	C15—C16—C17—Br3	-179.6 (2)
C6—C1—C2—C3	-1.0 (5)	C16—C17—C18—C19	0.4 (5)
O1—C1—C2—Br1	1.0 (4)	Br3—C17—C18—C19	179.2 (2)
C6—C1—C2—Br1	-179.9 (2)	C17—C18—C19—C20	1.5 (5)
C1—C2—C3—C4	0.7 (5)	C17—C18—C19—Br4	-179.2 (2)
Br1—C2—C3—C4	179.6 (2)	Cu1—O2—C20—C19	177.6 (2)
C2—C3—C4—C5	0.0 (5)	Cu1—O2—C20—C15	-3.4 (4)
C2—C3—C4—Br2	-179.4 (2)	C18—C19—C20—O2	176.3 (3)
C3—C4—C5—C6	-0.3 (5)	Br4—C19—C20—O2	-3.0 (4)
Br2—C4—C5—C6	179.1 (2)	C18—C19—C20—C15	-2.8 (5)
C4—C5—C6—C1	0.0 (5)	Br4—C19—C20—C15	177.9 (2)
C4—C5—C6—C7	178.6 (3)	C16—C15—C20—O2	-176.8 (3)
O1—C1—C6—C5	179.7 (3)	C14—C15—C20—O2	3.3 (5)
C2—C1—C6—C5	0.6 (4)	C16—C15—C20—C19	2.3 (4)
O1—C1—C6—C7	1.1 (5)	C14—C15—C20—C19	-177.6 (3)
C2—C1—C6—C7	-177.9 (3)	O1—Cu1—O3—C21	94.1 (3)
C8—N1—C7—C6	177.3 (3)	O2—Cu1—O3—C21	5.0 (3)
Cu1—N1—C7—C6	-3.4 (4)	N1—Cu1—O3—C21	-172.4 (3)
C5—C6—C7—N1	177.4 (3)	N2—Cu1—O3—C21	-88.6 (3)
C1—C6—C7—N1	-4.0 (5)	Cu1—O3—C21—N3	-66.7 (4)
C7—N1—C8—C9	1.1 (5)	C23—N3—C21—O3	-179.6 (4)
Cu1—N1—C8—C9	-178.4 (2)	C22—N3—C21—O3	-4.2 (5)
C7—N1—C8—C13	-178.9 (3)	O4—C24—N4—C26	0.4 (17)
Cu1—N1—C8—C13	1.6 (3)	O4—C24—N4—C25	-176.8 (12)
C13—C8—C9—C10	0.9 (5)	O4'—C24'—N4'—C25'	177 (4)
N1—C8—C9—C10	-179.1 (3)	O4'—C24'—N4'—C26'	1(6)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C5-H5\cdots O3^i$	0.95	2.47	3.319 (4)	148

C7—H7…O3 <sup>i</sup>	0.95	2.37	3.243 (4)	153
C14—H14…O4 <sup>ii</sup>	0.95	2.33	3.226 (14)	156
C22—H22A…O1	0.98	2.54	3.378 (4)	143

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+2, -y+1, -z+2$ .

Fig. 1

