

Dichloridobis(phenanthridine- κ N)-zinc(II)

 Zeinab Khoshtarkib,^a Amin Ebadi,^b Robabeh Alizadeh,^{c*} Roya Ahmadi^a and Vahid Amani^a

^aIslamic Azad University, Shahr-e-Rey Branch, Tehran, Iran, ^bDepartment of Chemistry, Islamic Azad University, Kazerun Branch, Kazerun, Fars, Iran, and ^cDamghan University of Basic Sciences, School of Chemistry, Damghan, Iran
Correspondence e-mail: alizadeh@dubs.ac.ir


Received 20 May 2009; accepted 22 May 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.086; data-to-parameter ratio = 20.5.

In the molecule of the title compound, $[\text{ZnCl}_2(\text{C}_{13}\text{H}_9\text{N})_2]$, the Zn^{II} atom is four-coordinated in a distorted tetrahedral configuration by two N atoms from two phenanthridine ligands and by two terminal Cl atoms. The dihedral angle between the planes of the phenanthridine ring systems is $69.92(3)^\circ$. An intramolecular $\text{C}-\text{H}\cdots\text{Cl}$ interaction results in the formation of a planar five-membered ring, which is oriented at a dihedral angle of $8.32(3)^\circ$ with respect to the adjacent phenanthridine ring system. In the crystal structure, $\pi-\pi$ contacts between the phenanthridine systems [centroid-centroid distances = 3.839 (2), 3.617 (1) and 3.682 (1) Å] may stabilize the structure. Two weak $\text{C}-\text{H}\cdots\pi$ interactions are also found.

Related literature

For related structures, see: Ahmadi *et al.* (2008); Çelik *et al.* (2004); Cui *et al.* (1998); Gruia *et al.* (2007); Khalighi *et al.* (2008); Khan & Tuck (1984); Khavasi *et al.* (2008); Kozhevnikov *et al.* (2006); Liu *et al.* (2004); Markowitz *et al.* (2006); Musie *et al.* (2004); Preston & Kennard (1969); Reimann *et al.* (1966); Shen *et al.* (2004); Steffen & Palenik (1977). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$[\text{ZnCl}_2(\text{C}_{13}\text{H}_9\text{N})_2]$	$V = 2130.0(9)$ Å ³
$M_r = 494.71$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 16.193(3)$ Å	$\mu = 1.42$ mm ⁻¹
$b = 10.101(2)$ Å	$T = 298$ K
$c = 14.491(3)$ Å	$0.45 \times 0.30 \times 0.22$ mm
$\beta = 116.02(3)^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	16947 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 1998)	5732 independent reflections
$T_{\min} = 0.610$, $T_{\max} = 0.740$	4612 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	280 parameters
$wR(F^2) = 0.086$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\text{max}} = 0.28$ e Å ⁻³
5732 reflections	$\Delta\rho_{\text{min}} = -0.39$ e Å ⁻³

Table 1

Selected geometric parameters (Å, °).

Cl1—Zn1	2.2234 (7)	N1—Zn1	2.0785 (17)
Cl2—Zn1	2.2456 (7)	N2—Zn1	2.0775 (17)
N2—Zn1—N1	105.19 (7)	N2—Zn1—Cl2	113.54 (5)
N2—Zn1—Cl1	108.18 (5)	N1—Zn1—Cl2	107.46 (6)
N1—Zn1—Cl1	106.23 (5)	Cl1—Zn1—Cl2	115.49 (3)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
Cl1—H1 \cdots Cl1	0.93	2.77	3.434 (3)	129
C17—H17 \cdots Cg6 ⁱ	0.93	2.82	3.535 (3)	134
C24—H24 \cdots Cg5 ⁱⁱ	0.93	2.81	3.508 (3)	132

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z - \frac{1}{2}$; (ii) $x, -y - \frac{1}{2}, z - \frac{1}{2}$. Cg5 and Cg6 are the centroids of the C15–C20 and C21–C26 rings, respectively.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

We are grateful to Damghan University of Basic Sciences for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2696).

References

- Ahmadi, R., Kalateh, K., Ebadi, A., Amani, V. & Khavasi, H. R. (2008). *Acta Cryst.* **E64**, m1266.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (1998). *SMART, SAINT and SADABS*. Bruker AXS, Madison, Wisconsin, USA.
- Çelik, Ö., İde, S., Kurt, M. & Yurdakul, Ş. (2004). *Acta Cryst.* **E60**, m1134–m1136.
- Cui, Y., Long, D., Chen, W. & Huang, J. (1998). *Acta Cryst.* **C54**, 1605–1607.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gruia, L. M., Rochon, F. D. & Beauchamp, A. L. (2007). *Inorg. Chim. Acta*, **360**, 1825–1840.
- Khalighi, A., Ahmadi, R., Amani, V. & Khavasi, H. R. (2008). *Acta Cryst.* **E64**, m1211–m1212.
- Khan, M. A. & Tuck, D. G. (1984). *Acta Cryst.* **C40**, 60–62.
- Khavasi, H. R., Abedi, A., Amani, V., Notash, B. & Safari, N. (2008). *Polyhedron*, **27**, 1848–1854.
- Kozhevnikov, D. N., Shabunina, O. V., Kopchuk, D. S., Slepukhin, P. A. & Kozhevnikov, V. N. (2006). *Tetrahedron Lett.* **47**, 7025–7029.
- Liu, Q. D., Wang, R. & Wang, S. (2004). *Dalton Trans.* pp. 2073–2079.
- Markowitz, B. M. E., Turnbull, M. M. & Awwadi, F. F. (2006). *Acta Cryst.* **E62**, m1278–m1280.
- Musie, G. T., Li, X. & Powell, D. R. (2004). *Acta Cryst.* **E60**, m471–m472.
- Preston, H. S. & Kennard, C. H. L. (1969). *J. Chem. Soc. A*, pp. 1965–1968.
- Reimann, C. W., Block, S. & Perloff, A. (1966). *Inorg. Chem.* **5**, 1185–1189.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shen, L., Li, M. C., Jin, Z. M., Hu, M. L. & Xuan, R. C. (2004). *Acta Cryst.* **E60**, m330–m331.
- Steffen, W. L. & Palenik, G. J. (1977). *Inorg. Chem.* **16**, 1119–1127.

supplementary materials

Acta Cryst. (2009). E65, m739-m740 [doi:10.1107/S160053680901959X]

Dichloridobis(phenanthridine- κ N)zinc(II)

Z. Khoshtarkib, A. Ebadi, R. Alizadeh, R. Ahmadi and V. Amani

Comment

There are several Zn^{II} complexes, with formula, [ZnCl₂(N)₂], such as [ZnCl₂(AMS)₂], (II) (Shen *et al.*, 2004), [ZnCl₂(4-CNpy)₂], (III) (Steffen & Palenik, 1977), [ZnCl₂(pht)₂], (IV) (Çelik *et al.*, 2004), [ZnCl₂(quin)₂], (V) (Cui *et al.*, 1998), [ZnCl₂(quino)₂], (VI) (Markowitz *et al.*, 2006) and [ZnCl₂(meim)₂], (VII) (Musie *et al.*, 2004) [where AMS is 3-Amino-5-methylisoxazole, 4-CNpy is 4-cyanopyridine, pht is phthalazine, quin is quinoline, quino is quinoxaline and meim is 1-methylimidazole] have been synthesized and characterized by single-crystal X-ray diffraction methods.

There are also several Zn^{II} complexes, with formula, [ZnCl₂(N—N)], such as [ZnCl₂(bipy)], (VIII) (Khan & Tuck, 1984), [ZnCl₂(biim)], (IX) (Gruia *et al.*, 2007), [ZnCl₂(phbipy)], (X) (Kozhevnikov *et al.*, 2006), [ZnCl₂(phen)], (XI) (Reimann *et al.*, 1966), [ZnCl₂(dmphen)], (XII) (Preston & Kennard, 1969), [ZnCl₂(dpdmbip)], (XIII) (Liu *et al.*, 2004), [ZnCl₂(dm4bt)], (XIV) (Khavasi *et al.*, 2008), [ZnCl₂(5,5'-dmbpy)], (XV) (Khalighi *et al.*, 2008) and [ZnCl₂(6-mbipy)], (XVI) (Ahmadi *et al.*, 2008) [where bipy is 2,2'-bipyridine, biim is 2,2'-biimidazole, phbipy is 5-phenyl-2,2'-bipyridine, phen is 1,10-phenanthroline, dmphen is 2,9-dimethyl-1,10-phenanthroline, dpdmbip is 4,4'-diphenyl-6,6'-dimethyl-2,2'-bipyrimidine, dm4bt is 2,2'-dimethyl-4,4'-bithiazole, 5,5'-dmbpy is 5,5'-dimethyl-2,2'-bipyridine and 6-mbipy is 6-methyl-2,2'-bipyridine] have been synthesized and characterized by single-crystal X-ray diffraction methods. We report herein the synthesis and crystal structure of the title compound, (I).

In the molecule of the title compound (Fig 1), Zn^{II} atom is four-coordinated in a distorted tetrahedral configuration by two N atoms from two phenanthridine and two terminal Cl atoms (Table 1). The bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Phenanthridine ring systems A (N1/C1-C13) and B (N2/C14-C26) are, of course, planar and the dihedral angle between them is A/B = 69.92 (3)°. Intramolecular C—H...Cl interaction (Table 2) results in the formation of a planar five-membered ring C (Zn1/Cl1/N1/C1/H1), which is oriented with respect to the adjacent phenanthridine ring system A at a dihedral angle of 8.32 (3)°.

In the crystal structure (Fig. 2), the π - π contacts between the phenanthridine rings, Cg2—Cg3ⁱ, Cg4—Cg6ⁱⁱ and Cg6—Cg6ⁱⁱ, [symmetry codes: (i) 1 - x, -y, -z, (ii) -x, -y, -z, where Cg2, Cg3, Cg4 and Cg6 are centroids of the rings (C2-C7), (C8-C13), (N2/C14/C15/C20/C21/C26) and (C21-C26), respectively] may stabilize the structure, with centroid-centroid distances of 3.839 (2), 3.617 (1) and 3.682 (1) Å, respectively. There also exist two weak C—H... π interactions (Table 2).

Experimental

For the preparation of the title compound, (I), a solution of phenanthridine (0.30 g, 1.66 mmol) in methanol (15 ml) was added to a solution of ZnCl₂ (0.11 g, 0.83 mmol) in acetonitrile (30 ml) and the resulting colorless solution was stirred for 30 min at 313 K, and then it was left to evaporate slowly at room temperature. After one week, colorless prismatic crystals of the title compound were isolated (yield; 0.31 g, 75.5%).

Refinement

H atoms were positioned geometrically, with C-H = 0.93 Å for aromatic H and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 40% probability level.




Fig. 2. A partial packing diagram of the title compound.

Dichloridobis(phenanthridine-κN)zinc(II)

Crystal data

$[\text{ZnCl}_2(\text{C}_{13}\text{H}_9\text{N}_1)_2]$

$M_r = 494.71$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.193(3) \text{ \AA}$

$b = 10.101(2) \text{ \AA}$

$c = 14.491(3) \text{ \AA}$

$\beta = 116.02(3)^\circ$

$V = 2130.0(9) \text{ \AA}^3$

$Z = 4$

$F_{000} = 1008$

$D_x = 1.543 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1987 reflections

$\theta = 2.5\text{--}29.3^\circ$

$\mu = 1.42 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Prism, colorless

$0.45 \times 0.30 \times 0.22 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

5732 independent reflections

4612 reflections with $I > 2\sigma(I)$

Monochromator: graphite $R_{\text{int}} = 0.041$
 $T = 298 \text{ K}$ $\theta_{\text{max}} = 29.3^\circ$
 φ and ω scans $\theta_{\text{min}} = 2.5^\circ$
 Absorption correction: multi-scan
 (SADABS; Bruker, 1998) $h = -22 \rightarrow 22$
 $T_{\text{min}} = 0.610$, $T_{\text{max}} = 0.740$ $k = -13 \rightarrow 13$
 16947 measured reflections $l = -19 \rightarrow 19$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.037$ H-atom parameters constrained
 $wR(F^2) = 0.086$ $w = 1/[\sigma^2(F_o^2) + (0.038P)^2 + 0.5741P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.09$ $(\Delta/\sigma)_{\text{max}} = 0.013$
 5732 reflections $\Delta\rho_{\text{max}} = 0.28 \text{ e } \text{\AA}^{-3}$
 280 parameters $\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.259400 (15)	0.26111 (2)	0.120119 (17)	0.03590 (7)
Cl1	0.25049 (4)	0.47913 (5)	0.09659 (5)	0.05264 (14)
Cl2	0.27473 (4)	0.19233 (6)	0.27420 (4)	0.05102 (14)
N1	0.37559 (11)	0.19993 (17)	0.10537 (13)	0.0373 (3)
N2	0.14942 (10)	0.17454 (16)	-0.00248 (12)	0.0338 (3)
C1	0.42820 (14)	0.2930 (2)	0.09724 (16)	0.0410 (4)
H1	0.4056	0.3791	0.0863	0.049*
C2	0.51762 (14)	0.2716 (2)	0.10399 (16)	0.0417 (4)
C3	0.57131 (17)	0.3780 (3)	0.09923 (19)	0.0528 (6)
H3	0.5481	0.4637	0.0902	0.063*
C4	0.65812 (18)	0.3558 (3)	0.1079 (2)	0.0626 (7)

supplementary materials

H4	0.6936	0.4260	0.1039	0.075*
C5	0.69249 (18)	0.2273 (3)	0.1228 (2)	0.0674 (8)
H5	0.7512	0.2124	0.1283	0.081*
C6	0.64215 (16)	0.1227 (3)	0.1293 (2)	0.0574 (6)
H6	0.6672	0.0380	0.1401	0.069*
C7	0.55249 (14)	0.1417 (2)	0.11974 (15)	0.0432 (5)
C8	0.49487 (14)	0.0368 (2)	0.12662 (15)	0.0409 (4)
C9	0.52121 (17)	-0.0978 (2)	0.13981 (17)	0.0503 (5)
H9	0.5782	-0.1218	0.1441	0.060*
C10	0.46418 (19)	-0.1931 (2)	0.14633 (18)	0.0548 (6)
H10	0.4828	-0.2812	0.1547	0.066*
C11	0.37861 (18)	-0.1601 (2)	0.14065 (18)	0.0517 (5)
H11	0.3404	-0.2259	0.1453	0.062*
C12	0.35073 (16)	-0.0305 (2)	0.12812 (16)	0.0448 (5)
H12	0.2939	-0.0084	0.1251	0.054*
C13	0.40733 (14)	0.0686 (2)	0.11980 (14)	0.0377 (4)
C14	0.16296 (13)	0.12723 (19)	-0.07864 (15)	0.0360 (4)
H14	0.2222	0.1316	-0.0736	0.043*
C15	0.09352 (13)	0.06958 (18)	-0.16859 (14)	0.0348 (4)
C16	0.11488 (16)	0.0160 (2)	-0.24580 (16)	0.0417 (4)
H16	0.1747	0.0203	-0.2389	0.050*
C17	0.04765 (18)	-0.0419 (2)	-0.33016 (18)	0.0486 (5)
H17	0.0615	-0.0771	-0.3810	0.058*
C18	-0.04216 (17)	-0.0482 (2)	-0.34024 (18)	0.0510 (5)
H18	-0.0876	-0.0881	-0.3981	0.061*
C19	-0.06479 (15)	0.0033 (2)	-0.26642 (17)	0.0454 (5)
H19	-0.1250	-0.0022	-0.2746	0.054*
C20	0.00307 (13)	0.06448 (18)	-0.17833 (14)	0.0343 (4)
C21	-0.01463 (13)	0.12123 (18)	-0.09734 (15)	0.0343 (4)
C22	-0.10280 (14)	0.1282 (2)	-0.10056 (17)	0.0430 (5)
H22	-0.1531	0.0952	-0.1574	0.052*
C23	-0.11556 (15)	0.1830 (2)	-0.02101 (19)	0.0481 (5)
H23	-0.1743	0.1872	-0.0247	0.058*
C24	-0.04179 (17)	0.2319 (2)	0.06441 (19)	0.0476 (5)
H24	-0.0510	0.2678	0.1183	0.057*
C25	0.04526 (15)	0.2277 (2)	0.07015 (17)	0.0417 (4)
H25	0.0947	0.2608	0.1279	0.050*
C26	0.05989 (12)	0.17376 (18)	-0.01046 (14)	0.0328 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.02853 (11)	0.03705 (12)	0.03880 (12)	-0.00174 (9)	0.01173 (8)	-0.00362 (9)
Cl1	0.0510 (3)	0.0376 (3)	0.0687 (4)	0.0014 (2)	0.0258 (3)	-0.0007 (2)
Cl2	0.0481 (3)	0.0596 (3)	0.0436 (3)	-0.0047 (3)	0.0185 (2)	0.0027 (2)
N1	0.0312 (8)	0.0388 (8)	0.0385 (8)	0.0034 (7)	0.0120 (7)	-0.0012 (7)
N2	0.0265 (7)	0.0339 (7)	0.0374 (8)	-0.0007 (6)	0.0107 (6)	-0.0010 (6)
C1	0.0341 (10)	0.0424 (10)	0.0442 (11)	0.0019 (8)	0.0149 (8)	-0.0014 (8)

C2	0.0317 (9)	0.0543 (12)	0.0367 (10)	-0.0003 (9)	0.0127 (8)	-0.0022 (9)
C3	0.0439 (12)	0.0640 (15)	0.0506 (13)	-0.0062 (11)	0.0207 (10)	-0.0017 (11)
C4	0.0429 (13)	0.087 (2)	0.0608 (15)	-0.0129 (13)	0.0250 (12)	-0.0031 (14)
C5	0.0333 (11)	0.104 (2)	0.0666 (16)	0.0047 (14)	0.0232 (11)	-0.0006 (15)
C6	0.0354 (11)	0.0780 (17)	0.0564 (14)	0.0117 (12)	0.0178 (10)	0.0020 (12)
C7	0.0311 (9)	0.0620 (13)	0.0325 (10)	0.0082 (9)	0.0103 (8)	-0.0010 (9)
C8	0.0339 (9)	0.0511 (11)	0.0306 (9)	0.0092 (9)	0.0075 (7)	-0.0005 (8)
C9	0.0453 (12)	0.0557 (13)	0.0419 (11)	0.0201 (11)	0.0116 (10)	0.0029 (10)
C10	0.0656 (16)	0.0453 (12)	0.0409 (11)	0.0180 (12)	0.0116 (11)	0.0035 (9)
C11	0.0556 (14)	0.0425 (11)	0.0462 (12)	-0.0001 (10)	0.0123 (10)	0.0037 (9)
C12	0.0395 (11)	0.0457 (11)	0.0430 (11)	0.0007 (9)	0.0125 (9)	-0.0004 (9)
C13	0.0329 (9)	0.0412 (10)	0.0317 (9)	0.0044 (8)	0.0074 (7)	-0.0023 (8)
C14	0.0286 (9)	0.0373 (9)	0.0394 (10)	-0.0009 (7)	0.0125 (8)	0.0010 (8)
C15	0.0346 (9)	0.0313 (9)	0.0351 (9)	-0.0016 (7)	0.0121 (8)	0.0008 (7)
C16	0.0436 (11)	0.0391 (10)	0.0431 (11)	0.0040 (9)	0.0198 (9)	0.0000 (8)
C17	0.0593 (14)	0.0412 (11)	0.0448 (12)	-0.0011 (10)	0.0223 (11)	-0.0077 (9)
C18	0.0519 (13)	0.0433 (11)	0.0459 (12)	-0.0085 (10)	0.0105 (10)	-0.0100 (9)
C19	0.0383 (10)	0.0408 (10)	0.0488 (12)	-0.0094 (9)	0.0116 (9)	-0.0039 (9)
C20	0.0313 (9)	0.0286 (8)	0.0375 (9)	-0.0026 (7)	0.0099 (7)	0.0025 (7)
C21	0.0306 (9)	0.0295 (8)	0.0405 (10)	-0.0020 (7)	0.0133 (8)	0.0036 (7)
C22	0.0307 (9)	0.0444 (11)	0.0513 (12)	-0.0046 (8)	0.0155 (9)	-0.0001 (9)
C23	0.0359 (10)	0.0465 (11)	0.0676 (15)	0.0004 (9)	0.0278 (10)	0.0024 (10)
C24	0.0468 (12)	0.0447 (11)	0.0588 (13)	0.0003 (10)	0.0302 (11)	-0.0060 (10)
C25	0.0378 (10)	0.0409 (10)	0.0455 (11)	-0.0024 (8)	0.0175 (9)	-0.0062 (8)
C26	0.0287 (8)	0.0290 (8)	0.0394 (9)	-0.0012 (7)	0.0138 (7)	0.0014 (7)

Geometric parameters (Å, °)

C11—Zn1	2.2234 (7)	C12—H12	0.9300
C12—Zn1	2.2456 (7)	C13—N1	1.405 (3)
N1—Zn1	2.0785 (17)	C14—N2	1.306 (2)
N2—Zn1	2.0775 (17)	C14—C15	1.420 (3)
C1—N1	1.308 (3)	C14—H14	0.9300
C1—C2	1.424 (3)	C15—C20	1.410 (3)
C1—H1	0.9300	C15—C16	1.416 (3)
C2—C3	1.402 (3)	C16—C17	1.361 (3)
C2—C7	1.407 (3)	C16—H16	0.9300
C3—C4	1.374 (4)	C17—C18	1.398 (4)
C3—H3	0.9300	C17—H17	0.9300
C4—C5	1.392 (4)	C18—C19	1.376 (3)
C4—H4	0.9300	C18—H18	0.9300
C5—C6	1.363 (4)	C19—C20	1.410 (3)
C5—H5	0.9300	C19—H19	0.9300
C6—C7	1.410 (3)	C20—C21	1.443 (3)
C6—H6	0.9300	C21—C22	1.410 (3)
C7—C8	1.444 (3)	C21—C26	1.411 (3)
C8—C9	1.412 (3)	C22—C23	1.374 (3)
C8—C13	1.414 (3)	C22—H22	0.9300
C9—C10	1.366 (4)	C23—C24	1.380 (3)

supplementary materials

C9—H9	0.9300	C23—H23	0.9300
C10—C11	1.392 (4)	C24—C25	1.376 (3)
C10—H10	0.9300	C24—H24	0.9300
C11—C12	1.371 (3)	C25—C26	1.400 (3)
C11—H11	0.9300	C25—H25	0.9300
C12—C13	1.398 (3)	C26—N2	1.403 (2)
N2—Zn1—N1	105.19 (7)	C11—C12—C13	120.3 (2)
N2—Zn1—C11	108.18 (5)	C11—C12—H12	119.8
N1—Zn1—C11	106.23 (5)	C13—C12—H12	119.8
N2—Zn1—C12	113.54 (5)	C12—C13—N1	118.57 (18)
N1—Zn1—C12	107.46 (6)	C12—C13—C8	120.43 (19)
C11—Zn1—C12	115.49 (3)	N1—C13—C8	121.00 (19)
C1—N1—C13	118.80 (18)	N2—C14—C15	124.52 (18)
C1—N1—Zn1	116.76 (14)	N2—C14—H14	117.7
C13—N1—Zn1	123.65 (14)	C15—C14—H14	117.7
C14—N2—C26	118.60 (16)	C20—C15—C16	120.69 (18)
C14—N2—Zn1	118.54 (13)	C20—C15—C14	118.46 (18)
C26—N2—Zn1	122.69 (13)	C16—C15—C14	120.84 (19)
N1—C1—C2	124.6 (2)	C17—C16—C15	119.8 (2)
N1—C1—H1	117.7	C17—C16—H16	120.1
C2—C1—H1	117.7	C15—C16—H16	120.1
C3—C2—C7	120.6 (2)	C16—C17—C18	119.9 (2)
C3—C2—C1	120.9 (2)	C16—C17—H17	120.0
C7—C2—C1	118.4 (2)	C18—C17—H17	120.0
C4—C3—C2	120.2 (3)	C19—C18—C17	121.4 (2)
C4—C3—H3	119.9	C19—C18—H18	119.3
C2—C3—H3	119.9	C17—C18—H18	119.3
C3—C4—C5	119.3 (3)	C18—C19—C20	120.1 (2)
C3—C4—H4	120.3	C18—C19—H19	119.9
C5—C4—H4	120.3	C20—C19—H19	119.9
C6—C5—C4	121.5 (2)	C15—C20—C19	118.01 (19)
C6—C5—H5	119.2	C15—C20—C21	118.19 (17)
C4—C5—H5	119.2	C19—C20—C21	123.80 (18)
C5—C6—C7	120.6 (3)	C22—C21—C26	117.92 (18)
C5—C6—H6	119.7	C22—C21—C20	123.54 (18)
C7—C6—H6	119.7	C26—C21—C20	118.54 (17)
C2—C7—C6	117.8 (2)	C23—C22—C21	121.0 (2)
C2—C7—C8	118.03 (19)	C23—C22—H22	119.5
C6—C7—C8	124.2 (2)	C21—C22—H22	119.5
C9—C8—C13	117.5 (2)	C22—C23—C24	120.5 (2)
C9—C8—C7	123.3 (2)	C22—C23—H23	119.8
C13—C8—C7	119.14 (19)	C24—C23—H23	119.8
C10—C9—C8	121.0 (2)	C25—C24—C23	120.2 (2)
C10—C9—H9	119.5	C25—C24—H24	119.9
C8—C9—H9	119.5	C23—C24—H24	119.9
C9—C10—C11	120.9 (2)	C24—C25—C26	120.4 (2)
C9—C10—H10	119.6	C24—C25—H25	119.8
C11—C10—H10	119.6	C26—C25—H25	119.8
C12—C11—C10	119.8 (2)	C25—C26—N2	118.43 (17)

C12—C11—H11	120.1	C25—C26—C21	119.96 (18)
C10—C11—H11	120.1	N2—C26—C21	121.61 (17)
N1—C1—C2—C3	-177.0 (2)	C18—C19—C20—C21	179.9 (2)
N1—C1—C2—C7	0.5 (3)	C15—C20—C21—C22	176.71 (18)
C7—C2—C3—C4	1.3 (3)	C19—C20—C21—C22	-3.7 (3)
C1—C2—C3—C4	178.7 (2)	C15—C20—C21—C26	-2.7 (3)
C2—C3—C4—C5	-0.8 (4)	C19—C20—C21—C26	176.82 (19)
C3—C4—C5—C6	-0.2 (4)	C26—C21—C22—C23	-0.8 (3)
C4—C5—C6—C7	0.8 (4)	C20—C21—C22—C23	179.8 (2)
C3—C2—C7—C6	-0.7 (3)	C21—C22—C23—C24	-0.4 (3)
C1—C2—C7—C6	-178.2 (2)	C22—C23—C24—C25	0.8 (4)
C3—C2—C7—C8	178.41 (19)	C23—C24—C25—C26	0.0 (3)
C1—C2—C7—C8	0.9 (3)	C24—C25—C26—N2	178.40 (19)
C5—C6—C7—C2	-0.4 (4)	C24—C25—C26—C21	-1.2 (3)
C5—C6—C7—C8	-179.4 (2)	C22—C21—C26—C25	1.5 (3)
C2—C7—C8—C9	178.4 (2)	C20—C21—C26—C25	-179.01 (17)
C6—C7—C8—C9	-2.5 (3)	C22—C21—C26—N2	-178.02 (17)
C2—C7—C8—C13	-1.5 (3)	C20—C21—C26—N2	1.4 (3)
C6—C7—C8—C13	177.6 (2)	C2—C1—N1—C13	-1.3 (3)
C13—C8—C9—C10	-0.4 (3)	C2—C1—N1—Zn1	168.86 (16)
C7—C8—C9—C10	179.7 (2)	C12—C13—N1—C1	-179.80 (18)
C8—C9—C10—C11	-0.3 (4)	C8—C13—N1—C1	0.7 (3)
C9—C10—C11—C12	0.2 (4)	C12—C13—N1—Zn1	10.7 (2)
C10—C11—C12—C13	0.8 (3)	C8—C13—N1—Zn1	-168.77 (14)
C11—C12—C13—N1	179.0 (2)	C15—C14—N2—C26	-2.6 (3)
C11—C12—C13—C8	-1.5 (3)	C15—C14—N2—Zn1	-177.82 (14)
C9—C8—C13—C12	1.3 (3)	C25—C26—N2—C14	-178.37 (18)
C7—C8—C13—C12	-178.80 (18)	C21—C26—N2—C14	1.2 (3)
C9—C8—C13—N1	-179.20 (18)	C25—C26—N2—Zn1	-3.3 (2)
C7—C8—C13—N1	0.7 (3)	C21—C26—N2—Zn1	176.23 (13)
N2—C14—C15—C20	1.2 (3)	C14—N2—Zn1—N1	-15.81 (16)
N2—C14—C15—C16	-177.53 (18)	C26—N2—Zn1—N1	169.13 (14)
C20—C15—C16—C17	-0.4 (3)	C14—N2—Zn1—Cl1	97.39 (14)
C14—C15—C16—C17	178.30 (19)	C26—N2—Zn1—Cl1	-77.67 (14)
C15—C16—C17—C18	0.0 (3)	C14—N2—Zn1—Cl2	-133.03 (13)
C16—C17—C18—C19	0.2 (4)	C26—N2—Zn1—Cl2	51.91 (15)
C17—C18—C19—C20	0.1 (3)	C1—N1—Zn1—N2	121.89 (15)
C16—C15—C20—C19	0.7 (3)	C13—N1—Zn1—N2	-68.46 (16)
C14—C15—C20—C19	-178.05 (18)	C1—N1—Zn1—Cl1	7.32 (16)
C16—C15—C20—C21	-179.75 (17)	C13—N1—Zn1—Cl1	176.97 (14)
C14—C15—C20—C21	1.5 (3)	C1—N1—Zn1—Cl2	-116.83 (14)
C18—C19—C20—C15	-0.6 (3)	C13—N1—Zn1—Cl2	52.82 (15)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...Cl1	0.93	2.77	3.434 (3)	129
C17—H17...Cg6 ⁱ	0.93	2.82	3.535 (3)	134

supplementary materials

C24—H24 \cdots Cg5ⁱⁱ

0.93

2.81

3.508 (3)

132

Symmetry codes: (i) $-x, y-1/2, -z-1/2$; (ii) $x, -y-1/2, z-1/2$.

Fig. 1




Fig. 2

