

Diaquabis(1,3-propanediamine)nickel(II) squarate tetrahydrate

Ersin Temel,^a Hakan Erer,^b Okan Zafer Yeşilel^b and Orhan Büyükgüngör^{a*}

^aDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, Kurupelit, TR-55139, Samsun, Turkey, and ^bDepartment of Chemistry, Faculty of Arts and Sciences, Eskişehir Osmangazi University, TR-26480, Eskişehir, Turkey

Correspondence e-mail: orhanb@omu.edu.tr


Received 28 May 2009; accepted 2 June 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.021; wR factor = 0.054; data-to-parameter ratio = 15.9.

The asymmetric unit of the title compound, $[\text{Ni}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2](\text{C}_4\text{O}_4)\cdot 4\text{H}_2\text{O}$, contains one-half of the diaquabis(1,3-propanediamine)nickel(II) cation, one-half of the centrosymmetric squarate anion and two uncoordinated water molecules. In the cation, the Ni^{II} atom is located on a crystallographic inversion centre and has a slightly distorted octahedral coordination geometry. The six-membered chelate ring adopts a chair conformation. $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the cation and anion through the water molecule, while $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the cation and anion and cation and water molecules. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into a three-dimensional network structure.

Related literature

For general background, see: Bertolasi *et al.* (2001); Gollogly & Hawkins (1972); Lam & Mak (2000); Liebeskind *et al.* (1993); Mathew *et al.* (2002); Reetz *et al.* (1994); Seitz & Imming (1992); Zaman *et al.* (2001). For related structures, see: Ghosh *et al.* (1997); Mukherjee *et al.* (1990); Pariya *et al.* (1995). For ring-puckering parameters, see: Cremer & Pople (1975). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$[\text{Ni}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2](\text{C}_4\text{O}_4)\cdot 4\text{H}_2\text{O}$ $V = 958.40$ (9) Å³
 $M_r = 427.09$ $Z = 2$
 Monoclinic, $P2_1/c$ $\text{Mo } K\alpha$ radiation
 $a = 8.0429$ (4) Å $\mu = 1.06$ mm⁻¹
 $b = 9.1752$ (5) Å $T = 296$ K
 $c = 14.6510$ (8) Å $0.75 \times 0.45 \times 0.05$ mm
 $\beta = 117.570$ (4)°

Data collection

Stoe IPDS II diffractometer 7288 measured reflections
 Absorption correction: integration 2204 independent reflections
 ($X\text{-RED32}$; Stoe & Cie, 2002) 2003 reflections with $I > 2\sigma(I)$
 $T_{\text{min}} = 0.638$, $T_{\text{max}} = 0.949$ $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.054$ $\Delta\rho_{\text{max}} = 0.29$ e Å⁻³
 $S = 1.06$ $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³
 2204 reflections
 139 parameters

Table 1

Selected geometric parameters (Å, °).

O1—Ni1	2.1429 (9)	N2—Ni1	2.0997 (10)
N1—Ni1	2.1090 (10)		
N1—Ni1—O1	88.86 (4)	N2—Ni1—N1	91.94 (4)
N2—Ni1—O1	91.46 (4)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1B \cdots O5 ⁱ	0.90	2.35	3.1715 (16)	152
N2—H2A \cdots O2 ⁱⁱ	0.90	2.33	3.2174 (14)	170
O1—H1F \cdots O2 ⁱⁱⁱ	0.77 (2)	2.08 (2)	2.8345 (15)	165 (2)
O4—H4A \cdots O2 ⁱⁱⁱ	0.80 (2)	2.10 (2)	2.8765 (16)	165 (2)
O4—H4B \cdots O2 ^{iv}	0.82 (3)	2.07 (3)	2.8965 (16)	178 (2)
O5—H5B \cdots O4 ^v	0.77 (2)	2.10 (3)	2.8730 (19)	177 (2)
N1—H1A \cdots O4	0.90	2.38	3.2442 (17)	160
N2—H2B \cdots O3	0.90	2.04	2.9333 (14)	174
O1—H1E \cdots O5	0.80 (2)	1.93 (2)	2.7311 (16)	175.8 (19)
O5—H5A \cdots O3	0.80 (2)	1.94 (2)	2.7296 (16)	166 (2)

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $x, y - 1, z$; (iv) $-x + 1, -y + 1, -z + 1$; (v) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences of Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS II diffractometer (purchased under grant No. F279 of the University Research Grant of Ondokuz Mayıs University).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2703).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bertolasi, V., Gilli, P., Ferretti, V. & Gilli, G. (2001). *Acta Cryst. B57*, 591–598.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Ghosh, S., Mukherjee, M., Mukherjee, A. K. & Ray Chaudhuri, N. (1997). *Acta Cryst. C53*, 1561–1564.
- Gollogly, J. R. & Hawkins, C. J. (1972). *Inorg. Chem.* **11**, 156–161.
- Lam, C. K. & Mak, T. C. W. (2000). *Tetrahedron*, **56**, 6657–6665.
- Liebeskind, L. S., Yu, M. S., Yu, R. H., Wang, J. & Hagen, K. S. (1993). *J. Am. Chem. Soc.* **115**, 9048–9050.
- Mathew, S., Paul, G., Shivasankar, K., Choudhury, A. & Rao, C. N. R. (2002). *J. Mol. Struct.* **641**, 263–279.
- Mukherjee, A. K., Mukherjee, M., Ray, S., Ghosh, A. & Chaudhuri, N. R. (1990). *J. Chem. Soc. Dalton Trans.* pp. 2347–2350.
- Pariya, C., Ghosh, S., Ghosh, A., Mukherjee, M., Mukherjee, A. K. & Chaudhuri, N. R. (1995). *J. Chem. Soc. Dalton Trans.* pp. 337–342.
- Reetz, M. T., Hoger, S. & Harms, K. (1994). *Angew. Chem. Int. Ed. Engl.* **33**, 181–183.
- Seitz, G. & Imming, P. (1992). *Chem. Rev.* **92**, 1227–1260.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Zaman, M. B., Tomura, M. & Yamashita, Y. (2001). *Acta Cryst. C57*, 621–624.

supplementary materials

Acta Cryst. (2009). E65, m735-m736 [doi:10.1107/S160053680902087X]

Diaquabis(1,3-propanediamine)nickel(II) squarate tetrahydrate

E. Temel, H. Erer, O. Z. Yesilel and O. Büyükgüngör

Comment

The conformation of six-membered rings arranged by the bidentate coordination of pen (1,3-propanediamine) to transition metals has long been of theoretical interest (Gollgoly & Hawkins, 1972). Despite this interest, only a limited number of such complexes have been structurally described. Because of their ability to undergo solid-state phase transitions, some nickel(II) complexes of bis(N-substituted-pen) have been studied in recent times (Mukherjee *et al.*, 1990; Pariya *et al.*, 1995; Ghosh *et al.*, 1997).

Squaric acid (H₂C₄O₄) has been of much interest because of its cyclic structure and possible aromaticity. Recently, considerable progress has been made in the crystal engineering of multidimensional arrays and networks containing metal ions as nodes. Squaric acid is a useful tool for constructing crystalline architectures, due to its rigid, planar four membered ring skeleton, and its proton donating and accepting capabilities for hydrogen bonding (Bertolasi *et al.*, 2001; Reetz *et al.*, 1994; Lam & Mak, 2000; Zaman *et al.*, 2001; Mathew *et al.*, 2002). In addition, squaric acid has been studied for potetial application in xerographic photoreceptors, organic solar cells and optical recording (Liebeskind *et al.*, 1993; Seitz & Imming, 1992).

The asymmetric unit of the title compound contains one centrosymmetric cation, where Ni^{II} is located on a crystallographic inversion centre, one centrosymmetric anion and two uncoordinated water molecules (Fig.1), in which the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. In cation, the Ni^{II} is hexacoordinated by two O atoms of two water molecules in a *trans* order and by four N atoms of two pen ligands at the equatorial positions (Table 1). It is suggested that the *trans* geometry is preferred, when the amine ligand is more bulky. Thus, the coordination environment of Ni^{II} is a slightly distorted octahedral. Intramolecular O-H...O hydrogen bonds (Table 2) link the cation and anion through the water molecule, while intramolecular N-H...O hydrogen bonds (Table 2) link the cation and anion and cation and water molecule. The six-membered chelate ring (Ni1/N1/N2/C1-C3) is not planar, having total puckering amplitude, Q_T, of 0.765 (3) Å and chair conformation [$\varphi = 166.25 (3)$ and $\theta = 40.42 (3)^\circ$] (Cremer & Pople, 1975).

In the crystal structure, intermolecular O-H...O and N-H...O hydrogen bonds (Table 2) link the molecules into chains (Fig. 2), in which they may be effective in the stabilization of the structure.

Experimental

For the preparation of the title compound, a solution of squaric acid (0.57 g, 5 mmol) in water (25 ml) was neutralized with sodium hydroxide (0.40 g, 10 mmol) and added dropwise with stirring to a solution of Ni(CH₃COO)₂·4(H₂O) (1.24 g, 5 mmol) in water (25 ml) at 323 K. The solution immediately became suspension and was stirred for 2 h. Then, 1,3-propanediamine (0.74 g, 10 mmol) in methanol (10 ml) was added dropwise to the obtained suspension. The clear solution was stirred for 2 h, and then cooled to room temperature. The crystals formed were filtered and washed with water (10 ml) and methanol (1:1), then dried in air. Anal. Calcd. : C 28.12, H 7.55, N 13.12%; Found C 28.06, H 7.61, N 13.18%.

Refinement

Atoms H1E, H1F, H4A, H4B, H5A and H5B (for H₂O) were located in difference syntheses and refined isotropically. The remaining H atoms were positioned geometrically with N-H = 0.90 Å (for NH₂) and C-H = 0.97 Å (for CH₂) and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 40% probability [symmetry code: (i) 1 - x, 1 - y, 1 - z].




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity [symmetry code: (i) 1 - x, 1 - y, 1 - z].

Diaquabis(1,3-propanediamine)nickel(II) squarate tetrahydrate

Crystal data

$[\text{Ni}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2](\text{C}_4\text{O}_4)\cdot 4\text{H}_2\text{O}$

$M_r = 427.09$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.0429$ (4) Å

$b = 9.1752$ (5) Å

$c = 14.6510$ (8) Å

$\beta = 117.570$ (4)°

$V = 958.40$ (9) Å³

$Z = 2$

$F_{000} = 456.0$

$D_x = 1.480$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2204 reflections

$\theta = 2.2\text{--}28.0^\circ$

$\mu = 1.06$ mm⁻¹

$T = 296$ K

Plate, violet

$0.75 \times 0.45 \times 0.05$ mm

Data collection

Stoe IPDS II
diffractometer

Monochromator: plane graphite

Detector resolution: 6.67 pixels mm⁻¹

$T = 296$ K

ω -scan rotation method

Absorption correction: integration
(X-RED32; Stoe & Cie, 2002)

2204 independent reflections

2003 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.020$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 2.7^\circ$

$h = -10 \rightarrow 10$

$T_{\min} = 0.638$, $T_{\max} = 0.949$
7288 measured reflections

$k = -11 \rightarrow 11$
 $l = -19 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.054$	$w = 1/[\sigma^2(F_o^2) + (0.0282P)^2 + 0.2118P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2204 reflections	$(\Delta/\sigma)_{\max} < 0.001$
139 parameters	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.5000	0.5000	0.02232 (7)
O1	0.64732 (15)	0.39000 (12)	0.43015 (8)	0.0383 (2)
H1E	0.713 (3)	0.432 (2)	0.4113 (14)	0.051 (5)*
H1F	0.692 (3)	0.315 (3)	0.4515 (15)	0.063 (6)*
O2	0.73874 (13)	1.09844 (11)	0.49933 (8)	0.0414 (2)
O3	0.86500 (14)	0.77497 (10)	0.46843 (9)	0.0434 (2)
O4	0.34274 (18)	0.07964 (14)	0.36302 (9)	0.0489 (3)
H4A	0.455 (3)	0.082 (2)	0.3910 (16)	0.067 (6)*
H4B	0.317 (3)	0.029 (3)	0.401 (2)	0.075 (7)*
O5	0.85562 (18)	0.53525 (13)	0.35580 (11)	0.0462 (3)
H5A	0.871 (3)	0.611 (3)	0.3859 (17)	0.066 (6)*
H5B	0.803 (3)	0.550 (3)	0.2975 (19)	0.072 (8)*
N1	0.24635 (14)	0.41813 (12)	0.38210 (8)	0.0317 (2)
H1A	0.2548	0.3203	0.3845	0.038*
H1B	0.1547	0.4421	0.3983	0.038*

supplementary materials

N2	0.47942 (14)	0.69084 (11)	0.41632 (8)	0.0294 (2)
H2A	0.4236	0.7588	0.4372	0.035*
H2B	0.5970	0.7221	0.4355	0.035*
C1	0.1833 (2)	0.46195 (17)	0.27461 (10)	0.0423 (3)
H1C	0.0568	0.4263	0.2326	0.051*
H1D	0.2640	0.4173	0.2497	0.051*
C2	0.1860 (2)	0.62587 (16)	0.26262 (11)	0.0438 (3)
H2C	0.1144	0.6502	0.1902	0.053*
H2D	0.1242	0.6710	0.2987	0.053*
C3	0.3806 (2)	0.68931 (16)	0.30294 (10)	0.0406 (3)
H3A	0.4520	0.6320	0.2776	0.049*
H3B	0.3718	0.7880	0.2774	0.049*
C4	0.88219 (16)	1.04411 (13)	0.49952 (9)	0.0269 (2)
C5	0.93844 (16)	0.89777 (13)	0.48539 (9)	0.0271 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.02176 (11)	0.02133 (11)	0.02386 (11)	0.00055 (7)	0.01055 (8)	0.00121 (7)
O1	0.0460 (6)	0.0308 (5)	0.0522 (6)	0.0070 (4)	0.0346 (5)	0.0041 (4)
O2	0.0319 (5)	0.0362 (5)	0.0643 (7)	0.0049 (4)	0.0291 (5)	0.0006 (5)
O3	0.0453 (5)	0.0286 (5)	0.0636 (7)	-0.0116 (4)	0.0314 (5)	-0.0089 (4)
O4	0.0416 (6)	0.0539 (7)	0.0491 (6)	-0.0045 (5)	0.0191 (5)	0.0039 (5)
O5	0.0530 (7)	0.0407 (6)	0.0538 (7)	-0.0009 (5)	0.0321 (6)	-0.0024 (5)
N1	0.0279 (5)	0.0315 (5)	0.0311 (5)	-0.0026 (4)	0.0097 (4)	-0.0008 (4)
N2	0.0309 (5)	0.0258 (5)	0.0317 (5)	0.0005 (4)	0.0147 (4)	0.0031 (4)
C1	0.0455 (8)	0.0434 (7)	0.0282 (6)	-0.0033 (6)	0.0089 (6)	-0.0053 (5)
C2	0.0453 (8)	0.0452 (8)	0.0278 (6)	0.0055 (6)	0.0058 (6)	0.0062 (6)
C3	0.0517 (8)	0.0407 (7)	0.0322 (6)	0.0026 (6)	0.0219 (6)	0.0086 (5)
C4	0.0251 (5)	0.0269 (5)	0.0296 (6)	0.0008 (4)	0.0134 (4)	0.0015 (4)
C5	0.0269 (5)	0.0263 (5)	0.0290 (6)	-0.0021 (4)	0.0138 (4)	-0.0006 (4)

Geometric parameters (\AA , $^\circ$)

Ni1—O1 ⁱ	2.1429 (9)	C1—N1	1.4695 (17)
Ni1—N1 ⁱ	2.1090 (10)	C1—C2	1.516 (2)
Ni1—N2 ⁱ	2.0997 (10)	C1—H1C	0.9700
O1—Ni1	2.1429 (9)	C1—H1D	0.9700
O1—H1E	0.80 (2)	C2—C3	1.511 (2)
O1—H1F	0.77 (2)	C2—H2C	0.9700
O4—H4A	0.80 (2)	C2—H2D	0.9700
O4—H4B	0.82 (3)	C3—N2	1.4728 (16)
O5—H5A	0.80 (2)	C3—H3A	0.9700
O5—H5B	0.77 (2)	C3—H3B	0.9700
N1—Ni1	2.1090 (10)	C4—O2	1.2557 (15)
N1—H1A	0.9000	C4—C5 ⁱⁱ	1.4557 (16)
N1—H1B	0.9000	C4—C5	1.4619 (17)
N2—Ni1	2.0997 (10)	C5—O3	1.2427 (15)

N2—H2A	0.9000	C5—C4 ⁱⁱ	1.4557 (16)
N2—H2B	0.9000		
O1—Ni1—O1 ⁱ	180.00 (5)	C3—N2—Ni1	120.41 (8)
N1 ⁱ —Ni1—N1	180.0	C3—N2—H2A	107.2
N1 ⁱ —Ni1—O1	91.14 (4)	C3—N2—H2B	107.2
N1—Ni1—O1	88.86 (4)	H2A—N2—H2B	106.9
N1 ⁱ —Ni1—O1 ⁱ	88.86 (4)	N1—C1—C2	112.30 (11)
N1—Ni1—O1 ⁱ	91.14 (4)	N1—C1—H1C	109.1
N2 ⁱ —Ni1—O1	88.54 (4)	C2—C1—H1C	109.1
N2—Ni1—O1	91.46 (4)	N1—C1—H1D	109.1
N2 ⁱ —Ni1—O1 ⁱ	91.46 (4)	C2—C1—H1D	109.1
N2—Ni1—O1 ⁱ	88.54 (4)	H1C—C1—H1D	107.9
N2 ⁱ —Ni1—N2	180.0	C3—C2—C1	113.88 (12)
N2 ⁱ —Ni1—N1 ⁱ	91.94 (4)	C3—C2—H2C	108.8
N2—Ni1—N1 ⁱ	88.06 (4)	C1—C2—H2C	108.8
N2 ⁱ —Ni1—N1	88.06 (4)	C3—C2—H2D	108.8
N2—Ni1—N1	91.94 (4)	C1—C2—H2D	108.8
Ni1—O1—H1E	122.4 (14)	H2C—C2—H2D	107.7
Ni1—O1—H1F	118.8 (15)	N2—C3—C2	111.42 (11)
H1E—O1—H1F	108.4 (19)	N2—C3—H3A	109.3
H4A—O4—H4B	104 (2)	C2—C3—H3A	109.3
H5A—O5—H5B	109 (2)	N2—C3—H3B	109.3
Ni1—N1—H1A	107.3	C2—C3—H3B	109.3
Ni1—N1—H1B	107.3	H3A—C3—H3B	108.0
C1—N1—Ni1	120.23 (9)	O2—C4—C5 ⁱⁱ	134.42 (12)
C1—N1—H1A	107.3	O2—C4—C5	135.11 (12)
C1—N1—H1B	107.3	C5 ⁱⁱ —C4—C5	90.46 (9)
H1A—N1—H1B	106.9	O3—C5—C4 ⁱⁱ	135.11 (12)
Ni1—N2—H2A	107.2	O3—C5—C4	135.35 (12)
Ni1—N2—H2B	107.2	C4 ⁱⁱ —C5—C4	89.54 (9)
C1—N1—Ni1—O1	-63.90 (10)	N1—C1—C2—C3	72.93 (17)
C1—N1—Ni1—O1 ⁱ	116.10 (10)	C1—C2—C3—N2	-73.63 (16)
C1—N1—Ni1—N2 ⁱ	-152.47 (10)	C2—C3—N2—Ni1	52.35 (14)
C1—N1—Ni1—N2	27.53 (10)	O2—C4—C5—O3	-0.2 (3)
C3—N2—Ni1—O1	60.31 (10)	C5 ⁱⁱ —C4—C5—O3	179.50 (19)
C3—N2—Ni1—O1 ⁱ	-119.69 (10)	O2—C4—C5—C4 ⁱⁱ	-179.72 (18)
C2—C1—N1—Ni1	-50.38 (16)	C5 ⁱⁱ —C4—C5—C4 ⁱⁱ	0.0

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1B \cdots O5 ⁱⁱⁱ	0.90	2.35	3.1715 (16)	152
N2—H2A \cdots O2 ^{iv}	0.90	2.33	3.2174 (14)	170

supplementary materials

O1—H1F···O2 ^v	0.77 (2)	2.08 (2)	2.8345 (15)	165 (2)
O4—H4A···O2 ^v	0.80 (2)	2.10 (2)	2.8765 (16)	165 (2)
O4—H4B···O2 ⁱ	0.82 (3)	2.07 (3)	2.8965 (16)	178 (2)
O5—H5B···O4 ^{vi}	0.77 (2)	2.10 (3)	2.8730 (19)	177 (2)
N1—H1A···O4	0.90	2.38	3.2442 (17)	160
N2—H2B···O3	0.90	2.04	2.9333 (14)	174
O1—H1E···O5	0.80 (2)	1.93 (2)	2.7311 (16)	175.8 (19)
O5—H5A···O3	0.80 (2)	1.94 (2)	2.7296 (16)	166 (2)

Symmetry codes: (iii) $x-1, y, z$; (iv) $-x+1, -y+2, -z+1$; (v) $x, y-1, z$; (i) $-x+1, -y+1, -z+1$; (vi) $-x+1, y+1/2, -z+1/2$.

Fig. 1




Fig. 2

