

catena-Poly[[ethanol- κ O)sodium(I)]-di- μ -aqua-[(*rac*-2'-hydroxy-1,1'-binaphthyl-2-yl phosphato- κ O)sodium]-tri- μ -aqua]

Yuya Tachibana* and Yasukatsu Maeda

Industrial Technology Center of Wakayama Prefecture, 60 Ogura, Wakayama 649-6261, Japan

Correspondence e-mail: yuya-tachibana@aist.go.jp


Received 28 May 2009; accepted 5 June 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.114; data-to-parameter ratio = 17.9.

The asymmetric unit of the polymeric title compound, $[\text{Na}_2(\text{C}_{20}\text{H}_{13}\text{O}_5\text{P})(\text{C}_2\text{H}_6\text{O})(\text{H}_2\text{O})_5]_n$, consists of two Na^{I} ions, one 2'-hydroxy-1,1'-binaphthyl-2-yl phosphate anion, one ethanol ligand and five water molecules of crystallization. Each Na^{I} ion has a distorted octahedral coordination geometry. The phosphate anion coordinates to one Na^{I} ion and the ethanol molecule coordinates to the other. The five water molecules bridge the Na^{I} ions, forming an inorganic chain structure along the b axis. The chains are connected by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds into an organic-inorganic hybrid layer parallel to (001).

Related literature

For organic-inorganic hybrid materials, see: Eckert & Ward (2001). For phosphate derivatives, see: Vioux *et al.* (2004).



Experimental

Crystal data

 $[\text{Na}_2(\text{C}_{20}\text{H}_{13}\text{O}_5\text{P})(\text{C}_2\text{H}_6\text{O})(\text{H}_2\text{O})_5]$
 $M_r = 546.40$

 Monoclinic, $P2_1/a$
 $a = 13.121$ (4) Å

 $b = 9.816$ (3) Å
 $c = 20.198$ (7) Å
 $\beta = 100.033$ (13)°
 $V = 2561.7$ (14) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 293$ K
 $0.60 \times 0.50 \times 0.10$ mm

Data collection

 Rigaku R-AXIS RAPID
 diffractometer
 Absorption correction: multi-scan
 (ABSCOR; Higashi, 1995)
 $T_{\text{min}} = 0.874$, $T_{\text{max}} = 0.980$

 24950 measured reflections
 5829 independent reflections
 5045 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.114$
 $S = 1.05$
 5829 reflections
 326 parameters
 12 restraints

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.46$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{O4}$	0.847 (17)	1.839 (17)	2.6687 (19)	166.3 (18)
$\text{O6}-\text{H2}\cdots\text{O11}^{\text{i}}$	0.852 (16)	2.043 (17)	2.854 (2)	159.0 (16)
$\text{O6}-\text{H3}\cdots\text{O9}^{\text{i}}$	0.848 (16)	1.956 (16)	2.7958 (19)	170.5 (16)
$\text{O7}-\text{H4}\cdots\text{O5}^{\text{ii}}$	0.853 (11)	1.873 (10)	2.7136 (18)	168.4 (19)
$\text{O7}-\text{H5}\cdots\text{O4}^{\text{i}}$	0.851 (14)	2.106 (14)	2.9473 (19)	169.9 (18)
$\text{O8}-\text{H6}\cdots\text{O1}^{\text{i}}$	0.847 (16)	1.955 (16)	2.775 (2)	162.7 (16)
$\text{O9}-\text{H8}\cdots\text{O4}$	0.852 (15)	1.875 (15)	2.7259 (18)	177.5 (12)
$\text{O9}-\text{H9}\cdots\text{O3}^{\text{iii}}$	0.851 (12)	1.850 (12)	2.6978 (18)	174.1 (16)
$\text{O10}-\text{H10}\cdots\text{O5}^{\text{iv}}$	0.850 (15)	2.197 (15)	3.012 (2)	161 (2)
$\text{O10}-\text{H11}\cdots\text{O3}^{\text{i}}$	0.849 (12)	2.197 (9)	2.984 (2)	154.1 (19)
$\text{O11}-\text{H12}\cdots\text{O3}^{\text{iv}}$	0.851 (9)	1.823 (11)	2.6711 (19)	174 (2)

 Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z$; (iii) $-x + 1, -y, -z$; (iv) $x, y + 1, z$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2004); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2427).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
 Eckert, H. & Ward, M. (2001). *Chem. Mater.* **13**, 3059–3060.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2004). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vioux, A., Bideau, J.-L., Mutin, P. H. & Leclercq, D. (2004). *Top. Curr. Chem.* **232**, 145–174.

supplementary materials

Acta Cryst. (2009). E65, m753 [doi:10.1107/S1600536809021394]

***catena*-Poly[[*(ethanol-κO)*sodium(I)]-di-*μ*-aqua-[(*rac*-2'-hydroxy-1,1'-binaphthyl-2-yl phosphato-*κO*)sodium]-tri-*μ*-aqua]**

Y. Tachibana and Y. Maeda

Comment

In recent years, studies on organic-inorganic hybrid layered materials have received a great deal of attention as electrical, magnetic and/or optical materials (Eckert & Ward, 2001). Some combinations are studied to form layered structure. Phosphate ester can form organic-inorganic hybrid network (Vioux *et al.*, 2004).

The title compound has inorganic layer formed with two sodium cation, phosphate anion and five waters, and organic layer formed by binaphthyl unit. The phosphate anion coordinates to one Na^I atom and the ethanol molecule coordinates to the other Na^I atom as shown in Fig. 1. Sodium ions in the compound are bridged by three water molecules, and one sodium ion connected to adjoining sodium ion through two water molecules to form a one-dimensional zigzag chain structure along the *b* axis as shown in Fig. 2. The chains build two-dimensional network along the *a-b* plane by hydrogen bonding, and form inorganic layer. Organic layers, build up with binaphthyl moiety, and inorganic layers form bilayered structure in the crystal. Then, three-dimensional organic-inorganic hybrid layered material is formed by self-assembly as shown in Fig. 3.

Experimental

Sodium hydride (1.5 g, 63 mmol) was dispersed into tetrahydrofuran (50 ml). (*rac*)-1,1'-Binaphthyl-2,2'-diyl hydrogenphosphate (1.0 g, 2.9 mmol) was added to the dispersion and stirred for 2 h at room temperature. The resulting mixture was filtrated and evaporated *in vacuo*. The residue was recrystallized from H₂O/ethanol.

Refinement

All C-bound H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and refined as riding atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. O-bound H atoms were located in a difference Fourier map and the positional parameters were refined, with distance restraints of O—H = 0.850 (5) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. A view of the title compound, showing 50% probability displacement ellipsoids.




Fig. 2. The one dimensional chain structure of the title compound bridged between sodium cations and water molecules. Hydrogen atoms have been omitted for clarity.




Fig. 3. The layered structure of the title compound. Hydrogen atoms have been omitted for clarity.

catena-Poly[[*(ethanol-κO)sodium(I)*]-di- μ -aqua- [*(rac-2'-hydroxy-1,1'-binaphthyl-2-yl phosphato-κO)sodium*]-tri- μ -aqua]

Crystal data

[Na₂(C₂₀H₁₃O₅P)(C₂H₆O)(H₂O)₅]

$M_r = 546.40$

Monoclinic, $P2_1/a$

Hall symbol: -P 2yab

$a = 13.121$ (4) Å

$b = 9.816$ (3) Å

$c = 20.198$ (7) Å

$\beta = 100.033$ (13)°

$V = 2561.7$ (14) Å³

$Z = 4$

$F_{000} = 1144$

$D_x = 1.417$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71075$ Å

Cell parameters from 19717 reflections

$\theta = 3.1$ – 27.5 °

$\mu = 0.20$ mm⁻¹

$T = 293$ K

Platelet, colourless

$0.60 \times 0.50 \times 0.10$ mm

Data collection

Rigaku R-Axis RAPID
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 5.00 pixels mm⁻¹

$T = 293$ K

ω scans

Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.874$, $T_{\max} = 0.980$

24950 measured reflections

5829 independent reflections

5045 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\max} = 27.5$ °

$\theta_{\min} = 3.1$ °

$h = -16 \rightarrow 16$

$k = -12 \rightarrow 12$

$l = -26 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.114$	$w = 1/[\sigma^2(F_o^2) + (0.0682P)^2 + 0.7208P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
5829 reflections	$(\Delta/\sigma)_{\max} = 0.006$
326 parameters	$\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
12 restraints	$\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.57919 (3)	-0.04823 (4)	0.124182 (17)	0.02333 (10)
Na1	0.66932 (5)	0.48279 (7)	0.07453 (3)	0.03971 (17)
Na2	0.74134 (4)	0.17009 (6)	0.04642 (3)	0.03330 (15)
O1	0.40206 (9)	0.09598 (13)	0.21979 (6)	0.0394 (3)
H1	0.4489 (12)	0.098 (2)	0.1958 (9)	0.047*
O2	0.61673 (8)	-0.09564 (10)	0.20212 (5)	0.0282 (2)
O3	0.50549 (8)	-0.16097 (12)	0.09628 (5)	0.0350 (2)
O4	0.52559 (9)	0.08848 (11)	0.12794 (5)	0.0338 (2)
O5	0.67384 (8)	-0.03767 (11)	0.09123 (5)	0.0318 (2)
O6	0.89693 (9)	0.06153 (12)	0.03525 (6)	0.0361 (3)
H2	0.9272 (14)	0.0031 (16)	0.0631 (8)	0.043*
H3	0.9421 (12)	0.1229 (15)	0.0342 (10)	0.043*
O7	0.82327 (8)	0.39178 (13)	0.03833 (6)	0.0352 (2)
H4	0.8329 (15)	0.410 (2)	-0.0014 (4)	0.042*
H5	0.8815 (8)	0.408 (2)	0.0629 (9)	0.042*
O8	0.72026 (10)	0.30040 (13)	0.14616 (6)	0.0392 (3)

supplementary materials

H6	0.7723 (11)	0.322 (2)	0.1753 (8)	0.047*
H7	0.6828 (13)	0.2551 (19)	0.1684 (9)	0.047*
O9	0.56093 (8)	0.26022 (11)	0.02832 (6)	0.0326 (2)
H8	0.5482 (14)	0.2058 (16)	0.0587 (7)	0.039*
H9	0.5390 (14)	0.2237 (19)	-0.0097 (5)	0.039*
O10	0.77621 (10)	0.69039 (13)	0.08084 (7)	0.0434 (3)
H10	0.7540 (16)	0.7654 (13)	0.0938 (11)	0.052*
H11	0.8397 (6)	0.685 (2)	0.0986 (11)	0.052*
O11	0.53790 (10)	0.58034 (12)	0.13535 (7)	0.0420 (3)
H12	0.5281 (17)	0.6642 (8)	0.1257 (11)	0.050*
C1	0.40429 (11)	-0.01837 (17)	0.25947 (8)	0.0318 (3)
C2	0.30914 (12)	-0.0869 (2)	0.25809 (9)	0.0424 (4)
H13	0.2496	-0.0541	0.2310	0.051*
C3	0.30408 (13)	-0.2003 (2)	0.29597 (9)	0.0437 (4)
H14	0.2409	-0.2441	0.2946	0.052*
C4	0.39363 (13)	-0.25268 (17)	0.33747 (8)	0.0350 (3)
C5	0.39111 (16)	-0.37427 (19)	0.37466 (9)	0.0443 (4)
H15	0.3285	-0.4192	0.3738	0.053*
C6	0.47905 (18)	-0.4266 (2)	0.41165 (10)	0.0522 (5)
H16	0.4769	-0.5089	0.4342	0.063*
C7	0.57301 (17)	-0.3559 (2)	0.41573 (10)	0.0518 (5)
H17	0.6325	-0.3908	0.4421	0.062*
C8	0.57852 (13)	-0.23620 (18)	0.38149 (8)	0.0387 (4)
H18	0.6412	-0.1901	0.3855	0.046*
C9	0.48927 (11)	-0.18220 (16)	0.34002 (7)	0.0296 (3)
C10	0.49298 (11)	-0.06224 (15)	0.30036 (7)	0.0273 (3)
C11	0.59188 (10)	0.01444 (15)	0.30256 (7)	0.0266 (3)
C12	0.62733 (11)	0.10678 (16)	0.35622 (7)	0.0314 (3)
C13	0.57337 (15)	0.1248 (2)	0.41085 (9)	0.0479 (4)
H19	0.5130	0.0760	0.4119	0.057*
C14	0.60945 (19)	0.2133 (3)	0.46178 (11)	0.0629 (6)
H20	0.5742	0.2225	0.4977	0.076*
C15	0.6995 (2)	0.2907 (3)	0.46050 (11)	0.0657 (6)
H21	0.7224	0.3520	0.4950	0.079*
C16	0.75278 (17)	0.2760 (2)	0.40923 (11)	0.0552 (5)
H22	0.8122	0.3276	0.4090	0.066*
C17	0.71954 (13)	0.18328 (17)	0.35570 (8)	0.0366 (3)
C18	0.77548 (12)	0.16267 (18)	0.30267 (9)	0.0378 (4)
H23	0.8356	0.2124	0.3018	0.045*
C19	0.74264 (11)	0.07110 (16)	0.25277 (8)	0.0324 (3)
H24	0.7815	0.0563	0.2191	0.039*
C20	0.64935 (10)	-0.00130 (14)	0.25234 (7)	0.0254 (3)
C21	0.50869 (18)	0.5732 (2)	0.20062 (11)	0.0533 (5)
H25A	0.4374	0.6027	0.1974	0.064*
H25B	0.5520	0.6343	0.2311	0.064*
C22	0.5196 (2)	0.4332 (3)	0.22760 (13)	0.0656 (6)
H26A	0.4797	0.3719	0.1963	0.098*
H26B	0.4952	0.4298	0.2697	0.098*
H26C	0.5912	0.4069	0.2344	0.098*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.02712 (18)	0.02280 (18)	0.02071 (17)	0.00115 (12)	0.00595 (13)	0.00013 (12)
Na1	0.0399 (3)	0.0389 (4)	0.0434 (4)	0.0078 (3)	0.0157 (3)	0.0124 (3)
Na2	0.0303 (3)	0.0315 (3)	0.0398 (3)	0.0013 (2)	0.0108 (2)	-0.0015 (2)
O1	0.0335 (6)	0.0470 (7)	0.0395 (6)	0.0102 (5)	0.0115 (5)	0.0121 (5)
O2	0.0382 (5)	0.0252 (5)	0.0213 (4)	0.0023 (4)	0.0055 (4)	0.0011 (4)
O3	0.0374 (6)	0.0348 (6)	0.0310 (5)	-0.0085 (5)	0.0014 (4)	-0.0008 (4)
O4	0.0436 (6)	0.0302 (5)	0.0298 (5)	0.0115 (4)	0.0129 (4)	0.0058 (4)
O5	0.0348 (5)	0.0337 (6)	0.0298 (5)	0.0000 (4)	0.0140 (4)	-0.0013 (4)
O6	0.0301 (5)	0.0311 (6)	0.0471 (7)	0.0012 (4)	0.0069 (5)	0.0030 (5)
O7	0.0312 (5)	0.0442 (7)	0.0317 (5)	-0.0035 (5)	0.0094 (4)	0.0010 (5)
O8	0.0485 (7)	0.0384 (6)	0.0303 (6)	-0.0044 (5)	0.0061 (5)	0.0050 (5)
O9	0.0353 (5)	0.0310 (6)	0.0316 (5)	-0.0027 (4)	0.0060 (4)	0.0041 (4)
O10	0.0425 (6)	0.0415 (7)	0.0460 (7)	0.0006 (5)	0.0072 (5)	-0.0023 (6)
O11	0.0536 (7)	0.0299 (6)	0.0456 (7)	-0.0019 (5)	0.0174 (6)	-0.0030 (5)
C1	0.0301 (7)	0.0380 (8)	0.0288 (7)	0.0010 (6)	0.0090 (6)	0.0006 (6)
C2	0.0279 (7)	0.0583 (11)	0.0403 (9)	-0.0027 (7)	0.0040 (6)	0.0043 (8)
C3	0.0341 (8)	0.0551 (11)	0.0430 (9)	-0.0138 (7)	0.0097 (7)	-0.0037 (8)
C4	0.0441 (8)	0.0363 (8)	0.0269 (7)	-0.0080 (6)	0.0128 (6)	-0.0057 (6)
C5	0.0632 (11)	0.0386 (9)	0.0338 (8)	-0.0153 (8)	0.0160 (8)	-0.0045 (7)
C6	0.0842 (14)	0.0346 (9)	0.0402 (9)	-0.0060 (9)	0.0172 (10)	0.0040 (7)
C7	0.0646 (12)	0.0474 (11)	0.0419 (10)	0.0110 (9)	0.0050 (9)	0.0099 (8)
C8	0.0402 (8)	0.0425 (9)	0.0335 (8)	0.0031 (7)	0.0067 (6)	0.0035 (7)
C9	0.0350 (7)	0.0322 (7)	0.0233 (6)	-0.0012 (6)	0.0100 (5)	-0.0029 (5)
C10	0.0280 (6)	0.0321 (7)	0.0233 (6)	-0.0007 (5)	0.0082 (5)	-0.0026 (5)
C11	0.0265 (6)	0.0290 (7)	0.0240 (6)	0.0011 (5)	0.0038 (5)	0.0011 (5)
C12	0.0334 (7)	0.0329 (8)	0.0274 (7)	0.0030 (6)	0.0036 (6)	-0.0035 (6)
C13	0.0491 (10)	0.0577 (11)	0.0386 (9)	0.0016 (8)	0.0127 (8)	-0.0150 (8)
C14	0.0748 (14)	0.0704 (15)	0.0446 (11)	0.0068 (12)	0.0132 (10)	-0.0261 (10)
C15	0.0786 (15)	0.0637 (14)	0.0491 (12)	0.0000 (12)	-0.0043 (11)	-0.0303 (10)
C16	0.0561 (11)	0.0489 (11)	0.0544 (11)	-0.0094 (9)	-0.0077 (9)	-0.0139 (9)
C17	0.0384 (8)	0.0325 (8)	0.0359 (8)	-0.0015 (6)	-0.0022 (6)	-0.0021 (6)
C18	0.0322 (7)	0.0380 (9)	0.0413 (9)	-0.0078 (6)	0.0014 (6)	0.0058 (7)
C19	0.0301 (7)	0.0378 (8)	0.0305 (7)	-0.0010 (6)	0.0087 (6)	0.0065 (6)
C20	0.0278 (6)	0.0251 (6)	0.0226 (6)	0.0022 (5)	0.0027 (5)	0.0022 (5)
C21	0.0634 (12)	0.0490 (11)	0.0548 (11)	0.0026 (9)	0.0304 (10)	-0.0034 (9)
C22	0.0689 (14)	0.0647 (14)	0.0716 (15)	0.0061 (11)	0.0359 (12)	0.0165 (12)

Geometric parameters (\AA , $^\circ$)

P1—O5	1.5112 (11)	C3—H14	0.9300
P1—O3	1.5124 (11)	C4—C5	1.414 (2)
P1—O4	1.5231 (11)	C4—C9	1.426 (2)
P1—O2	1.6330 (11)	C5—C6	1.361 (3)
Na1—O8	2.3256 (14)	C5—H15	0.9300

supplementary materials

Na1—O6 ⁱ	2.3660 (16)	C6—C7	1.405 (3)
Na1—O7	2.4347 (14)	C6—H16	0.9300
Na1—O10	2.4645 (16)	C7—C8	1.371 (3)
Na1—O11	2.4773 (15)	C7—H17	0.9300
Na1—O9	2.6842 (15)	C8—C9	1.418 (2)
Na2—O6	2.3484 (14)	C8—H18	0.9300
Na2—O8	2.4423 (15)	C9—C10	1.430 (2)
Na2—O7	2.4453 (15)	C10—C11	1.494 (2)
Na2—O5	2.4581 (14)	C11—C20	1.374 (2)
Na2—O9	2.4937 (14)	C11—C12	1.427 (2)
Na2—O10 ⁱⁱ	2.5483 (17)	C12—C17	1.426 (2)
O1—C1	1.377 (2)	C12—C13	1.422 (2)
O1—H1	0.847 (17)	C13—C14	1.366 (3)
O2—C20	1.3850 (17)	C13—H19	0.9300
O6—H2	0.852 (16)	C14—C15	1.408 (4)
O6—H3	0.848 (16)	C14—H20	0.9300
O7—H4	0.853 (11)	C15—C16	1.354 (3)
O7—H5	0.851 (14)	C15—H21	0.9300
O8—H6	0.847 (16)	C16—C17	1.423 (2)
O8—H7	0.847 (18)	C16—H22	0.9300
O9—H8	0.852 (15)	C17—C18	1.415 (3)
O9—H9	0.851 (12)	C18—C19	1.363 (2)
O10—H10	0.850 (15)	C18—H23	0.9300
O10—H11	0.849 (12)	C19—C20	1.414 (2)
O11—C21	1.437 (2)	C19—H24	0.9300
O11—H12	0.851 (9)	C21—C22	1.476 (3)
C1—C10	1.374 (2)	C21—H25A	0.9700
C1—C2	1.414 (2)	C21—H25B	0.9700
C2—C3	1.359 (3)	C22—H26A	0.9600
C2—H13	0.9300	C22—H26B	0.9600
C3—C4	1.416 (3)	C22—H26C	0.9600
Na1…Na2	3.2900 (13)	Na2…H3	2.728 (17)
Na1…Na2 ⁱ	3.4236 (12)	Na2…H8	2.612 (18)
O5—P1—O3	114.03 (7)	C10—C1—O1	122.37 (14)
O5—P1—O4	112.55 (6)	C10—C1—C2	121.02 (15)
O3—P1—O4	113.11 (7)	O1—C1—C2	116.57 (14)
O5—P1—O2	108.12 (6)	C3—C2—C1	120.54 (16)
O3—P1—O2	102.51 (6)	C3—C2—H13	119.7
O4—P1—O2	105.51 (6)	C1—C2—H13	119.7
O8—Na1—O6 ⁱ	148.58 (5)	C2—C3—C4	120.93 (15)
O8—Na1—O7	74.98 (5)	C2—C3—H14	119.5
O6 ⁱ —Na1—O7	91.55 (5)	C4—C3—H14	119.5
O8—Na1—O10	120.34 (5)	C3—C4—C5	121.83 (16)
O6 ⁱ —Na1—O10	83.76 (5)	C3—C4—C9	118.65 (15)
O7—Na1—O10	79.89 (5)	C5—C4—C9	119.49 (16)
O8—Na1—O11	97.96 (5)	C6—C5—C4	120.94 (18)
O6 ⁱ —Na1—O11	99.48 (5)	C6—C5—H15	119.5

O7—Na1—O11	167.51 (5)	C4—C5—H15	119.5
O10—Na1—O11	95.38 (5)	C5—C6—C7	119.82 (18)
O8—Na1—O9	70.56 (5)	C5—C6—H16	120.1
O6 ⁱ —Na1—O9	81.61 (4)	C7—C6—H16	120.1
O7—Na1—O9	90.80 (5)	C8—C7—C6	121.10 (18)
O10—Na1—O9	162.42 (5)	C8—C7—H17	119.5
O11—Na1—O9	96.58 (5)	C6—C7—H17	119.5
O6—Na2—O8	122.83 (5)	C7—C8—C9	120.48 (17)
O6—Na2—O7	89.87 (5)	C7—C8—H18	119.8
O8—Na2—O7	72.74 (5)	C9—C8—H18	119.8
O6—Na2—O5	91.94 (5)	C8—C9—C4	118.09 (15)
O8—Na2—O5	91.78 (5)	C8—C9—C10	122.24 (14)
O7—Na2—O5	162.52 (5)	C4—C9—C10	119.66 (14)
O6—Na2—O9	164.98 (5)	C1—C10—C9	119.15 (14)
O8—Na2—O9	72.19 (4)	C1—C10—C11	120.03 (14)
O7—Na2—O9	95.26 (5)	C9—C10—C11	120.82 (13)
O5—Na2—O9	87.40 (4)	C20—C11—C12	118.94 (13)
O6—Na2—O10 ⁱⁱ	82.29 (5)	C20—C11—C10	120.45 (13)
O8—Na2—O10 ⁱⁱ	141.97 (5)	C12—C11—C10	120.60 (13)
O7—Na2—O10 ⁱⁱ	80.01 (4)	C17—C12—C13	118.56 (15)
O5—Na2—O10 ⁱⁱ	117.47 (5)	C17—C12—C11	119.58 (14)
O9—Na2—O10 ⁱⁱ	84.71 (5)	C13—C12—C11	121.87 (15)
C1—O1—H1	113.9 (15)	C14—C13—C12	120.6 (2)
C20—O2—P1	121.22 (9)	C14—C13—H19	119.7
P1—O5—Na2	126.55 (6)	C12—C13—H19	119.7
Na2—O6—Na1 ⁱⁱ	93.14 (5)	C13—C14—C15	120.7 (2)
Na2—O6—H2	123.8 (14)	C13—C14—H20	119.6
Na1 ⁱⁱ —O6—H2	116.8 (15)	C15—C14—H20	119.6
Na2—O6—H3	107.7 (14)	C16—C15—C14	120.21 (18)
Na1 ⁱⁱ —O6—H3	110.7 (14)	C16—C15—H21	119.9
H2—O6—H3	104.2 (19)	C14—C15—H21	119.9
Na1—O7—Na2	84.78 (4)	C15—C16—C17	121.2 (2)
Na1—O7—H4	118.2 (14)	C15—C16—H22	119.4
Na2—O7—H4	112.7 (14)	C17—C16—H22	119.4
Na1—O7—H5	117.6 (14)	C18—C17—C12	118.90 (14)
Na2—O7—H5	119.4 (14)	C18—C17—C16	122.43 (17)
H4—O7—H5	104.1 (19)	C12—C17—C16	118.66 (17)
Na1—O8—Na2	87.23 (5)	C19—C18—C17	121.02 (14)
Na1—O8—H6	110.2 (15)	C19—C18—H23	119.5
Na2—O8—H6	120.8 (15)	C17—C18—H23	119.5
Na1—O8—H7	127.1 (14)	C18—C19—C20	119.86 (14)
Na2—O8—H7	108.8 (15)	C18—C19—H24	120.1
H6—O8—H7	104 (2)	C20—C19—H24	120.1
Na2—O9—Na1	78.81 (4)	C11—C20—O2	118.46 (12)
Na2—O9—H8	88.3 (13)	C11—C20—C19	121.64 (13)
Na1—O9—H8	114.7 (13)	O2—C20—C19	119.82 (13)
Na2—O9—H9	98.4 (13)	O11—C21—C22	111.06 (17)

supplementary materials

Na1—O9—H9	137.1 (14)	O11—C21—H25A	109.4
H8—O9—H9	107.9 (19)	C22—C21—H25A	109.4
Na1—O10—Na2 ⁱ	86.13 (5)	O11—C21—H25B	109.4
Na1—O10—H10	120.7 (15)	C22—C21—H25B	109.4
Na2 ⁱ —O10—H10	113.8 (16)	H25A—C21—H25B	108.0
Na1—O10—H11	118.5 (15)	C21—C22—H26A	109.5
Na2 ⁱ —O10—H11	109.3 (16)	C21—C22—H26B	109.5
H10—O10—H11	107 (2)	H26A—C22—H26B	109.5
C21—O11—Na1	139.11 (12)	C21—C22—H26C	109.5
C21—O11—H12	101.7 (16)	H26A—C22—H26C	109.5
Na1—O11—H12	110.5 (15)	H26B—C22—H26C	109.5
O5—P1—O2—C20	85.95 (11)	O9—Na1—O11—C21	88.50 (18)
O3—P1—O2—C20	-153.29 (10)	C10—C1—C2—C3	-1.8 (3)
O4—P1—O2—C20	-34.69 (12)	O1—C1—C2—C3	-179.66 (16)
O3—P1—O5—Na2	130.38 (8)	C1—C2—C3—C4	-0.2 (3)
O4—P1—O5—Na2	-0.19 (10)	C2—C3—C4—C5	-176.76 (17)
O2—P1—O5—Na2	-116.33 (7)	C2—C3—C4—C9	1.4 (3)
O6—Na2—O5—P1	168.16 (8)	C3—C4—C5—C6	176.72 (17)
O8—Na2—O5—P1	45.22 (8)	C9—C4—C5—C6	-1.5 (3)
O7—Na2—O5—P1	72.42 (18)	C4—C5—C6—C7	3.1 (3)
O9—Na2—O5—P1	-26.85 (8)	C5—C6—C7—C8	-1.8 (3)
O10 ⁱⁱ —Na2—O5—P1	-109.52 (8)	C6—C7—C8—C9	-1.1 (3)
O8—Na2—O6—Na1 ⁱⁱ	-178.06 (5)	C7—C8—C9—C4	2.7 (2)
O7—Na2—O6—Na1 ⁱⁱ	-108.82 (5)	C7—C8—C9—C10	-176.16 (16)
O5—Na2—O6—Na1 ⁱⁱ	88.57 (5)	C3—C4—C9—C8	-179.63 (15)
O9—Na2—O6—Na1 ⁱⁱ	1.4 (2)	C5—C4—C9—C8	-1.4 (2)
O10 ⁱⁱ —Na2—O6—Na1 ⁱⁱ	-28.89 (5)	C3—C4—C9—C10	-0.8 (2)
O8—Na1—O7—Na2	45.99 (4)	C5—C4—C9—C10	177.46 (14)
O6 ⁱ —Na1—O7—Na2	-105.25 (4)	O1—C1—C10—C9	-179.85 (13)
O10—Na1—O7—Na2	171.38 (5)	C2—C1—C10—C9	2.4 (2)
O11—Na1—O7—Na2	102.7 (2)	O1—C1—C10—C11	-0.4 (2)
O9—Na1—O7—Na2	-23.62 (4)	C2—C1—C10—C11	-178.07 (15)
O6—Na2—O7—Na1	-168.48 (5)	C8—C9—C10—C1	177.67 (15)
O8—Na2—O7—Na1	-43.84 (4)	C4—C9—C10—C1	-1.1 (2)
O5—Na2—O7—Na1	-72.42 (15)	C8—C9—C10—C11	-1.8 (2)
O9—Na2—O7—Na1	25.66 (4)	C4—C9—C10—C11	179.38 (13)
O10 ⁱⁱ —Na2—O7—Na1	109.33 (5)	C1—C10—C11—C20	-78.74 (19)
O6 ⁱ —Na1—O8—Na2	21.43 (11)	C9—C10—C11—C20	100.74 (17)
O7—Na1—O8—Na2	-45.89 (4)	C1—C10—C11—C12	100.25 (17)
O10—Na1—O8—Na2	-114.32 (6)	C9—C10—C11—C12	-80.28 (18)
O11—Na1—O8—Na2	144.64 (5)	C20—C11—C12—C17	1.8 (2)
O9—Na1—O8—Na2	50.43 (4)	C10—C11—C12—C17	-177.24 (14)
O6—Na2—O8—Na1	124.57 (5)	C20—C11—C12—C13	-178.02 (15)
O7—Na2—O8—Na1	46.30 (4)	C10—C11—C12—C13	3.0 (2)
O5—Na2—O8—Na1	-141.97 (5)	C17—C12—C13—C14	-0.2 (3)
O9—Na2—O8—Na1	-55.27 (4)	C11—C12—C13—C14	179.57 (19)

O10 ⁱⁱ —Na2—O8—Na1	0.10 (9)	C12—C13—C14—C15	1.4 (4)
O6—Na2—O9—Na1	-133.02 (18)	C13—C14—C15—C16	-1.4 (4)
O8—Na2—O9—Na1	46.47 (4)	C14—C15—C16—C17	0.2 (4)
O7—Na2—O9—Na1	-23.50 (4)	C13—C12—C17—C18	177.98 (16)
O5—Na2—O9—Na1	139.18 (4)	C11—C12—C17—C18	-1.8 (2)
O10 ⁱⁱ —Na2—O9—Na1	-102.92 (4)	C13—C12—C17—C16	-1.0 (2)
Na1 ⁱⁱ —Na2—O9—Na1	-131.92 (3)	C11—C12—C17—C16	179.20 (16)
O8—Na1—O9—Na2	-50.24 (4)	C15—C16—C17—C18	-177.9 (2)
O6 ⁱ —Na1—O9—Na2	114.95 (4)	C15—C16—C17—C12	1.0 (3)
O7—Na1—O9—Na2	23.51 (4)	C12—C17—C18—C19	-0.2 (2)
O10—Na1—O9—Na2	81.01 (16)	C16—C17—C18—C19	178.72 (17)
O11—Na1—O9—Na2	-146.40 (5)	C17—C18—C19—C20	2.3 (2)
O8—Na1—O10—Na2 ⁱ	131.19 (5)	C12—C11—C20—O2	177.15 (12)
O6 ⁱ —Na1—O10—Na2 ⁱ	-27.34 (4)	C10—C11—C20—O2	-3.9 (2)
O7—Na1—O10—Na2 ⁱ	65.36 (4)	C12—C11—C20—C19	0.3 (2)
O11—Na1—O10—Na2 ⁱ	-126.32 (5)	C10—C11—C20—C19	179.30 (13)
O9—Na1—O10—Na2 ⁱ	6.41 (17)	P1—O2—C20—C11	115.23 (13)
O8—Na1—O11—C21	17.30 (18)	P1—O2—C20—C19	-67.86 (16)
O6 ⁱ —Na1—O11—C21	171.05 (18)	C18—C19—C20—C11	-2.3 (2)
O7—Na1—O11—C21	-37.4 (3)	C18—C19—C20—O2	-179.16 (13)
O10—Na1—O11—C21	-104.40 (18)	Na1—O11—C21—C22	-38.2 (3)

Symmetry codes: (i) $-x+3/2, y+1/2, -z$; (ii) $-x+3/2, y-1/2, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O4	0.847 (17)	1.839 (17)	2.6687 (19)	166.3 (18)
O6—H2 \cdots O11 ⁱⁱⁱ	0.852 (16)	2.043 (17)	2.854 (2)	159.0 (16)
O6—H3 \cdots O9 ⁱⁱⁱ	0.848 (16)	1.956 (16)	2.7958 (19)	170.5 (16)
O7—H4 \cdots O5 ⁱ	0.853 (11)	1.873 (10)	2.7136 (18)	168.4 (19)
O7—H5 \cdots O4 ⁱⁱⁱ	0.851 (14)	2.106 (14)	2.9473 (19)	169.9 (18)
O8—H6 \cdots O1 ⁱⁱⁱ	0.847 (16)	1.955 (16)	2.775 (2)	162.7 (16)
O9—H8 \cdots O4	0.852 (15)	1.875 (15)	2.7259 (18)	177.5 (12)
O9—H9 \cdots O3 ^{iv}	0.851 (12)	1.850 (12)	2.6978 (18)	174.1 (16)
O10—H10 \cdots O5 ^v	0.850 (15)	2.197 (15)	3.012 (2)	161 (2)
O10—H11 \cdots O3 ⁱⁱⁱ	0.849 (12)	2.197 (9)	2.984 (2)	154.1 (19)
O11—H12 \cdots O3 ^v	0.851 (9)	1.823 (11)	2.6711 (19)	174 (2)

Symmetry codes: (iii) $x+1/2, -y+1/2, z$; (i) $-x+3/2, y+1/2, -z$; (iv) $-x+1, -y, -z$; (v) $x, y+1, z$.

Fig. 1




Fig. 2




Fig. 3

