

2-Amino-5-nitropyridinium tetraoxido-rhenate(VII) monohydrate

V. H. Rodrigues,^{a*} M. M. R. R. Costa,^a T. Dekola^b and E. de Matos Gomes^b

^aCEMDRX, Department of Physics, University of Coimbra, P-3004-516 Coimbra, Portugal, and ^bDepartamento de Física, Universidade do Minho, P-4710-057 Braga, Portugal

Correspondence e-mail: vhugo@fis.uc.pt


Received 13 July 2009; accepted 23 July 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.015; wR factor = 0.033; data-to-parameter ratio = 15.0.

All the residues of the title compound, $(\text{C}_5\text{H}_6\text{N}_3\text{O}_2)[\text{ReO}_4]\cdot\text{H}_2\text{O}$, are located on general crystallographic positions. The 2-amino-5-nitropyridinium cation has a typical planar conformation with one of the nitro O atoms $-0.058(5)\text{ \AA}$ out of plane; the amine H atoms are also a little out of the main ring plane towards the opposite side of the aforementioned O atom [by 0.02 (4) and 0.04 (4) \AA]. The perrhenate anion is nearly ideally tetrahedral. Three distinct $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds give rise to $C(8)$ zigzag chains running along [100]. $R_4^4(12)$ rings involving the two hydrogen bonds in which the water molecules interact with the perrhenate anions are also present.

Related literature

For the structural analyses of related 2-amino-5-nitropyridinium salts and their potential application as non-linear optical materials, see: Masse & Zyss (1991); Puig-Molina *et al.* (1998); Aakeröy *et al.* (1998); Pecaut *et al.* (1993).



Experimental

Crystal data

$(\text{C}_5\text{H}_6\text{N}_3\text{O}_2)[\text{ReO}_4]\cdot\text{H}_2\text{O}$	$V = 1031.08(6)\text{ \AA}^3$
$M_r = 408.34$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.6914(3)\text{ \AA}$	$\mu = 11.81\text{ mm}^{-1}$
$b = 9.1357(3)\text{ \AA}$	$T = 295\text{ K}$
$c = 13.4636(4)\text{ \AA}$	$0.10 \times 0.06 \times 0.05\text{ mm}$
$\beta = 120.120(2)^{\circ}$	

Data collection

Bruker APEXII diffractometer	30757 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	2374 independent reflections
$T_{\min} = 0.33$, $T_{\max} = 0.55$	2175 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.033$	$\Delta\rho_{\text{max}} = 0.88\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\text{min}} = -0.72\text{ e \AA}^{-3}$
2374 reflections	
158 parameters	
5 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^{\circ}$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O3	0.86	1.93	2.738 (3)	157
N1—H1 \cdots O6 ⁱ	0.86	2.63	3.049 (4)	112
N2—H2A \cdots O3	0.89 (3)	2.33 (3)	3.030 (4)	135 (3)
N2—H2A \cdots O4 ⁱⁱ	0.89 (3)	2.65 (3)	3.259 (4)	127 (3)
N2—H2B \cdots O5	0.89 (3)	1.98 (3)	2.846 (3)	164 (3)
O3—H3A \cdots O7 ⁱⁱⁱ	0.89 (3)	1.93 (3)	2.802 (4)	167 (3)
O3—H3B \cdots O4 ⁱⁱ	0.89 (3)	2.13 (3)	2.862 (4)	139 (3)

Symmetry codes: (i) $x - 1, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 1, -y, -z$; (iii) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

This work was supported by Fundação para a Ciência e a Tecnologia (FCT).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2177).

References

- Aakeröy, C. B., Beatty, A. M., Nieuwenhuyzen, M. & Zou, M. (1998). *J. Mater. Chem.* **8**, 1385–1389.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Masse, R. & Zyss, J. (1991). *Mol. Eng.* **1**, 141–152.
- Pecaut, J., Le Fur, Y. & Masse, R. (1993). *Acta Cryst. B* **49**, 535–541.
- Puig-Molina, A., Alvarez-Larena, A., Piniella, J. F., Howard, S. T. & Baert, F. (1998). *Struct. Chem.* **9**, 395–402.
- Sheldrick, G. M. (1996). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2009). E65, m1000 [doi:10.1107/S1600536809029365]

2-Amino-5-nitropyridinium tetraoxidorhenate(VII) monohydrate

V. H. Rodrigues, M. M. R. R. Costa, T. Dekola and E. de Matos Gomes

S1. Experimental

Crystals of the title compound were obtained by slow evaporation from a water solution of analytical grade reagents on a 1:1 molar ratio. The reagents used were 2-amino-5-nitropyridine (99.5%) and a perrhenic acid solution (65–75% water 99.5%), both purchased from Aldrich.

S2. Refinement

The structure was solved by direct methods using *SHELXS97*. All H atoms were first located on a difference Fourier map as a check for data quality; those bonded to C and N aromatic atoms were then placed at idealized positions and refined as riding [$C-H=0.93\text{ \AA}$, $N-H=0.86\text{ \AA}$, $U_{iso}(H)=1.2U_{eq}(C)$ and $U_{iso}(H)=1.2U_{eq}(N)$]. The coordinates of the amine H atoms were then refined only with the restraint that the $N-H$ distance would be 0.89 \AA within 0.2 \AA . The positions of the H atoms belonging to the water molecule were also restrained so that the intramolecular O—H distance would be 0.89 \AA and the H—H distance 1.40 \AA , both within 0.02 \AA , thus enforcing an acceptable geometry of the water molecule [$U_{iso}(H)=1.5U_{eq}(O)$].

Examination of the crystal structure with *PLATON* (Spek, 2009) showed that there are no solvent-accessible voids in the crystal lattice.





Figure 1


ORTEPII (Johnson, 1976) plot of the title compound. Displacement ellipsoids are drawn at the 50% level.

**Figure 2**

Zigzag chain running along the a axis. Small ring motifs made of a perrhenate anion and two water molecules are included in the chain. Hydrogen bonds are indicated with dashed lines.

2-Amino-5-nitropyridinium tetraoxidorhenate(VII) monohydrate

Crystal data



$M_r = 408.34$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.6914(3)$ Å

$b = 9.1357(3)$ Å

$c = 13.4636(4)$ Å

$\beta = 120.120(2)^\circ$

$V = 1031.08(6)$ Å³

$Z = 4$

$F(000) = 760$

$D_x = 2.631$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9920 reflections

$\theta = 2.4\text{--}32.3^\circ$

$\mu = 11.81$ mm⁻¹

$T = 295$ K

Prism, translucent colourless

0.10 × 0.06 × 0.05 mm

Data collection

Bruker APEXII
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.33$, $T_{\max} = 0.55$

30757 measured reflections

2374 independent reflections

2175 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.4^\circ$

$h = -12 \rightarrow 12$

$k = -11 \rightarrow 11$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.015$

$wR(F^2) = 0.033$

$S = 1.05$

2374 reflections

158 parameters

5 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0122P)^2 + 1.4451P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.88 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.72 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.00774 (16)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.2958 (3)	0.4684 (3)	0.0719 (2)	0.0338 (5)
H1	0.2145	0.4119	0.0493	0.041*
C2	0.4404 (3)	0.4063 (3)	0.1067 (2)	0.0315 (6)
N2	0.4512 (3)	0.2632 (3)	0.1047 (3)	0.0434 (6)
H2A	0.367 (3)	0.204 (3)	0.080 (3)	0.052*
H2B	0.543 (2)	0.219 (4)	0.124 (3)	0.052*
C3	0.5736 (3)	0.4998 (3)	0.1440 (3)	0.0352 (6)
H3	0.6741	0.4605	0.1684	0.042*
C4	0.5549 (3)	0.6463 (3)	0.1442 (2)	0.0362 (6)
H4	0.6420	0.7086	0.1688	0.043*
C5	0.4020 (3)	0.7029 (3)	0.1067 (2)	0.0332 (6)
N3	0.3785 (4)	0.8592 (3)	0.1068 (2)	0.0437 (6)
O1	0.4946 (4)	0.9377 (3)	0.1387 (3)	0.0717 (9)
O2	0.2459 (3)	0.9055 (3)	0.0781 (3)	0.0632 (7)
C6	0.2737 (3)	0.6133 (3)	0.0712 (2)	0.0336 (6)
H6	0.1728	0.6517	0.0469	0.040*
O3	0.1012 (3)	0.2314 (2)	0.0301 (2)	0.0538 (6)
H3A	0.083 (5)	0.174 (3)	0.076 (2)	0.081*
H3B	0.079 (5)	0.174 (3)	-0.030 (2)	0.081*
Re	0.905465 (12)	0.037384 (13)	0.215124 (10)	0.03195 (6)
O4	0.8222 (3)	-0.1153 (3)	0.1338 (2)	0.0680 (7)
O5	0.7677 (3)	0.1758 (3)	0.1707 (3)	0.0632 (7)
O6	0.9713 (4)	-0.0024 (3)	0.3551 (2)	0.0674 (8)
O7	1.0618 (3)	0.0893 (3)	0.1987 (3)	0.0664 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0264 (11)	0.0290 (12)	0.0452 (15)	-0.0007 (9)	0.0174 (11)	0.0031 (11)
C2	0.0289 (12)	0.0319 (14)	0.0329 (15)	0.0032 (11)	0.0150 (11)	0.0054 (12)
N2	0.0375 (13)	0.0281 (13)	0.0627 (18)	0.0052 (11)	0.0237 (14)	0.0057 (13)

C3	0.0262 (13)	0.0400 (16)	0.0369 (16)	0.0004 (11)	0.0142 (12)	0.0010 (13)
C4	0.0312 (13)	0.0411 (17)	0.0341 (16)	-0.0090 (12)	0.0147 (12)	-0.0035 (13)
C5	0.0417 (15)	0.0277 (14)	0.0296 (15)	0.0020 (12)	0.0175 (12)	0.0023 (12)
N3	0.0597 (17)	0.0299 (13)	0.0415 (15)	0.0009 (12)	0.0254 (13)	0.0003 (11)
O1	0.080 (2)	0.0345 (13)	0.108 (3)	-0.0181 (13)	0.0530 (19)	-0.0083 (14)
O2	0.0674 (17)	0.0398 (14)	0.0761 (19)	0.0152 (13)	0.0313 (15)	0.0003 (13)
C6	0.0320 (13)	0.0329 (15)	0.0351 (15)	0.0054 (12)	0.0161 (12)	0.0046 (13)
O3	0.0643 (16)	0.0382 (13)	0.0750 (19)	-0.0090 (12)	0.0471 (15)	-0.0043 (13)
Re	0.02646 (7)	0.03566 (8)	0.03418 (8)	0.00296 (4)	0.01554 (5)	0.00215 (5)
O4	0.0773 (19)	0.0537 (16)	0.0667 (18)	-0.0151 (14)	0.0315 (15)	-0.0156 (14)
O5	0.0447 (13)	0.0583 (16)	0.0843 (19)	0.0226 (12)	0.0307 (13)	0.0131 (15)
O6	0.085 (2)	0.0780 (19)	0.0394 (15)	0.0150 (16)	0.0313 (14)	0.0090 (13)
O7	0.0398 (13)	0.083 (2)	0.086 (2)	0.0057 (13)	0.0389 (14)	0.0225 (17)

Geometric parameters (\AA , ^\circ)

N1—C6	1.341 (4)	C5—C6	1.360 (4)
N1—C2	1.358 (3)	C5—N3	1.446 (4)
N1—H1	0.8600	N3—O1	1.216 (4)
C2—N2	1.313 (4)	N3—O2	1.217 (4)
C2—C3	1.413 (4)	C6—H6	0.9300
N2—H2A	0.89 (3)	O3—H3A	0.89 (3)
N2—H2B	0.89 (3)	O3—H3B	0.89 (3)
C3—C4	1.351 (4)	Re—O6	1.698 (3)
C3—H3	0.9300	Re—O7	1.704 (2)
C4—C5	1.402 (4)	Re—O4	1.705 (3)
C4—H4	0.9300	Re—O5	1.714 (2)
C6—N1—C2	123.2 (2)	C6—C5—N3	118.6 (3)
C6—N1—H1	118.4	C4—C5—N3	120.1 (3)
C2—N1—H1	118.4	O1—N3—O2	123.3 (3)
N2—C2—N1	119.2 (3)	O1—N3—C5	117.7 (3)
N2—C2—C3	122.8 (3)	O2—N3—C5	119.0 (3)
N1—C2—C3	118.0 (3)	N1—C6—C5	118.5 (2)
C2—N2—H2A	123 (2)	N1—C6—H6	120.7
C2—N2—H2B	122 (2)	C5—C6—H6	120.7
H2A—N2—H2B	115 (3)	H3A—O3—H3B	104 (3)
C4—C3—C2	120.0 (3)	O6—Re—O7	109.89 (15)
C4—C3—H3	120.0	O6—Re—O4	109.31 (15)
C2—C3—H3	120.0	O7—Re—O4	108.01 (15)
C3—C4—C5	118.9 (3)	O6—Re—O5	109.77 (14)
C3—C4—H4	120.6	O7—Re—O5	109.69 (13)
C5—C4—H4	120.6	O4—Re—O5	110.15 (14)
C6—C5—C4	121.3 (3)	 	
C6—N1—C2—N2	179.9 (3)	C4—C5—N3—O1	1.0 (4)
C6—N1—C2—C3	0.1 (4)	C6—C5—N3—O2	1.9 (4)
N2—C2—C3—C4	-179.9 (3)	C4—C5—N3—O2	-177.2 (3)

N1—C2—C3—C4	−0.1 (4)	C2—N1—C6—C5	0.1 (4)
C2—C3—C4—C5	−0.2 (4)	C4—C5—C6—N1	−0.4 (4)
C3—C4—C5—C6	0.4 (4)	N3—C5—C6—N1	−179.5 (3)
C3—C4—C5—N3	179.5 (3)	N1—C2—N2—H2A	1 (3)
C6—C5—N3—O1	−179.9 (3)	C3—C2—N2—H2B	−3 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O3	0.86	1.93	2.738 (3)	157
N1—H1···O6 ⁱ	0.86	2.63	3.049 (4)	112
N2—H2A···O3	0.89 (3)	2.33 (3)	3.030 (4)	135 (3)
N2—H2A···O4 ⁱⁱ	0.89 (3)	2.65 (3)	3.259 (4)	127 (3)
N2—H2B···O5	0.89 (3)	1.98 (3)	2.846 (3)	164 (3)
O3—H3A···O7 ⁱⁱⁱ	0.89 (3)	1.93 (3)	2.802 (4)	167 (3)
O3—H3B···O4 ⁱⁱ	0.89 (3)	2.13 (3)	2.862 (4)	139 (3)

Symmetry codes: (i) $x-1, -y+1/2, z-1/2$; (ii) $-x+1, -y, -z$; (iii) $x-1, y, z$.