

(E)-2-(2-Nitroprop-1-enyl)furan

Pedro Valerga,^{a*} M. Carmen Puerta,^a Zenaida Rodríguez Negrín,^b Nilo Castañedo Cancio^b and Miguel Palma Lovillo^c

^aDepartamento de Ciencia de los Materiales e Ingeniería Metalúrgica, Facultad de Ciencias, Campus Universitario del Río San Pedro, Puerto Real 11510, Spain,
^bCentro de Bioactivos Químicos, Universidad Central Marta Abreu de Las Villas, Cuba, and ^cDepartamento de Química Analítica, Facultad de Ciencias, Campus Universitario del Río San Pedro, Puerto Real 11510, Spain
Correspondence e-mail: pedro.valerga@uca.es


Received 25 July 2009; accepted 29 July 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.049; wR factor = 0.130; data-to-parameter ratio = 15.9.

Crystals of the title compound, $\text{C}_7\text{H}_7\text{NO}_3$, under Mo $K\alpha$ radiation sublime in less than 1 h at room temperature. However, it was possible to collect data at 100 K. It crystallized as the *E* isomer only. A double-bond conjugation in the furan ring is extended to the nitroalkenyl group. Molecular associations were realized in the crystal through $\text{N}\cdots\pi$ [3.545 (2) Å] interactions involving the furan ring and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For general background to (nitro-alkenyl)-furan compounds, see: Yan *et al.* (2008); Ono N. (2006); Vallejos *et al.* (2005); Negrín *et al.* (2003); Negrín *et al.* (2002), Estrada *et al.* (1999); Agafonov *et al.* (1991); Gruntfest *et al.* (1972). For related structures, see: Valerga *et al.* (2009); Martínez-Bescos *et al.* (2008); Novoa-de-Armas *et al.* (1997); Pomés *et al.* (1995).



Experimental

Crystal data

$\text{C}_7\text{H}_7\text{NO}_3$
 $M_r = 153.14$
Monoclinic, $P2_1/n$
 $a = 7.1061 (14)\text{ \AA}$
 $b = 9.4394 (19)\text{ \AA}$
 $c = 10.743 (2)\text{ \AA}$
 $\beta = 101.86 (3)^\circ$

$V = 705.2 (3)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.12\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.45 \times 0.30 \times 0.18\text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2004)
 $T_{\min} = 0.898$, $T_{\max} = 1.000$
(expected range = 0.880–0.980)

5648 measured reflections
1620 independent reflections
1497 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.130$
 $S = 1.07$
1620 reflections

102 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.34\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2···O3 ⁱ	0.95	2.47	3.3037 (19)	147
C5—H5···O3 ⁱ	0.95	3.03	3.770 (2)	136
C4—H4···O2 ⁱⁱ	0.95	2.65	3.2980 (19)	126
C4—H4···O3 ⁱⁱ	0.95	2.58	3.516 (2)	170
C7—H7C···O2 ⁱⁱⁱ	0.98	2.70	3.310 (2)	121

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x, y + 1, z$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL*.

We thank the SCYCT (Universidad de Cádiz) for the X-ray data collection and the Consejería de Innovación, Ciencia y Empresa de la Junta de Andalucía, for financial support. ZRN thanks the AUIP and Aula Iberoamericana for the stay at UCA.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2230).

References

- Agafonov, N. E., Sredishev, I. P., Dudin, A. V., Kutin, A. A., Stashina, G. A. & Zhulin, V. M. (1991). *Izv. Akad. Nauk SSSR Ser. Khim.* **2**, 426–433.
- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Estrada, E., Gomez, M., Castañedo, N. & Perez, C. (1999). *J. Mol. Struct. THEOCHEM*, **468**, 193–200. [Amended journal title OK?].
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gruntfest, M. G., Potemkin, G. F., Kolodyazhnyi, Y. V., Zverev, V. V., Nazarova, Z. N. & Osipov, O. A. (1972). *Zh. Org. Khim.* **8**, 404–411.
- Martínez-Bescos, P., Cagide-Fagin, F., Roa, L. F., Ortiz-Lara, J. C., Kierus, K., Ozores-Viturro, L., Fernández-González, M. & Alonso, R. (2008). *J. Org. Chem.* **73**, 3745–3753.
- Negrín, Z. R., Martínez, B. N. H., Meseguer, G. P., Placeres, E. G. & Molina, M. I. D. (2003). *Cent. Azúcar*, **30**, 30–34.
- Negrín, Z. R., Placeres, E. G., Martínez, B. N. H., Meseguer, G. P. & Montenegro, O. N. (2002). *Cent. Azúcar*, **29**, 79–86.
- Novoa-de-Armas, H., Pomes-Hernández, R., Duque-Rodríguez, J. & Toscano, R. A. (1997). *Z. Kristallogr.* **212**, 63–63.
- Ono, N. (2006). *Sci. Synth.* **33**, 337–370.
- Pomés, R., Duque, J., Novoa, H., Castañedo, N. & Toscano, A. (1995). *Acta Cryst. C51*, 1368–1369.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.

- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Valerga, P., Puerta, M. C., Rodríguez Negrín, Z., Castañedo Cancio, N. & Palma Lovillo, M. (2009). *Acta Cryst. E* **65**, o1979.
- Vallejos, G., Fierro, A., Rezende, M. C., Sepúlveda-Bozab, S. & Reyes-Parada, M. (2005). *Bioorg. Med. Chem.* **14**, 4450–4457.
- Yan, S., Gao, Y., Xing, R., Shen, Y., Liu, Y., Wu, P. & Wu, H. (2008). *Tetrahedron Lett.* **64**, 6294–6299.

supporting information

Acta Cryst. (2009). E65, o2060–o2061 [doi:10.1107/S1600536809030141]

(*E*)-2-(2-Nitroprop-1-enyl)furan

Pedro Valerga, M. Carmen Puerta, Zenaida Rodríguez Negrín, Nilo Castañedo Cancio and Miguel Palma Lovillo

S1. Comment

Some (nitro-alkenyl)-furan compounds show antibacterial and antifungal activities and were described and patented as drugs ingredients for veterinarian and agricultural purposes. We recently started the structural study of this group of furyl-nitroolefins. The title compound crystallized exclusively in its *E* configuration (Fig. 1). The conjugated system of double bonds in furan ring is extended to the alkenyl group being C2—C3 and C1—C5 bond lengths 1.422 (2) and 1.431 (2) Å, significatively shorter than single C—C bonds. Distances N1—O2 = 1.232 (2) and N1—O3= 1.231 (2) indicate conjugation with a delocalized double bond. Alkenyl C5 and C6 sp^2 carbons are coplanar with the furan ring as shown by an angle of 3.9 (1) $^\circ$ between ring plane and C5 C6 C7 N1 plane. Crystal packing shows N··· π interactions involving the furan ring: N···Cg (1/2-x, -1/2+y, 1/2-z) distance is 3.545 (2) Å (Fig. 2) and CH···O hydrogen bonds (Table 1).

S2. Experimental

2-(2-Nitro-propen-1-yl)-furan, also called UC-244, was obtained using the Knoevenagel's condensation method by reaction of furfural, an aromatic compound from acid hydrolysis of sugar cane residuals (straw, sawdust, etc.) and nitroethane in the presence of isobutylamine as a catalyst. To obtain a product with purity higher than 99% the method was optimized studying temperature, contact and reaction times as variables. The purification was achieved using activated coal and ethanol. The yellow crystals should be protected from the light and heating. ^1H NMR (CDCl_3) δ (ppm): 2.511 (3H, s, $-\text{CH}_3$), 6.533 (1H, dd, 2J = 3.6 Hz and 2J = 1.6 Hz, $-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 6.781 (1H, d, 2J = 3.6Hz, $-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 7.599 (1H, d, 2J = 1.6 Hz, $-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 7.775 (1H, s, $\text{HC}=\text{CMe}$) $^{13}\text{C}\{\text{H}\}$ NMR (CDCl_3) δ (ppm): 13.656 ($-\text{C}\text{H}_3$), 112.688 ($-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 119.100 ($-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 120.346 ($-\text{C}=\text{C}(\text{Me})\text{NO}_2$), 144.097 ($-\text{C}=\text{C}(\text{Me})\text{NO}_2$), 146.104 ($-\text{O}-\text{CH}=\text{CH}-\text{CH}=$), 147.679 ($\text{C}_{\text{ring}}-\text{CH}-\text{C}=\text{C}(\text{Me})\text{NO}_2$).

S3. Refinement

All H atoms were positioned geometrically and treated as riding ($\text{C}-\text{H}$ = 0.99 Å for methylene and $\text{C}-\text{H}$ = 0.93 Å otherwise). $U_{\text{iso}}(\text{H})$ = 1.2 $U_{\text{eq}}(\text{C})$ of the carrier atom. In the absence of any significant anomalous scatters, the Friedel pairs were merged before final refinements.





Figure 1

ORTEP representation of I with the atom labelling scheme and displacement ellipsoids at the 50% probability level.

**Figure 2**

Packing diagram of *E*-2-(2-nitro-propen-1-yl)-furan.

(*E*)-2-(2-Nitroprop-1-enyl)furan

Crystal data

$C_7H_7NO_3$

$M_r = 153.14$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 7.1061 (14)$ Å

$b = 9.4394 (19)$ Å

$c = 10.743 (2)$ Å

$\beta = 101.86 (3)^\circ$

$V = 705.2 (3)$ Å³

$Z = 4$

$F(000) = 320$

$D_x = 1.442 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2619 reflections

$\theta = 2.9\text{--}27.6^\circ$

$\mu = 0.12 \text{ mm}^{-1}$

$T = 100$ K

Irregular, yellow

$0.45 \times 0.30 \times 0.18$ mm

Data collection

Bruker SMART APEX
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
1700 ω scan frames, 0.3°, 10s
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2004)
 $T_{\min} = 0.898$, $T_{\max} = 1.000$

5648 measured reflections
1620 independent reflections
1497 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.9^\circ$
 $h = -9 \rightarrow 8$
 $k = -12 \rightarrow 12$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.130$
 $S = 1.07$
1620 reflections
102 parameters
0 restraints
0 constraints

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.069P)^2 + 0.2975P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$
Extinction correction: *SHELXL*,
 $F_c^* = kF_c[1 + 0.001xF_c^2l^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.012 (3)

Special details

Experimental. Refinement of F^2 against unique set of reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against unique set of reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.08533 (15)	0.64923 (11)	0.35499 (10)	0.0257 (3)
O2	0.22698 (17)	0.12565 (12)	0.41245 (11)	0.0328 (3)
O3	-0.00888 (16)	0.10699 (11)	0.25033 (11)	0.0311 (3)
N1	0.09873 (17)	0.17838 (13)	0.33088 (11)	0.0227 (3)
C1	0.19682 (19)	0.55649 (15)	0.43728 (13)	0.0200 (3)
C2	0.31539 (19)	0.62992 (15)	0.53073 (13)	0.0211 (3)
H2	0.4068	0.5911	0.5993	0.025*
C3	0.2760 (2)	0.77609 (16)	0.50590 (15)	0.0272 (4)
H3	0.3360	0.8539	0.5545	0.033*
C4	0.1374 (2)	0.78256 (15)	0.40026 (16)	0.0278 (4)
H4	0.0828	0.8679	0.3620	0.033*
C5	0.18569 (19)	0.40627 (15)	0.41909 (12)	0.0194 (3)

H5	0.2709	0.3514	0.4802	0.023*
C6	0.07004 (19)	0.33308 (14)	0.32658 (12)	0.0197 (3)
C7	-0.0833 (2)	0.38425 (16)	0.22071 (14)	0.0266 (3)
H7A	-0.1245	0.4791	0.2409	0.040*
H7B	-0.1932	0.3193	0.2092	0.040*
H7C	-0.0336	0.3881	0.1422	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0287 (6)	0.0186 (5)	0.0277 (5)	0.0010 (4)	0.0008 (4)	0.0023 (4)
O2	0.0399 (7)	0.0183 (5)	0.0339 (6)	0.0058 (5)	-0.0073 (5)	0.0003 (4)
O3	0.0372 (7)	0.0217 (6)	0.0304 (6)	-0.0050 (4)	-0.0023 (5)	-0.0073 (4)
N1	0.0263 (6)	0.0186 (6)	0.0223 (6)	-0.0007 (5)	0.0028 (5)	-0.0024 (5)
C1	0.0210 (7)	0.0172 (7)	0.0222 (7)	0.0015 (5)	0.0051 (5)	0.0019 (5)
C2	0.0197 (7)	0.0216 (7)	0.0217 (7)	-0.0002 (5)	0.0038 (5)	-0.0009 (5)
C3	0.0296 (8)	0.0210 (7)	0.0324 (8)	-0.0057 (6)	0.0100 (6)	-0.0069 (6)
C4	0.0321 (8)	0.0145 (7)	0.0383 (8)	0.0012 (6)	0.0105 (6)	0.0031 (6)
C5	0.0199 (7)	0.0173 (7)	0.0206 (6)	0.0013 (5)	0.0034 (5)	0.0010 (5)
C6	0.0217 (7)	0.0164 (6)	0.0211 (7)	0.0014 (5)	0.0047 (5)	0.0005 (5)
C7	0.0292 (8)	0.0254 (7)	0.0224 (7)	0.0037 (6)	-0.0014 (6)	-0.0020 (6)

Geometric parameters (\AA , ^\circ)

O1—C4	1.3724 (18)	C3—C4	1.342 (2)
O1—C1	1.3739 (17)	C3—H3	0.9500
O2—N1	1.2320 (16)	C4—H4	0.9500
O3—N1	1.2311 (16)	C5—C6	1.3434 (19)
N1—C6	1.4739 (18)	C5—H5	0.9500
C1—C2	1.3602 (19)	C6—C7	1.4850 (19)
C1—C5	1.431 (2)	C7—H7A	0.9800
C2—C3	1.422 (2)	C7—H7B	0.9800
C2—H2	0.9500	C7—H7C	0.9800
C4—O1—C1	106.17 (12)	C3—C4—H4	124.6
O3—N1—O2	122.73 (12)	O1—C4—H4	124.6
O3—N1—C6	117.28 (11)	C6—C5—C1	128.22 (13)
O2—N1—C6	119.99 (11)	C6—C5—H5	115.9
C2—C1—O1	109.76 (13)	C1—C5—H5	115.9
C2—C1—C5	127.88 (13)	C5—C6—N1	115.26 (12)
O1—C1—C5	122.34 (12)	C5—C6—C7	129.82 (13)
C1—C2—C3	106.73 (13)	N1—C6—C7	114.92 (12)
C1—C2—H2	126.6	C6—C7—H7A	109.5
C3—C2—H2	126.6	C6—C7—H7B	109.5
C4—C3—C2	106.53 (13)	H7A—C7—H7B	109.5
C4—C3—H3	126.7	C6—C7—H7C	109.5
C2—C3—H3	126.7	H7A—C7—H7C	109.5
C3—C4—O1	110.81 (13)	H7B—C7—H7C	109.5

C4—O1—C1—C2	−0.25 (15)	O1—C1—C5—C6	−1.4 (2)
C4—O1—C1—C5	−178.62 (12)	C1—C5—C6—N1	177.53 (12)
O1—C1—C2—C3	0.10 (15)	C1—C5—C6—C7	−2.8 (2)
C5—C1—C2—C3	178.36 (13)	O3—N1—C6—C5	177.57 (12)
C1—C2—C3—C4	0.08 (16)	O2—N1—C6—C5	−2.79 (19)
C2—C3—C4—O1	−0.24 (17)	O3—N1—C6—C7	−2.15 (18)
C1—O1—C4—C3	0.31 (16)	O2—N1—C6—C7	177.49 (12)
C2—C1—C5—C6	−179.46 (13)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C2—H2···O3 ⁱ	0.95	2.47	3.3037 (19)	147
C5—H5···O3 ⁱ	0.95	3.03	3.770 (2)	136
C4—H4···O2 ⁱⁱ	0.95	2.65	3.2980 (19)	126
C4—H4···O3 ⁱⁱ	0.95	2.58	3.516 (2)	170
C7—H7C···O2 ⁱⁱⁱ	0.98	2.70	3.310 (2)	121

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x, y+1, z$; (iii) $x-1/2, -y+1/2, z-1/2$.