

## 2-(2H-Benzotriazol-2-yl)-4-methylphenyl diphenylphosphinate

Yi-Chang Liu, Chia-Her Lin and Bao-Tsan Ko\*

Department of Chemistry, Chung Yuan Christian University, Chung-Li 320, Taiwan  
Correspondence e-mail: btko@cycu.edu.tw


Received 21 July 2009; accepted 28 July 2009

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  
 $R$  factor = 0.032;  $wR$  factor = 0.086; data-to-parameter ratio = 13.2.

In the title molecule,  $\text{C}_{25}\text{H}_{20}\text{N}_3\text{O}_2\text{P}$ , the dihedral angle between the mean planes of the benzotriazol ring system and the  $N$ -bonded benzene ring is  $45.8(2)^\circ$ . All but one of the angles at the P atom show slight distortions from an ideal tetrahedral geometry.

### Related literature

For background to the use of 2-(2H-benzotriazol-2H-yl)-phenol (BTP-H) derivatives, see: Li *et al.* (2009); Tsai *et al.* (2009). For related structures, see: Al-Farhan (1992); Cheng *et al.* (2007).



### Experimental

#### Crystal data

$\text{C}_{25}\text{H}_{20}\text{N}_3\text{O}_2\text{P}$

$M_r = 425.41$

Orthorhombic,  $Pca2_1$   
 $a = 12.7691(7)\text{ \AA}$   
 $b = 9.4064(5)\text{ \AA}$   
 $c = 18.6362(10)\text{ \AA}$   
 $V = 2238.4(2)\text{ \AA}^3$

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.15\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.30 \times 0.20 \times 0.15\text{ mm}$

#### Data collection

Bruker SMART-1000 CCD  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $(S) = 0.963$ ,  $T_{\max} = 0.974$

11920 measured reflections  
3684 independent reflections  
3343 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.086$   
 $S = 1.05$   
3684 reflections  
280 parameters  
1 restraint

H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.14\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
1412 Friedel pairs  
Flack parameter: 0.01 (7)

**Table 1**  
Selected bond angles ( $^\circ$ ).

O2—P—O1	115.53 (10)	O2—P—C20	112.37 (9)
O2—P—C14	113.48 (8)	O1—P—C20	105.04 (8)
O1—P—C14	100.15 (8)	C14—P—C20	109.30 (8)

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We gratefully acknowledge the financial support in part from the National Science Council, Taiwan (NSC97-2113-M-033-005-MY2) and in part from the project of the specific research fields in the Chung Yuan Christian University, Taiwan (No. CYCU-97-CR-CH).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2871).

### References

- Al-Farhan, K. A. (1992). *J. Crystallogr. Spectrosc. Res.* **22**, 687–689.
- Bruker (1999). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheng, Y.-H., Weng, C.-M. & Hong, F.-E. (2007). *Tetrahedron*, **63**, 12277–12285.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Li, C.-Y., Lin, C.-H. & Ko, B.-T. (2009). *Acta Cryst. E* **65**, m670.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tsai, C.-Y., Lin, C.-H. & Ko, B.-T. (2009). *Acta Cryst. E* **65**, m619.

# supporting information

*Acta Cryst.* (2009). E65, o2058 [doi:10.1107/S1600536809029870]

## 2-(2H-Benzotriazol-2-yl)-4-methylphenyl diphenylphosphinate

**Yi-Chang Liu, Chia-Her Lin and Bao-Tsan Ko**

### S1. Comment

2-(2H-benzotriazol-2H-yl)phenol (BTP-H) derivatives are widely used as ultraviolet (UV) absorbers for the protection of commercially important synthetic plastics and fibers against the UV light in industry. In terms of coordination chemistry, the benzotriazol-phenolate group can provide N, O-bidentate chelation to stabilize the transition metal or main group metal complexes. For instance, our group has successfully synthesized and structural characterized the Pd complex (II) with 4-methyl-2-(2H-benzotriazol-2-yl)-phenolate ligands (Tsai *et al.*, 2009). Recently, we also reported the synthesis and crystal structure of an Al(III) complex with the 4-methyl-2-(2H-benzotriazol-2-yl)-phenolate ligand (Li *et al.*, 2009). Most recently, Cheng *et al.* (2007) reported some palladium complexes of monodentate phosphinite ligands and these complexes in the presence of  $\text{Pd}(\text{OAc})_2$  have been demonstrated effectively to catalyze Suzuki–Miyaura cross-coupling reactions. Therefore, our group is interested in the synthesis and preparation of the phosphinite functionalized benzotriazol-phenolate ligands derived from BTP-H. Here, we report the synthesis and crystal structure of the title compound, (**I**), a potential ligand for the preparation of metal complexes.


The molecule of (**I**) is composed of a benzotriazol-phenolate moiety and a diphenylphosphine oxide functionalized group (Fig. 1). The dihedral angle between the planes of the benzotriazole moiety and the benzene ring of the phenoxy group is  $45.8(2)^\circ$ . The P atom is bonded one O atom of the phosphine oxide, one phenoxy O atom and two C atoms from two phenyl groups, forming a slightly distorted tetrahedral environment. The P–O and P=O bond distances bond distances are similar to those found in the crystal structure of triphenylphosphine oxide (Al-Farhan, 1992).

### S2. Experimental


The title compound **I** was synthesized by the following procedure (Fig. 2): To a rapidly stirred solution of 4-methyl-2-(2H-benzotriazol-2-yl)phenol (2.48 g, 10.0 mmol) in toluene (20 ml),  $\text{Ph}_2\text{PCl}$  (1.8 ml, 10.0 mmol) and  $\text{NEt}_3$  (20.0 mmol, 2.02 g) was slowly added. The mixed solution was stirred at 363 K for 18 h. Subsequently, the  $\text{HNEt}_3\text{Cl}$  salt was filtered and the resulting solution was dried under reduced pressure. The resulting oily product was re-dissolved in MeOH (20 ml) and  $\text{H}_2\text{O}_2$  (1 ml) was added. The final solution was stirred at room temperature for another 1 h and the volatile components were removed in *vacuo*. The residue was extracted with ethyl acetate (50 ml) and the extract was dried under vacuum to give oily, white solids. Colorless crystals were obtained on cooling the saturated  $\text{Et}_2\text{O}$  solution at 253 K overnight.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , p.p.m.):  $\delta$  7.10–7.97 (17H, m, ArH), 2.35 (3H, s,  $\text{CH}_3$ ).

### S3. Refinement

The H atoms were placed in idealized positions and constrained to ride on their parent atoms, with C—H = 0.93 and 0.96 Å with  $U_{\text{iso}}(\text{H}) = 1.2$  and  $1.5U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of **I** with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

The synthetic procedure of the title compound.

**2-(2*H*-Benzotriazol-2-yl)-4-methylphenyl diphenylphosphinate***Crystal data*

$C_{25}H_{20}N_3O_2P$   
 $M_r = 425.41$   
Orthorhombic,  $Pca2_1$   
Hall symbol: P 2c -2ac  
 $a = 12.7691 (7) \text{ \AA}$   
 $b = 9.4064 (5) \text{ \AA}$   
 $c = 18.6362 (10) \text{ \AA}$   
 $V = 2238.4 (2) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 888$   
 $D_x = 1.262 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 6495 reflections  
 $\theta = 2.7\text{--}26.0^\circ$   
 $\mu = 0.15 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Block, colourless  
 $0.30 \times 0.20 \times 0.15 \text{ mm}$

*Data collection*

Bruker SMART-1000 CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.963$ ,  $T_{\max} = 0.974$

11920 measured reflections  
3684 independent reflections  
3343 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$   
 $\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.2^\circ$   
 $h = -15 \rightarrow 14$   
 $k = -11 \rightarrow 11$   
 $l = -19 \rightarrow 22$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.086$   
 $S = 1.05$   
3684 reflections  
280 parameters  
1 restraint  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.058P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.14 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$   
Absolute structure: Flack (1983), 1412 Friedel  
pairs  
Absolute structure parameter: 0.01 (7)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
P	-0.78562 (3)	-0.27388 (4)	-0.41236 (3)	0.04851 (13)
O1	-0.74629 (11)	-0.25123 (13)	-0.49386 (7)	0.0523 (3)
O2	-0.89556 (10)	-0.23744 (14)	-0.39862 (10)	0.0724 (4)

N1	-0.54417 (13)	-0.30745 (18)	-0.54995 (9)	0.0573 (4)
N2	-0.58778 (12)	-0.21611 (16)	-0.59589 (8)	0.0510 (3)
N3	-0.56809 (15)	-0.23649 (17)	-0.66558 (9)	0.0617 (4)
C1	-0.73324 (14)	-0.11822 (19)	-0.52534 (11)	0.0498 (4)
C2	-0.65289 (15)	-0.10047 (19)	-0.57421 (10)	0.0536 (4)
C3	-0.63716 (18)	0.0309 (2)	-0.60682 (12)	0.0641 (5)
H3B	-0.5823	0.0422	-0.6391	0.077*
C4	-0.7020 (2)	0.1451 (2)	-0.59187 (13)	0.0675 (6)
C5	-0.78319 (19)	0.1244 (2)	-0.54483 (14)	0.0710 (6)
H5A	-0.8283	0.1996	-0.5351	0.085*
C6	-0.80000 (17)	-0.0052 (2)	-0.51143 (12)	0.0652 (5)
H6A	-0.8558	-0.0163	-0.4798	0.078*
C7	-0.49173 (14)	-0.3965 (2)	-0.59453 (10)	0.0543 (4)
C8	-0.4308 (2)	-0.5185 (3)	-0.57796 (13)	0.0768 (6)
H8A	-0.4209	-0.5492	-0.5310	0.092*
C9	-0.3876 (2)	-0.5879 (3)	-0.63507 (16)	0.0861 (7)
H9A	-0.3471	-0.6683	-0.6265	0.103*
C10	-0.4020 (2)	-0.5427 (3)	-0.70636 (15)	0.0862 (7)
H10A	-0.3709	-0.5943	-0.7432	0.103*
C11	-0.45980 (19)	-0.4263 (3)	-0.72317 (13)	0.0762 (6)
H11A	-0.4683	-0.3963	-0.7704	0.091*
C12	-0.50631 (15)	-0.3531 (2)	-0.66534 (11)	0.0561 (4)
C13	-0.6860 (3)	0.2874 (3)	-0.6283 (2)	0.0984 (9)
H13A	-0.7377	0.3535	-0.6114	0.148*
H13B	-0.6929	0.2760	-0.6793	0.148*
H13C	-0.6173	0.3227	-0.6173	0.148*
C14	-0.75764 (14)	-0.45832 (18)	-0.40115 (10)	0.0511 (4)
C15	-0.66460 (14)	-0.5202 (2)	-0.42229 (12)	0.0612 (5)
H15A	-0.6144	-0.4658	-0.4459	0.073*
C16	-0.64533 (18)	-0.6626 (2)	-0.40869 (16)	0.0746 (6)
H16A	-0.5822	-0.7036	-0.4226	0.089*
C17	-0.7204 (3)	-0.7429 (2)	-0.37445 (17)	0.0840 (8)
H17A	-0.7086	-0.8391	-0.3663	0.101*
C18	-0.8118 (3)	-0.6822 (3)	-0.35251 (19)	0.0982 (9)
H18A	-0.8609	-0.7367	-0.3281	0.118*
C19	-0.83227 (19)	-0.5408 (3)	-0.36614 (16)	0.0811 (7)
H19A	-0.8956	-0.5008	-0.3520	0.097*
C20	-0.69692 (13)	-0.17140 (17)	-0.35845 (9)	0.0460 (4)
C21	-0.73139 (19)	-0.1249 (2)	-0.29154 (12)	0.0636 (6)
H21A	-0.7985	-0.1470	-0.2757	0.076*
C22	-0.6643 (2)	-0.0451 (3)	-0.24858 (12)	0.0791 (7)
H22A	-0.6865	-0.0153	-0.2035	0.095*
C23	-0.56668 (2)	-0.0102 (2)	-0.27184 (15)	0.0770 (6)
H23A	-0.5234	0.0449	-0.2430	0.092*
C24	-0.53189 (18)	-0.0559 (2)	-0.33777 (13)	0.0677 (5)
H24A	-0.4653	-0.0310	-0.3535	0.081*
C25	-0.59558 (14)	-0.1389 (2)	-0.38062 (11)	0.0562 (4)
H25A	-0.5708	-0.1731	-0.4243	0.067*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
P	0.0399 (2)	0.0488 (2)	0.0569 (3)	-0.00062 (15)	0.0029 (2)	-0.0003 (2)
O1	0.0566 (7)	0.0499 (6)	0.0504 (7)	0.0003 (5)	-0.0023 (6)	0.0000 (5)
O2	0.0446 (7)	0.0726 (9)	0.0998 (14)	0.0035 (6)	0.0074 (7)	0.0013 (8)
N1	0.0617 (9)	0.0667 (10)	0.0436 (8)	0.0101 (8)	-0.0017 (7)	0.0031 (7)
N2	0.0576 (9)	0.0532 (8)	0.0424 (8)	0.0008 (6)	-0.0003 (6)	0.0025 (6)
N3	0.0772 (11)	0.0662 (10)	0.0416 (9)	0.0011 (8)	0.0019 (8)	0.0047 (7)
C1	0.0529 (10)	0.0484 (10)	0.0481 (10)	0.0013 (7)	-0.0086 (8)	-0.0012 (8)
C2	0.0621 (10)	0.0524 (10)	0.0462 (10)	0.0023 (8)	-0.0080 (8)	0.0007 (8)
C3	0.0749 (13)	0.0586 (12)	0.0588 (12)	-0.0026 (9)	-0.0006 (9)	0.0065 (9)
C4	0.0864 (14)	0.0510 (11)	0.0650 (14)	0.0023 (9)	-0.0146 (11)	0.0061 (10)
C5	0.0842 (14)	0.0547 (11)	0.0742 (15)	0.0183 (10)	-0.0125 (12)	-0.0007 (10)
C6	0.0687 (12)	0.0643 (12)	0.0625 (13)	0.0107 (9)	-0.0018 (10)	0.0021 (10)
C7	0.0533 (10)	0.0600 (10)	0.0498 (10)	0.0030 (8)	-0.0007 (8)	-0.0018 (8)
C8	0.0810 (13)	0.0845 (15)	0.0650 (13)	0.0261 (12)	0.0004 (12)	0.0052 (12)
C9	0.0862 (16)	0.0871 (16)	0.0849 (18)	0.0310 (13)	0.0016 (14)	-0.0089 (14)
C10	0.0905 (17)	0.0927 (17)	0.0752 (17)	0.0148 (14)	0.0108 (13)	-0.0277 (14)
C11	0.0877 (16)	0.0905 (17)	0.0504 (12)	0.0048 (13)	0.0065 (11)	-0.0092 (11)
C12	0.0598 (10)	0.0610 (11)	0.0476 (10)	-0.0026 (8)	0.0029 (8)	-0.0012 (8)
C13	0.124 (2)	0.0571 (13)	0.114 (2)	0.0010 (14)	0.0021 (19)	0.0205 (14)
C14	0.0516 (8)	0.0477 (8)	0.0540 (11)	-0.0075 (7)	-0.0029 (8)	-0.0012 (8)
C15	0.0549 (10)	0.0556 (10)	0.0731 (14)	0.0019 (7)	0.0007 (9)	0.0058 (10)
C16	0.0793 (13)	0.0602 (11)	0.0842 (15)	0.0131 (10)	-0.0135 (13)	-0.0024 (12)
C17	0.110 (2)	0.0502 (12)	0.0916 (19)	-0.0074 (12)	-0.0226 (15)	0.0083 (12)
C18	0.112 (2)	0.0664 (14)	0.116 (3)	-0.0274 (15)	0.0132 (18)	0.0180 (15)
C19	0.0705 (13)	0.0671 (14)	0.106 (2)	-0.0147 (10)	0.0195 (13)	0.0097 (13)
C20	0.0529 (9)	0.0382 (8)	0.0467 (9)	0.0036 (7)	0.0034 (7)	-0.0001 (7)
C21	0.0804 (16)	0.0569 (12)	0.0534 (13)	0.0099 (10)	0.0133 (10)	0.0009 (9)
C22	0.122 (2)	0.0656 (14)	0.0500 (12)	0.0125 (14)	-0.0011 (13)	-0.0161 (11)
C23	0.0982 (17)	0.0576 (12)	0.0752 (15)	-0.0024 (11)	-0.0260 (14)	-0.0107 (11)
C24	0.0624 (12)	0.0676 (13)	0.0733 (14)	-0.0109 (9)	-0.0138 (10)	-0.0027 (11)
C25	0.0527 (9)	0.0622 (11)	0.0536 (10)	-0.0032 (8)	0.0018 (8)	-0.0064 (8)

Geometric parameters ( $\text{\AA}$ ,  $\text{^\circ}$ )

P—O2	1.4675 (14)	C11—C12	1.410 (3)
P—O1	1.6138 (15)	C11—H11A	0.9300
P—C14	1.7835 (18)	C13—H13A	0.9600
P—C20	1.7948 (18)	C13—H13B	0.9600
O1—C1	1.392 (2)	C13—H13C	0.9600
N1—N2	1.335 (2)	C14—C15	1.380 (3)
N1—C7	1.357 (2)	C14—C19	1.391 (3)
N2—N3	1.337 (2)	C15—C16	1.385 (3)
N2—C2	1.427 (2)	C15—H15A	0.9300
N3—C12	1.351 (3)	C16—C17	1.377 (4)
C1—C2	1.382 (3)	C16—H16A	0.9300

C1—C6	1.387 (3)	C17—C18	1.362 (4)
C2—C3	1.392 (3)	C17—H17A	0.9300
C3—C4	1.385 (3)	C18—C19	1.379 (4)
C3—H3B	0.9300	C18—H18A	0.9300
C4—C5	1.371 (4)	C19—H19A	0.9300
C4—C13	1.515 (3)	C20—C25	1.392 (3)
C5—C6	1.385 (3)	C20—C21	1.393 (3)
C5—H5A	0.9300	C21—C22	1.392 (4)
C6—H6A	0.9300	C21—H21A	0.9300
C7—C12	1.394 (3)	C22—C23	1.359 (4)
C7—C8	1.420 (3)	C22—H22A	0.9300
C8—C9	1.365 (4)	C23—C24	1.376 (4)
C8—H8A	0.9300	C23—H23A	0.9300
C9—C10	1.407 (4)	C24—C25	1.381 (3)
C9—H9A	0.9300	C24—H24A	0.9300
C10—C11	1.357 (4)	C25—H25A	0.9300
C10—H10A	0.9300		
O2—P—O1	115.53 (10)	N3—C12—C7	108.62 (16)
O2—P—C14	113.48 (8)	N3—C12—C11	129.8 (2)
O1—P—C14	100.15 (8)	C7—C12—C11	121.60 (19)
O2—P—C20	112.37 (9)	C4—C13—H13A	109.5
O1—P—C20	105.04 (8)	C4—C13—H13B	109.5
C14—P—C20	109.30 (8)	H13A—C13—H13B	109.5
C1—O1—P	123.55 (12)	C4—C13—H13C	109.5
N2—N1—C7	102.17 (15)	H13A—C13—H13C	109.5
N1—N2—N3	116.91 (16)	H13B—C13—H13C	109.5
N1—N2—C2	123.51 (16)	C15—C14—C19	119.22 (18)
N3—N2—C2	119.59 (15)	C15—C14—P	123.30 (13)
N2—N3—C12	102.88 (15)	C19—C14—P	117.40 (16)
C2—C1—C6	119.13 (19)	C14—C15—C16	120.55 (19)
C2—C1—O1	118.37 (16)	C14—C15—H15A	119.7
C6—C1—O1	122.45 (18)	C16—C15—H15A	119.7
C1—C2—C3	120.14 (18)	C17—C16—C15	119.5 (2)
C1—C2—N2	121.79 (16)	C17—C16—H16A	120.3
C3—C2—N2	117.97 (18)	C15—C16—H16A	120.3
C4—C3—C2	121.0 (2)	C18—C17—C16	120.3 (2)
C4—C3—H3B	119.5	C18—C17—H17A	119.8
C2—C3—H3B	119.5	C16—C17—H17A	119.8
C5—C4—C3	118.1 (2)	C17—C18—C19	120.8 (2)
C5—C4—C13	120.9 (2)	C17—C18—H18A	119.6
C3—C4—C13	121.0 (3)	C19—C18—H18A	119.6
C4—C5—C6	122.0 (2)	C18—C19—C14	119.6 (2)
C4—C5—H5A	119.0	C18—C19—H19A	120.2
C6—C5—H5A	119.0	C14—C19—H19A	120.2
C5—C6—C1	119.7 (2)	C25—C20—C21	119.38 (18)
C5—C6—H6A	120.2	C25—C20—P	122.57 (13)
C1—C6—H6A	120.2	C21—C20—P	118.05 (15)

N1—C7—C12	109.42 (16)	C22—C21—C20	119.3 (2)
N1—C7—C8	129.50 (19)	C22—C21—H21A	120.3
C12—C7—C8	121.07 (18)	C20—C21—H21A	120.3
C9—C8—C7	116.0 (2)	C23—C22—C21	120.7 (2)
C9—C8—H8A	122.0	C23—C22—H22A	119.7
C7—C8—H8A	122.0	C21—C22—H22A	119.7
C8—C9—C10	122.6 (2)	C22—C23—C24	120.4 (2)
C8—C9—H9A	118.7	C22—C23—H23A	119.8
C10—C9—H9A	118.7	C24—C23—H23A	119.8
C11—C10—C9	122.2 (2)	C23—C24—C25	120.1 (2)
C11—C10—H10A	118.9	C23—C24—H24A	119.9
C9—C10—H10A	118.9	C25—C24—H24A	119.9
C10—C11—C12	116.5 (2)	C24—C25—C20	119.98 (19)
C10—C11—H11A	121.7	C24—C25—H25A	120.0
C12—C11—H11A	121.7	C20—C25—H25A	120.0
O2—P—O1—C1	69.76 (16)	N1—C7—C12—N3	0.0 (2)
C14—P—O1—C1	-167.96 (14)	C8—C7—C12—N3	179.2 (2)
C20—P—O1—C1	-54.64 (15)	N1—C7—C12—C11	179.55 (19)
C7—N1—N2—N3	-0.5 (2)	C8—C7—C12—C11	-1.2 (3)
C7—N1—N2—C2	179.47 (16)	C10—C11—C12—N3	-179.2 (2)
N1—N2—N3—C12	0.5 (2)	C10—C11—C12—C7	1.3 (3)
C2—N2—N3—C12	-179.50 (16)	O2—P—C14—C15	169.14 (18)
P—O1—C1—C2	145.53 (15)	O1—P—C14—C15	45.41 (19)
P—O1—C1—C6	-36.9 (2)	C20—P—C14—C15	-64.59 (19)
C6—C1—C2—C3	2.4 (3)	O2—P—C14—C19	-14.0 (2)
O1—C1—C2—C3	-179.93 (17)	O1—P—C14—C19	-137.76 (18)
C6—C1—C2—N2	-174.01 (18)	C20—P—C14—C19	112.24 (19)
O1—C1—C2—N2	3.7 (3)	C19—C14—C15—C16	-0.2 (3)
N1—N2—C2—C1	-47.2 (3)	P—C14—C15—C16	176.57 (17)
N3—N2—C2—C1	132.8 (2)	C14—C15—C16—C17	0.7 (4)
N1—N2—C2—C3	136.33 (19)	C15—C16—C17—C18	-1.6 (4)
N3—N2—C2—C3	-43.7 (2)	C16—C17—C18—C19	2.1 (5)
C1—C2—C3—C4	-0.9 (3)	C17—C18—C19—C14	-1.6 (5)
N2—C2—C3—C4	175.64 (18)	C15—C14—C19—C18	0.6 (4)
C2—C3—C4—C5	-1.0 (3)	P—C14—C19—C18	-176.3 (2)
C2—C3—C4—C13	-179.0 (2)	O2—P—C20—C25	-151.46 (15)
C3—C4—C5—C6	1.4 (4)	O1—P—C20—C25	-25.08 (17)
C13—C4—C5—C6	179.3 (3)	C14—P—C20—C25	81.63 (17)
C4—C5—C6—C1	0.1 (4)	O2—P—C20—C21	29.44 (18)
C2—C1—C6—C5	-2.0 (3)	O1—P—C20—C21	155.82 (14)
O1—C1—C6—C5	-179.60 (18)	C14—P—C20—C21	-97.47 (16)
N2—N1—C7—C12	0.33 (19)	C25—C20—C21—C22	1.0 (3)
N2—N1—C7—C8	-178.8 (2)	P—C20—C21—C22	-179.91 (17)
N1—C7—C8—C9	179.6 (2)	C20—C21—C22—C23	1.2 (3)
C12—C7—C8—C9	0.6 (3)	C21—C22—C23—C24	-1.4 (4)
C7—C8—C9—C10	0.0 (4)	C22—C23—C24—C25	-0.6 (4)
C8—C9—C10—C11	0.2 (5)	C23—C24—C25—C20	2.7 (3)

---

C9—C10—C11—C12	−0.8 (4)	C21—C20—C25—C24	−2.9 (3)
N2—N3—C12—C7	−0.3 (2)	P—C20—C25—C24	178.04 (16)
N2—N3—C12—C11	−179.8 (2)		

---