

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(1Z)-1-(2,4-Dichlorophenyl)ethan-1-one semicarbazone

Hoong-Kun Fun,^{a,*} Kasthuri Balasubramani,^a A. M. Vijesh,^b Shridhar Malladi^c and Arun M. Isloor^c^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 Universiti Sains Malaysia, Penang, Malaysia, ^bSeQuent Scientific Limited, No. 120 A&B, Industrial Area, Baikampady, New Bangalore, Karnataka 575 011, India, and^cDepartment of Chemistry, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India

Correspondence e-mail: hkfun@usm.my


Received 7 July 2009; accepted 28 July 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.072; wR factor = 0.192; data-to-parameter ratio = 28.2.

In the title compound, $\text{C}_9\text{H}_9\text{Cl}_2\text{N}_3\text{O}$, the semicarbazone group is approximately planar, with an r.m.s deviation from the mean plane of 0.011 (2) Å. The dihedral angle between the least-squares planes through the semicarbazone group and the benzene ring is 38.76 (9)°. The crystal structure is further stabilized by $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonding.

Related literature

For applications of semicarbazone derivatives, see: Warren *et al.* (1977); Chandra & Gupta (2005); Jain *et al.* (2002); Pilgram (1978); Yogeewari *et al.* (2004); For semicarbazide preparations, see: Furniss *et al.* (1978). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_9\text{H}_9\text{Cl}_2\text{N}_3\text{O}$ $M_r = 246.09$ Monoclinic, $C2/c$ $a = 37.8079$ (17) Å $b = 3.8097$ (2) Å $c = 14.4920$ (7) Å

* Thomson Reuters ResearcherID: A-3561-2009.

[§] Department of Chemistry, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India.

$\beta = 98.852$ (2)°
 $V = 2062.52$ (17) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 0.60$ mm⁻¹
 $T = 100$ K
 $0.42 \times 0.14 \times 0.04$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.707$, $T_{\max} = 0.974$

32124 measured reflections
 4202 independent reflections
 3654 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.072$ $wR(F^2) = 0.192$ $S = 1.11$

4202 reflections

149 parameters

H atoms treated by a mixture of
 independent and constrained
 refinement

 $\Delta\rho_{\text{max}} = 3.37$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.82$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H1N2}\cdots\text{O1}^{\text{i}}$	0.85 (3)	2.07 (3)	2.907 (2)	168 (3)
$\text{N3}-\text{H2N3}\cdots\text{O1}^{\text{ii}}$	0.81 (4)	2.13 (4)	2.924 (2)	164 (4)
$\text{C9}-\text{H9A}\cdots\text{O1}^{\text{iii}}$	0.96	2.59	3.465 (2)	152

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x, y, -z + \frac{1}{2}$; (iii) $-x, -y + 2, -z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and KBS thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. KBS thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HKF also thanks Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012. AMI is grateful to the Head of the Department of Chemistry and the Director, NITK, Surathkal, India, for providing research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2153).

References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chandra, S. & Gupta, L. K. (2005). *Spectrochim. Acta Part A*, **62**, 1089–1094.
- Furniss, B. S., Hannaford, A. J., Rogers, V., Smith, P. W. G. & Tatchell, A. R. (1978). *Vogel's Textbook of Practical Organic Chemistry*, 4th ed., p. 1112. London: ELBS.
- Jain, V. K., Handa, A., Pandya, R., Shrivastav, P. & Agrawal, Y. K. (2002). *React. Funct. Polym.* **51**, 101–110.
- Pilgram, K. H. G. (1978). US Patent No. 4 108 399.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Warren, J. D., Woodward, D. L. & Hargreaves, R. T. (1977). *J. Med. Chem.* **20**, 1520–1521.
- Yogeewari, P., Sriram, D., Pandeya, S. N. & Stables, J. P. (2004). *Farmaco*, **59**, 609–613.

supplementary materials

Acta Cryst. (2009). E65, o2072 [doi:10.1107/S1600536809029900]

(1Z)-1-(2,4-Dichlorophenyl)ethan-1-one semicarbazone

H.-K. Fun, K. Balasubramani, A. M. Vijesh, S. Malladi and A. M. Isloor

Comment

In organic chemistry, a semicarbazone is a derivative of an aldehyde or ketone formed by a condensation between a ketone or aldehyde and semicarbazide. They find immense applications in the field of synthetic chemistry, such as medicinal chemistry (Warren *et al.*, 1977), organometallics (Chandra & Gupta, 2005), polymers (Jain *et al.*, 2002) and herbicides (Pilgram, 1978). 4-Sulphamoylphenyl semicarbazones were synthesized and were found to possess anticonvulsant activity (Yogeeswari *et al.*, 2004). Keeping in view of their biological importance, we hereby reporting crystal structure of the semicarbazone of commercial importance.

The semicarbazone group (Fig. 1) (C9/C6/C7/N1/N2/C8/O1/N3) is approximately planar, with an r.m.s deviation of 0.011 (2) Å for atom N1, while the dihedral angle between the least-squares plane through the semicarbazone group and the benzene ring is 38.76 (9)°. The molecules are linked *via* N—H···O hydrogen bonds to generate $R_2^2(8)$ ring motifs (Bernstein *et al.* 1995). These motifs are further connected through C—H···O hydrogen bonds to form a one-dimensional chain along the [0 1 0] direction (Fig. 2).

Experimental

3.16 g (28.3 mmol) of semicarbazide hydrochloride and 2.83 g (34.5 mmol) of crystallized sodium acetate was dissolved in 25 ml of water (Furniss *et al.*, 1978). The reaction mixture was stirred at room temperature for 10 minutes. (5.0 g, 26.5 mmol) of 2,4-dichloroacetophenone in 25 ml of ethanol was then added and the mixture stirred well for 6 h. The separated semicarbazone was filtered, washed with chilled water and recrystallized from an ethanol-DMF mixture. Yield was found to be 5.23 g, 86.02%. *M.p.* 501–503 K.

Refinement

H atoms were positioned geometrically (C—H = 0.93–0.96 Å) and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{methyl C})$. A rotating-group model was used for the methyl groups. The nitrogen H atoms were located from the difference Fourier map [$\text{N—H} = 0.85(3)–0.91(3)$ Å] and allowed to refine freely.

Figures




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom numbering scheme.




Fig. 2. An one-dimensional chain of (I) with $R_2^2(8)$ ring motifs along the [010] direction. Dashed lines indicate the hydrogen bonding.

(1Z)-1-(2,4-Dichlorophenyl)ethan-1-one semicarbazone

Crystal data

$C_9H_9Cl_2N_3O$

$M_r = 246.09$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 37.8079$ (17) Å

$b = 3.8097$ (2) Å

$c = 14.4920$ (7) Å

$\beta = 98.852$ (2)°

$V = 2062.52$ (17) Å³

$Z = 8$

$F_{000} = 1008$

$D_x = 1.585$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9961 reflections

$\theta = 2.9\text{--}34.0^\circ$

$\mu = 0.60$ mm⁻¹

$T = 100$ K

Plate, colorless

$0.42 \times 0.14 \times 0.04$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100$ K

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.707$, $T_{\max} = 0.974$

32124 measured reflections

4202 independent reflections

3654 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 34.1^\circ$

$\theta_{\min} = 1.1^\circ$

$h = -59 \rightarrow 58$

$k = -5 \rightarrow 5$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.072$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.192$	$w = 1/[\sigma^2(F_o^2) + (0.1176P)^2 + 5.115P]$
$S = 1.11$	where $P = (F_o^2 + 2F_c^2)/3$
4202 reflections	$(\Delta/\sigma)_{\max} = 0.001$
149 parameters	$\Delta\rho_{\max} = 3.37 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.82 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.254197 (12)	1.14825 (12)	0.14159 (3)	0.01486 (13)
C12	0.149699 (12)	0.64254 (12)	-0.11406 (3)	0.01413 (13)
O1	-0.00572 (4)	0.4635 (5)	0.11828 (10)	0.0162 (3)
N1	0.08094 (4)	0.7191 (5)	0.08948 (11)	0.0127 (3)
N2	0.04458 (4)	0.6666 (5)	0.06771 (12)	0.0141 (3)
N3	0.04484 (5)	0.5433 (7)	0.22395 (13)	0.0222 (4)
C1	0.14889 (5)	1.0482 (6)	0.14140 (13)	0.0132 (3)
H1A	0.1327	1.0954	0.1819	0.016*
C2	0.18500 (5)	1.1187 (5)	0.17051 (14)	0.0132 (3)
H2A	0.1928	1.2131	0.2293	0.016*
C3	0.20913 (5)	1.0451 (5)	0.10974 (13)	0.0120 (3)
C4	0.19805 (5)	0.9013 (5)	0.02261 (13)	0.0124 (3)
H4A	0.2145	0.8508	-0.0172	0.015*
C5	0.16165 (5)	0.8336 (5)	-0.00436 (13)	0.0112 (3)
C6	0.13617 (5)	0.9086 (5)	0.05322 (13)	0.0112 (3)
C7	0.09697 (5)	0.8539 (5)	0.02541 (13)	0.0116 (3)
C8	0.02647 (5)	0.5559 (6)	0.13727 (14)	0.0152 (3)
C9	0.07833 (5)	0.9765 (5)	-0.06811 (13)	0.0123 (3)
H9A	0.0571	1.1034	-0.0601	0.018*
H9B	0.0720	0.7772	-0.1078	0.018*
H9C	0.0940	1.1272	-0.0962	0.018*
H1N2	0.0320 (9)	0.660 (8)	0.014 (2)	0.016 (7)*
H1N3	0.0663 (10)	0.636 (10)	0.231 (3)	0.028 (9)*

supplementary materials

H2N3 0.0324 (10) 0.484 (10) 0.262 (3) 0.028 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0092 (2)	0.0180 (2)	0.0168 (2)	-0.00196 (13)	0.00036 (15)	-0.00011 (14)
C12	0.0141 (2)	0.0179 (2)	0.0105 (2)	-0.00169 (14)	0.00232 (15)	-0.00251 (14)
O1	0.0090 (6)	0.0283 (8)	0.0111 (6)	-0.0027 (5)	0.0010 (5)	0.0002 (5)
N1	0.0076 (6)	0.0192 (7)	0.0111 (7)	-0.0008 (5)	0.0009 (5)	0.0005 (5)
N2	0.0084 (6)	0.0238 (8)	0.0097 (7)	-0.0026 (5)	0.0005 (5)	0.0012 (5)
N3	0.0114 (7)	0.0461 (12)	0.0088 (7)	-0.0060 (8)	0.0002 (6)	0.0043 (7)
C1	0.0100 (7)	0.0191 (8)	0.0104 (7)	-0.0019 (6)	0.0016 (6)	-0.0020 (6)
C2	0.0114 (7)	0.0163 (8)	0.0114 (7)	-0.0007 (6)	0.0005 (6)	-0.0018 (6)
C3	0.0082 (7)	0.0140 (8)	0.0135 (8)	-0.0007 (6)	0.0006 (6)	0.0011 (6)
C4	0.0119 (7)	0.0126 (7)	0.0130 (8)	-0.0005 (6)	0.0027 (6)	-0.0004 (6)
C5	0.0113 (7)	0.0127 (7)	0.0097 (7)	0.0001 (5)	0.0021 (6)	-0.0005 (5)
C6	0.0104 (7)	0.0130 (7)	0.0101 (7)	-0.0007 (6)	0.0015 (5)	0.0013 (6)
C7	0.0099 (7)	0.0141 (8)	0.0105 (7)	-0.0007 (5)	0.0009 (6)	-0.0003 (5)
C8	0.0109 (7)	0.0234 (9)	0.0116 (8)	-0.0011 (7)	0.0025 (6)	0.0017 (7)
C9	0.0107 (7)	0.0140 (8)	0.0114 (7)	-0.0011 (6)	-0.0010 (6)	0.0017 (6)

Geometric parameters (\AA , $^\circ$)

C11—C3	1.7403 (19)	C1—H1A	0.9300
C12—C5	1.7436 (19)	C2—C3	1.391 (3)
O1—C8	1.256 (2)	C2—H2A	0.9300
N1—C7	1.291 (2)	C3—C4	1.381 (3)
N1—N2	1.377 (2)	C4—C5	1.395 (3)
N2—C8	1.369 (2)	C4—H4A	0.9300
N2—H1N2	0.85 (3)	C5—C6	1.398 (3)
N3—C8	1.339 (3)	C6—C7	1.489 (3)
N3—H1N3	0.88 (4)	C7—C9	1.503 (3)
N3—H2N3	0.81 (4)	C9—H9A	0.9600
C1—C2	1.392 (3)	C9—H9B	0.9600
C1—C6	1.399 (3)	C9—H9C	0.9600
C7—N1—N2	117.12 (16)	C4—C5—C6	122.38 (17)
C8—N2—N1	118.08 (16)	C4—C5—C12	116.02 (14)
C8—N2—H1N2	113 (2)	C6—C5—C12	121.59 (15)
N1—N2—H1N2	128 (2)	C5—C6—C1	116.83 (17)
C8—N3—H1N3	115 (2)	C5—C6—C7	123.95 (17)
C8—N3—H2N3	112 (3)	C1—C6—C7	119.21 (16)
H1N3—N3—H2N3	131 (4)	N1—C7—C6	114.76 (16)
C2—C1—C6	122.24 (17)	N1—C7—C9	124.43 (17)
C2—C1—H1A	118.9	C6—C7—C9	120.65 (16)
C6—C1—H1A	118.9	O1—C8—N3	122.71 (18)
C3—C2—C1	118.53 (17)	O1—C8—N2	120.16 (18)
C3—C2—H2A	120.7	N3—C8—N2	117.11 (18)
C1—C2—H2A	120.7	C7—C9—H9A	109.5

C4—C3—C2	121.52 (17)	C7—C9—H9B	109.5
C4—C3—C11	118.62 (14)	H9A—C9—H9B	109.5
C2—C3—C11	119.83 (15)	C7—C9—H9C	109.5
C3—C4—C5	118.48 (17)	H9A—C9—H9C	109.5
C3—C4—H4A	120.8	H9B—C9—H9C	109.5
C5—C4—H4A	120.8		
C7—N1—N2—C8	-173.82 (19)	C12—C5—C6—C7	-3.6 (3)
C6—C1—C2—C3	-0.4 (3)	C2—C1—C6—C5	1.6 (3)
C1—C2—C3—C4	-0.7 (3)	C2—C1—C6—C7	-177.46 (18)
C1—C2—C3—C11	177.08 (15)	N2—N1—C7—C6	179.31 (17)
C2—C3—C4—C5	0.7 (3)	N2—N1—C7—C9	4.0 (3)
C11—C3—C4—C5	-177.15 (14)	C5—C6—C7—N1	137.72 (19)
C3—C4—C5—C6	0.5 (3)	C1—C6—C7—N1	-43.3 (3)
C3—C4—C5—C12	-178.54 (15)	C5—C6—C7—C9	-46.8 (3)
C4—C5—C6—C1	-1.6 (3)	C1—C6—C7—C9	132.2 (2)
C12—C5—C6—C1	177.39 (15)	N1—N2—C8—O1	-171.2 (2)
C4—C5—C6—C7	177.37 (18)	N1—N2—C8—N3	7.1 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H1N2 \cdots O1 ⁱ	0.85 (3)	2.07 (3)	2.907 (2)	168 (3)
N3—H2N3 \cdots O1 ⁱⁱ	0.81 (4)	2.13 (4)	2.924 (2)	164 (4)
C9—H9A \cdots O1 ⁱⁱⁱ	0.96	2.59	3.465 (2)	152

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, y, -z+1/2$; (iii) $-x, -y+2, -z$.

Fig. 1




Fig. 2

