

4-[4-(Dimethylamino)benzylidene]-2,6-dimethylcyclohexa-2,5-dienone

Nathalie Hampel, Dorothea Richter, Armin R. Ofial,
Herbert Mayr and Peter Mayer*

Ludwig-Maximilians-Universität, Department, Butenandtstrasse 5–13, 81377 München, Germany

Correspondence e-mail: pemay@cup.uni-muenchen.de


Received 2 July 2009; accepted 3 August 2009

Key indicators: single-crystal X-ray study; $T = 200\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.049; wR factor = 0.131; data-to-parameter ratio = 16.3.

The title compound, $C_{17}H_{19}\text{NO}$, crystallized with two molecules per asymmetric unit. $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds lead to infinite chains along [100]. According to graph-set theory, the descriptor $C_1^1(13)C_1^1(13)$ can be assigned.

Related literature

For a related structure, see: Kawai *et al.* (2004). For background to graph set analysis, see: Bernstein *et al.* (1995); Etter *et al.* (1990). For the preparation, see: Richter *et al.* (2009).



Experimental

Crystal data

$C_{17}H_{19}\text{NO}$	$V = 2818.74 (11)\text{ \AA}^3$
$M_r = 253.34$	$Z = 8$
Monoclinic, $P2_1/c$	$\text{Mo } K\alpha$ radiation
$a = 14.5357 (3)\text{ \AA}$	$\mu = 0.07\text{ mm}^{-1}$
$b = 7.2759 (2)\text{ \AA}$	$T = 200\text{ K}$
$c = 27.5473 (5)\text{ \AA}$	$0.24 \times 0.20 \times 0.19\text{ mm}$
$\beta = 104.6463 (14)^\circ$	

Data collection

Nonius KappaCCD diffractometer
Absorption correction: none
18635 measured reflections

5725 independent reflections
3685 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.131$
 $S = 1.02$
5725 reflections

351 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C17—H17B \cdots O1 ⁱ	0.98	2.51	3.456 (2)	163
C34—H34B \cdots O2 ⁱ	0.98	2.36	3.328 (2)	169

Symmetry code: (i) $x + 1, y, z$.

Data collection: *COLLECT* (Hooft, 2004); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *PLATON* (Spek, 2009).

The authors thank Professor Peter Klüfers for generous allocation of diffractometer time.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FL2253).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst. B* **46**, 256–262.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hooft, R. W. W. (2004). *COLLECT*. Bruker–Nonius BV, Delft, The Netherlands.
- Kawai, H., Nagasu, T., Takeda, T., Fujiwara, K., Tsuji, T., Ohkita, M., Nishida, J. & Suzuki, T. (2004). *Tetrahedron Lett.* **45**, 4553–4558.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Richter, D., Hampel, N., Singer, T., Ofial, A. R. & Mayr, H. (2009). *Eur. J. Org. Chem.* pp. 3203–3211.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2009). E65, o2102 [doi:10.1107/S1600536809030748]

4-[4-(Dimethylamino)benzylidene]-2,6-dimethylcyclohexa-2,5-dienone

Nathalie Hampel, Dorothea Richter, Armin R. Ofial, Herbert Mayr and Peter Mayer

S1. Comment

The asymmetric unit of (I) contains two complete molecules of the title compound. Figure 1 shows one of the two independent molecules.

A major difference between the two symmetrically independent molecules is found in the angle formed by the planes of the two C₆-rings within a molecule. This angle is found to be 35.19 (7)^o between the planes in one molecule but only 20.00 (7)^o between the planes in the other molecule (Fig. 2). With a bulky naphthyl substituent at the C atom linking the two rings, an angle of 43.16 (6)^o is observed [Kawai *et al.* (2004)].


The molecular packing, which is shown in Figure 3, is dominated by two C—H···O hydrogen bonds leading to infinite chains along [100]. Each of the chains is built up by 13 atoms and contains one donor atom and one acceptor atom. According to graph set theory [Bernstein *et al.* (1995), Etter *et al.* (1990)] the descriptor C₁¹(13)C₁¹(13) can be assigned. The strands are cross-linked by very weak C—H···C contacts with H···C distances of at least 2.79 Å (Table 1).

S2. Experimental


The title compound was prepared under an atmosphere of dry N₂ from (4-hydroxy-3,5-dimethylphenyl)[4-(dimethylamino)phenyl]methanol [Richter *et al.* (2009)] (200 mg, 0.737 mmol) that was dissolved in dry CH₂Cl₂ (60 ml) and cooled to 0 °C. Then ethereal HBF₄-solution (0.110 ml, 0.811 mmol) was added at 0 °C. After 5 min, NEt₃ (0.133 ml, 0.958 mmol) was added. The cooling bath was removed and stirring was continued for 3 h before the mixture was washed with water (3 times). The organic layer was dried (MgSO₄) and the solvent was removed under reduced pressure. Crystals were obtained by slow cooling of a warm solution of the title compound in acetonitrile. Yield: 149 mg (80%), mp 127–128 °C.

S3. Refinement


The H atoms were positioned geometrically (C—H = 0.98 Å for CH₃, 0.95 Å for CH) and treated as riding on their parent atoms [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH, $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for CH₃].

**Figure 1**

The molecular structure of the title compound (one molecule out of two of the asymmetric unit), with atom labels and anisotropic displacement ellipsoids (drawn at 50% probability level) for non-H atoms.

**Figure 2**

Schematic representation of the different orientations of the aromatic planes in the two molecules, viewed along the C–H bond of the linking group and parallel to the dienone ring planes. For clarity all atoms connected to the aromatic rings have been omitted.

**Figure 3**

The packing of the title compound, viewed along [010]. Dashed lines indicate hydrogen bonds of the type C—H···O. The solid, red bonds link the 13 atoms involved in one of the two different hydrogen bonds leading to chains along [100].

4-[4-(Dimethylamino)benzylidene]-2,6-dimethylcyclohexa-2,5-dienone

Crystal data

C₁₇H₁₉NO
*M*_r = 253.34
 Monoclinic, *P*2₁/*c*
 Hall symbol: -P 2ybc
a = 14.5357 (3) Å
b = 7.2759 (2) Å
c = 27.5473 (5) Å
 β = 104.6463 (14) $^\circ$
V = 2818.74 (11) Å³
Z = 8

F(000) = 1088
*D*_x = 1.194 (1) Mg m⁻³
 Mo $K\alpha$ radiation, λ = 0.71073 Å
 Cell parameters from 10514 reflections
 θ = 3.1–26.4°
 μ = 0.07 mm⁻¹
T = 200 K
 Block, red
 0.24 × 0.20 × 0.19 mm

Data collection

Nonius KappaCCD
 diffractometer
 Radiation source: rotating anode
 MONTEL, graded multilayered X-ray optics
 monochromator
 Detector resolution: 9 pixels mm⁻¹
 CCD; rotation images; thick slices, ϕ/ω scan
 18635 measured reflections

5725 independent reflections
 3685 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$
 $\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 3.2^\circ$
 $h = -18 \rightarrow 18$
 $k = -9 \rightarrow 9$
 $l = -33 \rightarrow 34$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.049$$

$$wR(F^2) = 0.131$$

$$S = 1.02$$

5725 reflections

351 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.058P)^2 + 0.2713P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on all data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.31746 (8)	0.77657 (19)	0.46546 (5)	0.0626 (4)
N1	0.37909 (9)	0.7322 (2)	0.45672 (5)	0.0497 (4)
C1	-0.24976 (11)	0.7798 (2)	0.44563 (6)	0.0428 (4)
C2	-0.26523 (11)	0.7680 (2)	0.39087 (6)	0.0441 (4)
C3	-0.19006 (11)	0.7670 (2)	0.37098 (6)	0.0425 (4)
H3	-0.2014	0.7639	0.3355	0.051*
C4	-0.09286 (11)	0.7703 (2)	0.40056 (6)	0.0374 (4)
C5	-0.07886 (11)	0.7914 (2)	0.45418 (5)	0.0372 (4)
H5	-0.0157	0.7992	0.4747	0.045*
C6	-0.15154 (11)	0.8003 (2)	0.47623 (5)	0.0372 (4)
C7	-0.36678 (12)	0.7568 (3)	0.35963 (7)	0.0621 (5)
H7A	-0.3675	0.7544	0.3240	0.093*
H7B	-0.4022	0.8642	0.3664	0.093*
H7C	-0.3964	0.6446	0.3682	0.093*
C8	-0.13802 (12)	0.8369 (2)	0.53118 (6)	0.0477 (4)
H8A	-0.0699	0.8458	0.5475	0.072*
H8B	-0.1660	0.7362	0.5463	0.072*
H8C	-0.1693	0.9527	0.5357	0.072*
C9	-0.02032 (11)	0.7623 (2)	0.37695 (6)	0.0394 (4)
H9	-0.0399	0.7699	0.3414	0.047*
C10	0.08090 (10)	0.7442 (2)	0.39752 (5)	0.0350 (4)
C11	0.14350 (11)	0.8184 (2)	0.37151 (5)	0.0395 (4)
H11	0.1179	0.8716	0.3394	0.047*
C12	0.24043 (11)	0.8172 (2)	0.39054 (6)	0.0413 (4)
H12	0.2798	0.8735	0.3720	0.050*

C13	0.28220 (11)	0.7344 (2)	0.43683 (6)	0.0385 (4)
C14	0.22023 (11)	0.6473 (2)	0.46155 (5)	0.0395 (4)
H14	0.2461	0.5826	0.4919	0.047*
C15	0.12351 (11)	0.6539 (2)	0.44275 (5)	0.0370 (4)
H15	0.0840	0.5956	0.4608	0.044*
C16	0.41687 (13)	0.6822 (3)	0.50911 (7)	0.0713 (6)
H16A	0.3946	0.7704	0.5305	0.107*
H16B	0.4865	0.6838	0.5173	0.107*
H16C	0.3949	0.5586	0.5149	0.107*
C17	0.43927 (13)	0.8433 (3)	0.43384 (8)	0.0752 (6)
H17A	0.4262	0.8142	0.3980	0.113*
H17B	0.5061	0.8173	0.4500	0.113*
H17C	0.4262	0.9737	0.4380	0.113*
O2	-0.37910 (8)	0.25472 (19)	0.34392 (5)	0.0634 (4)
N2	0.33071 (9)	0.24688 (19)	0.33231 (5)	0.0443 (4)
C18	-0.30784 (12)	0.2279 (2)	0.32776 (6)	0.0444 (4)
C19	-0.31707 (11)	0.1739 (2)	0.27539 (6)	0.0410 (4)
C20	-0.23808 (10)	0.1486 (2)	0.25927 (6)	0.0386 (4)
H20	-0.2453	0.1116	0.2254	0.046*
C21	-0.14294 (10)	0.1744 (2)	0.29038 (5)	0.0348 (4)
C22	-0.13547 (11)	0.2196 (2)	0.34218 (6)	0.0387 (4)
H22	-0.0739	0.2329	0.3641	0.046*
C23	-0.21173 (11)	0.2438 (2)	0.36091 (6)	0.0404 (4)
C24	-0.41593 (11)	0.1501 (3)	0.24209 (7)	0.0547 (5)
H24A	-0.4121	0.1181	0.2081	0.082*
H24B	-0.4515	0.2651	0.2411	0.082*
H24C	-0.4485	0.0516	0.2554	0.082*
C25	-0.20350 (13)	0.2812 (3)	0.41543 (6)	0.0535 (5)
H25A	-0.1363	0.2953	0.4331	0.080*
H25B	-0.2309	0.1783	0.4300	0.080*
H25C	-0.2379	0.3944	0.4188	0.080*
C26	-0.06843 (10)	0.1560 (2)	0.26859 (5)	0.0348 (4)
H26	-0.0867	0.1124	0.2350	0.042*
C27	0.03223 (10)	0.1899 (2)	0.28665 (5)	0.0322 (4)
C28	0.09255 (10)	0.1161 (2)	0.25908 (5)	0.0344 (4)
H28	0.0648	0.0509	0.2291	0.041*
C29	0.18928 (10)	0.1334 (2)	0.27339 (5)	0.0356 (4)
H29	0.2268	0.0777	0.2538	0.043*
C30	0.23412 (10)	0.2326 (2)	0.31684 (5)	0.0337 (4)
C31	0.17424 (10)	0.3170 (2)	0.34342 (5)	0.0352 (4)
H31	0.2015	0.3904	0.3720	0.042*
C32	0.07772 (10)	0.2950 (2)	0.32885 (5)	0.0340 (4)
H32	0.0399	0.3529	0.3479	0.041*
C33	0.37588 (11)	0.3514 (3)	0.37678 (6)	0.0547 (5)
H33A	0.3545	0.4795	0.3724	0.082*
H33B	0.4451	0.3468	0.3820	0.082*
H33C	0.3586	0.2987	0.4060	0.082*
C34	0.39029 (11)	0.1576 (3)	0.30456 (7)	0.0613 (5)

H34A	0.3789	0.0247	0.3038	0.092*
H34B	0.4572	0.1823	0.3209	0.092*
H34C	0.3751	0.2051	0.2702	0.092*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0423 (7)	0.0853 (11)	0.0637 (8)	-0.0013 (7)	0.0201 (6)	0.0060 (7)
N1	0.0368 (8)	0.0599 (10)	0.0510 (9)	-0.0005 (7)	0.0087 (7)	-0.0015 (7)
C1	0.0380 (10)	0.0389 (10)	0.0521 (10)	0.0030 (7)	0.0124 (8)	0.0058 (8)
C2	0.0385 (10)	0.0414 (11)	0.0488 (10)	0.0007 (8)	0.0045 (8)	0.0039 (8)
C3	0.0431 (10)	0.0449 (11)	0.0361 (9)	0.0003 (8)	0.0036 (7)	0.0007 (7)
C4	0.0378 (9)	0.0333 (9)	0.0386 (9)	0.0011 (7)	0.0053 (7)	-0.0010 (7)
C5	0.0364 (9)	0.0357 (10)	0.0375 (9)	0.0031 (7)	0.0055 (7)	-0.0011 (7)
C6	0.0435 (10)	0.0293 (9)	0.0387 (9)	0.0022 (7)	0.0099 (7)	0.0004 (7)
C7	0.0396 (10)	0.0792 (15)	0.0615 (12)	-0.0030 (10)	0.0018 (8)	0.0074 (10)
C8	0.0514 (11)	0.0480 (11)	0.0455 (10)	0.0000 (8)	0.0158 (8)	-0.0053 (8)
C9	0.0430 (10)	0.0396 (10)	0.0343 (8)	0.0027 (7)	0.0075 (7)	-0.0001 (7)
C10	0.0376 (9)	0.0342 (9)	0.0333 (8)	0.0021 (7)	0.0093 (7)	-0.0037 (7)
C11	0.0481 (11)	0.0388 (10)	0.0329 (8)	0.0061 (8)	0.0129 (7)	0.0021 (7)
C12	0.0440 (10)	0.0403 (10)	0.0444 (9)	0.0017 (8)	0.0199 (8)	0.0006 (8)
C13	0.0370 (9)	0.0378 (10)	0.0410 (9)	0.0029 (7)	0.0108 (7)	-0.0065 (7)
C14	0.0435 (10)	0.0419 (10)	0.0330 (8)	0.0071 (8)	0.0096 (7)	0.0011 (7)
C15	0.0418 (10)	0.0361 (10)	0.0356 (8)	0.0032 (7)	0.0144 (7)	0.0003 (7)
C16	0.0454 (11)	0.1020 (18)	0.0574 (12)	0.0055 (11)	-0.0036 (9)	-0.0013 (11)
C17	0.0449 (12)	0.0843 (17)	0.0953 (16)	-0.0098 (11)	0.0155 (11)	0.0064 (13)
O2	0.0438 (7)	0.0777 (10)	0.0766 (9)	-0.0019 (6)	0.0296 (7)	-0.0172 (7)
N2	0.0300 (8)	0.0539 (10)	0.0487 (8)	-0.0022 (6)	0.0097 (6)	-0.0082 (7)
C18	0.0397 (10)	0.0411 (11)	0.0572 (11)	-0.0015 (8)	0.0210 (8)	-0.0019 (8)
C19	0.0327 (9)	0.0379 (10)	0.0519 (10)	-0.0006 (7)	0.0099 (7)	-0.0001 (8)
C20	0.0365 (9)	0.0383 (10)	0.0398 (9)	-0.0006 (7)	0.0073 (7)	-0.0029 (7)
C21	0.0339 (9)	0.0324 (9)	0.0375 (8)	0.0014 (7)	0.0081 (7)	0.0002 (7)
C22	0.0337 (9)	0.0408 (10)	0.0400 (9)	-0.0033 (7)	0.0066 (7)	0.0003 (7)
C23	0.0436 (10)	0.0369 (10)	0.0437 (9)	-0.0036 (7)	0.0165 (8)	-0.0023 (7)
C24	0.0359 (10)	0.0649 (13)	0.0630 (11)	-0.0020 (9)	0.0117 (8)	-0.0051 (9)
C25	0.0613 (12)	0.0562 (13)	0.0482 (10)	-0.0044 (9)	0.0237 (9)	-0.0043 (9)
C26	0.0331 (9)	0.0354 (9)	0.0345 (8)	0.0013 (7)	0.0060 (7)	-0.0011 (7)
C27	0.0324 (9)	0.0325 (9)	0.0314 (8)	0.0024 (7)	0.0074 (6)	0.0025 (6)
C28	0.0370 (9)	0.0363 (9)	0.0296 (8)	-0.0012 (7)	0.0080 (6)	0.0004 (6)
C29	0.0365 (9)	0.0372 (10)	0.0366 (8)	0.0007 (7)	0.0159 (7)	-0.0013 (7)
C30	0.0284 (8)	0.0368 (10)	0.0360 (8)	0.0004 (7)	0.0084 (6)	0.0042 (7)
C31	0.0357 (9)	0.0363 (10)	0.0327 (8)	-0.0024 (7)	0.0070 (7)	-0.0022 (7)
C32	0.0335 (9)	0.0352 (9)	0.0342 (8)	0.0021 (7)	0.0104 (7)	-0.0008 (7)
C33	0.0362 (10)	0.0668 (14)	0.0552 (10)	-0.0054 (9)	0.0006 (8)	-0.0090 (9)
C34	0.0337 (10)	0.0803 (15)	0.0735 (13)	-0.0009 (9)	0.0201 (9)	-0.0154 (11)

Geometric parameters (\AA , $\text{^{\circ}}$)

O1—C1	1.2406 (18)	O2—C18	1.2418 (18)
N1—C13	1.376 (2)	N2—C30	1.3642 (18)
N1—C17	1.447 (2)	N2—C34	1.446 (2)
N1—C16	1.455 (2)	N2—C33	1.450 (2)
C1—C2	1.470 (2)	C18—C23	1.468 (2)
C1—C6	1.471 (2)	C18—C19	1.468 (2)
C2—C3	1.341 (2)	C19—C20	1.344 (2)
C2—C7	1.511 (2)	C19—C24	1.507 (2)
C3—C4	1.442 (2)	C20—C21	1.443 (2)
C3—H3	0.9500	C20—H20	0.9500
C4—C9	1.373 (2)	C21—C26	1.371 (2)
C4—C5	1.447 (2)	C21—C22	1.441 (2)
C5—C6	1.347 (2)	C22—C23	1.347 (2)
C5—H5	0.9500	C22—H22	0.9500
C6—C8	1.500 (2)	C23—C25	1.501 (2)
C7—H7A	0.9800	C24—H24A	0.9800
C7—H7B	0.9800	C24—H24B	0.9800
C7—H7C	0.9800	C24—H24C	0.9800
C8—H8A	0.9800	C25—H25A	0.9800
C8—H8B	0.9800	C25—H25B	0.9800
C8—H8C	0.9800	C25—H25C	0.9800
C9—C10	1.443 (2)	C26—C27	1.4426 (19)
C9—H9	0.9500	C26—H26	0.9500
C10—C11	1.401 (2)	C27—C28	1.404 (2)
C10—C15	1.406 (2)	C27—C32	1.409 (2)
C11—C12	1.374 (2)	C28—C29	1.367 (2)
C11—H11	0.9500	C28—H28	0.9500
C12—C13	1.402 (2)	C29—C30	1.409 (2)
C12—H12	0.9500	C29—H29	0.9500
C13—C14	1.410 (2)	C30—C31	1.412 (2)
C14—C15	1.371 (2)	C31—C32	1.368 (2)
C14—H14	0.9500	C31—H31	0.9500
C15—H15	0.9500	C32—H32	0.9500
C16—H16A	0.9800	C33—H33A	0.9800
C16—H16B	0.9800	C33—H33B	0.9800
C16—H16C	0.9800	C33—H33C	0.9800
C17—H17A	0.9800	C34—H34A	0.9800
C17—H17B	0.9800	C34—H34B	0.9800
C17—H17C	0.9800	C34—H34C	0.9800
C13—N1—C17	119.38 (14)	C30—N2—C34	120.48 (13)
C13—N1—C16	119.30 (14)	C30—N2—C33	120.93 (13)
C17—N1—C16	116.99 (14)	C34—N2—C33	118.59 (13)
O1—C1—C2	121.20 (15)	O2—C18—C23	120.91 (15)
O1—C1—C6	120.85 (15)	O2—C18—C19	121.12 (15)
C2—C1—C6	117.93 (14)	C23—C18—C19	117.94 (13)

C3—C2—C1	119.39 (14)	C20—C19—C18	119.19 (14)
C3—C2—C7	123.14 (16)	C20—C19—C24	123.09 (15)
C1—C2—C7	117.47 (15)	C18—C19—C24	117.72 (14)
C2—C3—C4	123.54 (15)	C19—C20—C21	123.81 (14)
C2—C3—H3	118.2	C19—C20—H20	118.1
C4—C3—H3	118.2	C21—C20—H20	118.1
C9—C4—C3	119.46 (14)	C26—C21—C22	125.83 (13)
C9—C4—C5	124.14 (14)	C26—C21—C20	118.16 (13)
C3—C4—C5	116.32 (14)	C22—C21—C20	116.00 (13)
C6—C5—C4	122.78 (14)	C23—C22—C21	123.07 (14)
C6—C5—H5	118.6	C23—C22—H22	118.5
C4—C5—H5	118.6	C21—C22—H22	118.5
C5—C6—C1	119.69 (14)	C22—C23—C18	119.83 (14)
C5—C6—C8	123.10 (14)	C22—C23—C25	122.80 (15)
C1—C6—C8	117.18 (14)	C18—C23—C25	117.34 (14)
C2—C7—H7A	109.5	C19—C24—H24A	109.5
C2—C7—H7B	109.5	C19—C24—H24B	109.5
H7A—C7—H7B	109.5	H24A—C24—H24B	109.5
C2—C7—H7C	109.5	C19—C24—H24C	109.5
H7A—C7—H7C	109.5	H24A—C24—H24C	109.5
H7B—C7—H7C	109.5	H24B—C24—H24C	109.5
C6—C8—H8A	109.5	C23—C25—H25A	109.5
C6—C8—H8B	109.5	C23—C25—H25B	109.5
H8A—C8—H8B	109.5	H25A—C25—H25B	109.5
C6—C8—H8C	109.5	C23—C25—H25C	109.5
H8A—C8—H8C	109.5	H25A—C25—H25C	109.5
H8B—C8—H8C	109.5	H25B—C25—H25C	109.5
C4—C9—C10	130.31 (14)	C21—C26—C27	132.61 (14)
C4—C9—H9	114.8	C21—C26—H26	113.7
C10—C9—H9	114.8	C27—C26—H26	113.7
C11—C10—C15	115.83 (13)	C28—C27—C32	115.29 (13)
C11—C10—C9	119.57 (13)	C28—C27—C26	117.75 (13)
C15—C10—C9	124.60 (14)	C32—C27—C26	126.93 (13)
C12—C11—C10	122.60 (14)	C29—C28—C27	123.11 (13)
C12—C11—H11	118.7	C29—C28—H28	118.4
C10—C11—H11	118.7	C27—C28—H28	118.4
C11—C12—C13	121.08 (14)	C28—C29—C30	120.92 (13)
C11—C12—H12	119.5	C28—C29—H29	119.5
C13—C12—H12	119.5	C30—C29—H29	119.5
N1—C13—C12	122.09 (14)	N2—C30—C29	121.54 (13)
N1—C13—C14	121.14 (14)	N2—C30—C31	121.71 (13)
C12—C13—C14	116.74 (14)	C29—C30—C31	116.75 (13)
C15—C14—C13	121.43 (14)	C32—C31—C30	121.21 (13)
C15—C14—H14	119.3	C32—C31—H31	119.4
C13—C14—H14	119.3	C30—C31—H31	119.4
C14—C15—C10	122.08 (14)	C31—C32—C27	122.56 (13)
C14—C15—H15	119.0	C31—C32—H32	118.7
C10—C15—H15	119.0	C27—C32—H32	118.7

N1—C16—H16A	109.5	N2—C33—H33A	109.5
N1—C16—H16B	109.5	N2—C33—H33B	109.5
H16A—C16—H16B	109.5	H33A—C33—H33B	109.5
N1—C16—H16C	109.5	N2—C33—H33C	109.5
H16A—C16—H16C	109.5	H33A—C33—H33C	109.5
H16B—C16—H16C	109.5	H33B—C33—H33C	109.5
N1—C17—H17A	109.5	N2—C34—H34A	109.5
N1—C17—H17B	109.5	N2—C34—H34B	109.5
H17A—C17—H17B	109.5	H34A—C34—H34B	109.5
N1—C17—H17C	109.5	N2—C34—H34C	109.5
H17A—C17—H17C	109.5	H34A—C34—H34C	109.5
H17B—C17—H17C	109.5	H34B—C34—H34C	109.5
O1—C1—C2—C3	-178.22 (16)	O2—C18—C19—C20	179.15 (16)
C6—C1—C2—C3	3.3 (2)	C23—C18—C19—C20	-2.7 (2)
O1—C1—C2—C7	1.5 (2)	O2—C18—C19—C24	-0.6 (2)
C6—C1—C2—C7	-176.94 (15)	C23—C18—C19—C24	177.49 (15)
C1—C2—C3—C4	2.3 (3)	C18—C19—C20—C21	-1.1 (2)
C7—C2—C3—C4	-177.45 (16)	C24—C19—C20—C21	178.72 (15)
C2—C3—C4—C9	177.69 (16)	C19—C20—C21—C26	-175.74 (15)
C2—C3—C4—C5	-5.3 (2)	C19—C20—C21—C22	3.6 (2)
C9—C4—C5—C6	179.44 (15)	C26—C21—C22—C23	176.92 (16)
C3—C4—C5—C6	2.6 (2)	C20—C21—C22—C23	-2.4 (2)
C4—C5—C6—C1	2.9 (2)	C21—C22—C23—C18	-1.3 (2)
C4—C5—C6—C8	-175.02 (15)	C21—C22—C23—C25	176.74 (16)
O1—C1—C6—C5	175.68 (15)	O2—C18—C23—C22	-178.00 (16)
C2—C1—C6—C5	-5.8 (2)	C19—C18—C23—C22	3.9 (2)
O1—C1—C6—C8	-6.3 (2)	O2—C18—C23—C25	3.9 (2)
C2—C1—C6—C8	172.17 (15)	C19—C18—C23—C25	-174.24 (15)
C3—C4—C9—C10	-172.71 (15)	C22—C21—C26—C27	-6.6 (3)
C5—C4—C9—C10	10.5 (3)	C20—C21—C26—C27	172.68 (15)
C4—C9—C10—C11	-151.45 (17)	C21—C26—C27—C28	165.06 (16)
C4—C9—C10—C15	29.2 (3)	C21—C26—C27—C32	-16.9 (3)
C15—C10—C11—C12	-5.2 (2)	C32—C27—C28—C29	4.1 (2)
C9—C10—C11—C12	175.42 (14)	C26—C27—C28—C29	-177.62 (13)
C10—C11—C12—C13	2.4 (2)	C27—C28—C29—C30	-1.7 (2)
C17—N1—C13—C12	10.3 (2)	C34—N2—C30—C29	-0.8 (2)
C16—N1—C13—C12	166.30 (17)	C33—N2—C30—C29	179.24 (15)
C17—N1—C13—C14	-171.83 (16)	C34—N2—C30—C31	179.45 (15)
C16—N1—C13—C14	-15.9 (2)	C33—N2—C30—C31	-0.5 (2)
C11—C12—C13—N1	-179.73 (15)	C28—C29—C30—N2	178.22 (14)
C11—C12—C13—C14	2.3 (2)	C28—C29—C30—C31	-2.0 (2)
N1—C13—C14—C15	177.92 (14)	N2—C30—C31—C32	-177.05 (14)
C12—C13—C14—C15	-4.1 (2)	C29—C30—C31—C32	3.2 (2)
C13—C14—C15—C10	1.3 (2)	C30—C31—C32—C27	-0.7 (2)
C11—C10—C15—C14	3.3 (2)	C28—C27—C32—C31	-2.9 (2)
C9—C10—C15—C14	-177.29 (14)	C26—C27—C32—C31	179.02 (14)

Hydrogen-bond geometry (Å, °)

$D\text{---}H\cdots A$	$D\text{---}H$	$H\cdots A$	$D\cdots A$	$D\text{---}H\cdots A$
C17—H17 <i>B</i> ···O1 ⁱ	0.98	2.51	3.456 (2)	163
C34—H34 <i>B</i> ···O2 ⁱ	0.98	2.36	3.328 (2)	169

Symmetry code: (i) $x+1, y, z$.