

trans-Dioxidotetrapyridinerhenium(V) triiodide

Miłosz Siczek, Marta S. Krawczyk* and Tadeusz Lis

University of Wrocław, Faculty of Chemistry, 14 Joliot-Curie St, 50-383 Wrocław, Poland

Correspondence e-mail: martakrawczyk.re@gmail.com


Received 31 July 2009; accepted 2 August 2009

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.019; wR factor = 0.031; data-to-parameter ratio = 30.9.

In the title salt, $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]\text{I}_3$, the cation and anion are both located on centres of symmetry. The Re^{V} atom adopts a *trans*- ReO_2N_4 octahedral coordination and short intramolecular $\text{C}-\text{H}\cdots\text{O}$ contacts occur within the cation. In the crystal, the cations form layers perpendicular to $[100]$ and a weak $\text{C}-\text{H}\cdots\text{O}$ interaction links the cations.

Related literature

For related structures containing the same cation, see: Calvo *et al.* (1971); Lock & Turner (1978); Luck & O'Neill (2001). For further synthetic details, see: Johnson *et al.* (1967). For background to aromatic π - π stacking, see: Janiak (2000).



Experimental

Crystal data

$[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]\text{I}_3$
 $M_r = 915.30$
 Triclinic, $P\bar{1}$
 $a = 7.993$ (3) Å
 $b = 9.100$ (3) Å
 $c = 9.356$ (3) Å
 $\alpha = 92.45$ (4)°
 $\beta = 102.41$ (4)°

$\gamma = 104.10$ (4)°
 $V = 641.3$ (4) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 8.37$ mm⁻¹
 $T = 100$ K
 $0.10 \times 0.10 \times 0.07$ mm

Data collection

Oxford Diffraction Xcalibur PX
 KM-4-CCD diffractometer
 Absorption correction: analytical
 (*CrysAlis RED*; Oxford
 Diffraction, 2006)
 $T_{\text{min}} = 0.411$, $T_{\text{max}} = 0.656$

11244 measured reflections
 4298 independent reflections
 3593 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.019$
 $wR(F^2) = 0.031$
 $S = 1.04$
 4298 reflections

139 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.40$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.08$ e Å⁻³

Table 1

Selected bond lengths (Å).

Re—O	1.7649 (18)	Re—N2	2.1442 (18)
Re—N1	2.1411 (19)	I1—I2	2.9222 (12)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C15—H15 \cdots O	0.95	2.39	2.914 (3)	114
C25—H25 \cdots O	0.95	2.38	2.906 (3)	115
C11—H11 \cdots O ⁱ	0.95	2.39	2.913 (3)	115
C21—H21 \cdots O ⁱ	0.95	2.37	2.908 (3)	115
C22—H22 \cdots O ⁱⁱ	0.95	2.41	3.309 (3)	157

 Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and *pubCIF* (Westrip, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5027).

References

- Calvo, C., Krishnamachari, N. & Lock, C. J. L. (1971). *J. Cryst. Mol. Struct.* **1**, 161–172.
 Janiak, C. (2000). *J. Chem. Soc. Dalton Trans.* pp. 3885–3896.
 Johnson, N. P., Lock, C. J. L. & Wilkinson, G. (1967). *Inorg. Synth.* **9**, 145–148.
 Lock, C. J. L. & Turner, G. (1978). *Acta Cryst.* **B34**, 923–927.
 Luck, R. L. & O'Neill, R. S. (2001). *Polyhedron*, **28**, 773–782.
 Oxford Diffraction (2006). *CrysAlis RED* and *CrysAlis CCD*. Oxford Diffraction Poland, Wrocław, Poland.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). *pubCIF*. In preparation.

supporting information

Acta Cryst. (2009). E65, m1057 [doi:10.1107/S1600536809030724]

***trans*-Dioxidotetrapyridinerhenium(V) triiodide**

Miłosz Siczek, Marta S. Krawczyk and Tadeusz Lis

S1. Comment

The crystal structure of a salt containing $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]^+$ cation was first investigated by Calvo *et al.*, (1971). The authors obtained dioxidotetra(pyridine)rhenium(V) chloride dihydrate in the reaction between trichloridooxidobis(tri-phenylphosphine)rhenium(V) (Johnson *et al.*, 1967) and hot pyridine used in excess. The crystal structure of $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]\text{Cl}\cdot 2\text{H}_2\text{O}$ was redetermined by Lock & Turner (1978). The cation $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]^+$ was also described by Luck & O'Neill (2001) as $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4[\text{OH}]\cdot 1.75\text{H}_2\text{O}$ salt. This salt was prepared by dissolving $\text{ReCl}(\text{H}_2)(\text{PMePh}_2)_4$ in the mixture of benzene, pyridine, water and hexane.

The crystal structure of *trans*-dioxidotetra(pyridine)rhenium(V) triiodide comprises of $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]^+$ cations and I_3^- anions (Fig. 1). Both ions are located on centres of symmetry. The cation is a distorted octahedron, with two *oxido* (terminal) ligands in *trans* arrangement and four pyridine ligands in equatorial positions.

The average Re—O and Re—N bond distances equal 1.765 (2), 2.143 (2) Å, respectively, and are in good agreement with values reported by Calvo *et al.*, (1971), Lock & Turner (1978) and Luck & O'Neill (2001). Moreover, comparing the values of O—Re—O angle comparatively small differences between previous and present results can be observed. In the crystal structure reported here this angle equals 180° and reported for other salts is 171 (1)° (Calvo *et al.*, 1971) and 174.5 (4)° (Lock & Turner, 1978). Similarly, the value of N—Re—N_{*trans*} angles in $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]\text{I}_3$ equals 180° and the analogous complex cations that have been determined previously have near linear arrangement of the N—Re—N_{*trans*} moiety. These angles are 176 (2) and 170 (1)° (Calvo *et al.*, 1971), and 173.9 (4) and 175.2 (6)° (Lock & Turner, 1978). The comparatively weak intramolecular hydrogen bonds such as C—H...O can be observed (Fig. 2, Table 2).

The molecular packing in the crystal structure can be described as layers perpendicular to [100] direction which consist of the complex cations (Fig. 3). The I_3^- anions are located between the layers of $[\text{ReO}_2(\text{C}_5\text{H}_5\text{N})_4]^+$ cations. In the crystal packing there are intermolecular stacking interactions between pyridine rings with centroid-centroid distance of 3.831 (2) Å and a slip angle 25°. These values are comparable with the corresponding values reported for transition-metal pyridine fragments (Janiak, 2000). (The ring centroid contacts range between 3.4 and 3.8 Å and the angle averages 27°).

S2. Experimental

Rhenium(III) iodide 0.2982 g (0.1753 mmol) was refluxed in dry pyridine (5 ml) (62 mmol) for 3 h at 423 K. The mixture was allowed to evaporate in air at high temperature to give a greenish brown precipitate. The complex was recrystallized from methanol to yield orange blocks of (I).

S3. Refinement

All hydrogen atoms were placed in calculated positions and refined using riding model [C—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. The highest peak and the deepest hole in the final difference map were 1.07 Å from N1 and 0.78 Å from Re, respectively.




Figure 1

The molecular structure of (I) showing ellipsoids drawn at the 30% probability level. The unlabelled atoms of the cation are generated by the symmetry operation $(1-x, 1-y, 1-z)$ and the unlabelled I atom by $(2-x, 2-y, 2-z)$.




Figure 2

A part of the crystal structure showing formation of C—H...O hydrogen bonding. [symmetry code (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$.]




Figure 3

A packing diagram of (I) showing layers of cations and anions. Hydrogen atoms are omitted for clarity.

trans-dioxidotetra(pyridine)rhenium(V) triiodide

Crystal data

[ReO₂(C₅H₅N)₄]₃I₃

$M_r = 915.30$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.993$ (3) Å

$b = 9.100$ (3) Å

$c = 9.356$ (3) Å

$\alpha = 92.45$ (4)°

$\beta = 102.41$ (4)°

$\gamma = 104.10$ (4)°

$V = 641.3$ (4) Å³

$Z = 1$

$F(000) = 418$

$D_x = 2.370$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 11676 reflections

$\theta = 4.5\text{--}38.4^\circ$

$\mu = 8.37$ mm⁻¹

$T = 100$ K

Block, orange

$0.10 \times 0.10 \times 0.07$ mm

Data collection

Oxford Diffraction Xcalibur PX KM-4-CCD diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: analytical

(*CrysAlis RED*; Oxford Diffraction, 2006)

$T_{\min} = 0.411$, $T_{\max} = 0.656$

11244 measured reflections

4298 independent reflections

3593 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 32.5^\circ$, $\theta_{\min} = 4.5^\circ$

$h = -12 \rightarrow 11$

$k = -9 \rightarrow 13$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.019$

$wR(F^2) = 0.031$

$S = 1.04$

4298 reflections

139 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.008P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.003$
 $\Delta\rho_{\max} = 1.40 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -1.08 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Re	0.5000	0.5000	0.5000	0.01065 (3)
I1	0.92874 (2)	1.171742 (18)	0.747033 (17)	0.02194 (4)
I2	1.0000	1.0000	1.0000	0.01924 (5)
O	0.7122 (2)	0.62887 (16)	0.53698 (16)	0.0144 (3)
N1	0.5561 (2)	0.4374 (2)	0.71983 (19)	0.0132 (4)
C11	0.4386 (3)	0.3225 (2)	0.7640 (2)	0.0159 (5)
H11	0.3318	0.2707	0.6957	0.019*
C12	0.4708 (3)	0.2798 (3)	0.9045 (3)	0.0213 (5)
H12	0.3873	0.1988	0.9315	0.026*
C13	0.6234 (3)	0.3541 (3)	1.0059 (3)	0.0225 (6)
H13	0.6468	0.3251	1.1031	0.027*
C14	0.7421 (3)	0.4718 (3)	0.9631 (3)	0.0223 (5)
H14	0.8480	0.5260	1.0310	0.027*
C15	0.7045 (3)	0.5096 (3)	0.8203 (2)	0.0176 (5)
H15	0.7872	0.5903	0.7918	0.021*
N2	0.3956 (2)	0.6744 (2)	0.58055 (19)	0.0126 (4)
C21	0.2226 (3)	0.6460 (3)	0.5840 (2)	0.0148 (5)
H21	0.1481	0.5467	0.5506	0.018*
C22	0.1494 (3)	0.7552 (3)	0.6340 (2)	0.0173 (5)
H22	0.0268	0.7312	0.6337	0.021*
C23	0.2564 (3)	0.8991 (3)	0.6844 (3)	0.0196 (5)
H23	0.2092	0.9762	0.7192	0.024*
C24	0.4346 (3)	0.9288 (3)	0.6833 (3)	0.0229 (6)
H24	0.5119	1.0265	0.7190	0.027*
C25	0.4992 (3)	0.8152 (2)	0.6299 (3)	0.0189 (5)
H25	0.6211	0.8374	0.6279	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Re	0.00779 (7)	0.01136 (7)	0.01158 (6)	0.00084 (5)	0.00184 (5)	-0.00043 (5)
I1	0.01743 (9)	0.02567 (9)	0.02152 (8)	0.00406 (7)	0.00338 (7)	0.00427 (7)
I2	0.01598 (12)	0.02278 (12)	0.01852 (11)	0.00268 (10)	0.00608 (9)	0.00035 (9)
O	0.0099 (8)	0.0144 (8)	0.0166 (8)	0.0008 (7)	0.0020 (7)	-0.0025 (7)
N1	0.0123 (10)	0.0131 (9)	0.0145 (9)	0.0044 (8)	0.0026 (8)	-0.0003 (8)
C11	0.0131 (12)	0.0146 (11)	0.0182 (11)	-0.0002 (10)	0.0043 (10)	-0.0013 (9)
C12	0.0291 (15)	0.0172 (12)	0.0199 (12)	0.0060 (12)	0.0101 (11)	0.0051 (10)
C13	0.0314 (16)	0.0256 (13)	0.0152 (11)	0.0152 (13)	0.0064 (11)	0.0039 (10)
C14	0.0191 (14)	0.0289 (14)	0.0170 (12)	0.0088 (12)	-0.0019 (10)	-0.0021 (11)
C15	0.0141 (12)	0.0178 (12)	0.0181 (11)	0.0005 (10)	0.0029 (10)	-0.0013 (10)
N2	0.0098 (10)	0.0145 (9)	0.0123 (9)	0.0015 (8)	0.0019 (8)	0.0004 (8)
C21	0.0110 (12)	0.0148 (11)	0.0144 (11)	-0.0006 (10)	-0.0009 (9)	-0.0027 (9)
C22	0.0112 (12)	0.0230 (13)	0.0182 (11)	0.0046 (11)	0.0040 (10)	0.0020 (10)
C23	0.0192 (13)	0.0187 (12)	0.0236 (12)	0.0080 (11)	0.0076 (11)	-0.0006 (10)
C24	0.0171 (13)	0.0140 (12)	0.0361 (14)	0.0007 (11)	0.0087 (12)	-0.0061 (11)
C25	0.0127 (12)	0.0163 (12)	0.0268 (13)	0.0003 (10)	0.0067 (10)	-0.0006 (10)

Geometric parameters (\AA , $^\circ$)

Re—O ⁱ	1.7649 (18)	C13—H13	0.9500
Re—O	1.7649 (18)	C14—C15	1.382 (3)
Re—N1 ⁱ	2.1411 (19)	C14—H14	0.9500
Re—N1	2.1411 (19)	C15—H15	0.9500
Re—N2 ⁱ	2.1442 (18)	N2—C25	1.344 (3)
Re—N2	2.1442 (18)	N2—C21	1.351 (3)
I1—I2	2.9222 (12)	C21—C22	1.382 (3)
I2—I1 ⁱⁱ	2.9222 (12)	C21—H21	0.9500
N1—C15	1.346 (3)	C22—C23	1.378 (3)
N1—C11	1.368 (3)	C22—H22	0.9500
C11—C12	1.375 (3)	C23—C24	1.386 (3)
C11—H11	0.9500	C23—H23	0.9500
C12—C13	1.375 (4)	C24—C25	1.383 (3)
C12—H12	0.9500	C24—H24	0.9500
C13—C14	1.383 (3)	C25—H25	0.9500
O ⁱ —Re—O	180.0	C12—C13—H13	120.8
O ⁱ —Re—N1 ⁱ	89.50 (8)	C14—C13—H13	120.8
O—Re—N1 ⁱ	90.50 (8)	C15—C14—C13	119.2 (3)
O ⁱ —Re—N1	90.50 (8)	C15—C14—H14	120.4
O—Re—N1	89.50 (8)	C13—C14—H14	120.4
N1 ⁱ —Re—N1	180.0	N1—C15—C14	123.0 (2)
O ⁱ —Re—N2 ⁱ	89.76 (7)	N1—C15—H15	118.5
O—Re—N2 ⁱ	90.24 (7)	C14—C15—H15	118.5
N1 ⁱ —Re—N2 ⁱ	88.04 (7)	C25—N2—C21	117.65 (18)
N1—Re—N2 ⁱ	91.96 (7)	C25—N2—Re	121.51 (15)

O ⁱ —Re—N2	90.24 (7)	C21—N2—Re	120.85 (15)
O—Re—N2	89.76 (7)	N2—C21—C22	122.8 (2)
N1 ⁱ —Re—N2	91.96 (7)	N2—C21—H21	118.6
N1—Re—N2	88.04 (7)	C22—C21—H21	118.6
N2 ⁱ —Re—N2	180.0	C23—C22—C21	119.2 (2)
I1—I2—I1 ⁱⁱ	180.0	C23—C22—H22	120.4
C15—N1—C11	117.36 (19)	C21—C22—H22	120.4
C15—N1—Re	122.15 (15)	C22—C23—C24	118.4 (2)
C11—N1—Re	120.48 (16)	C22—C23—H23	120.8
N1—C11—C12	121.8 (2)	C24—C23—H23	120.8
N1—C11—H11	119.1	C25—C24—C23	119.5 (2)
C12—C11—H11	119.1	C25—C24—H24	120.2
C13—C12—C11	120.3 (2)	C23—C24—H24	120.2
C13—C12—H12	119.9	N2—C25—C24	122.4 (2)
C11—C12—H12	119.9	N2—C25—H25	118.8
C12—C13—C14	118.5 (2)	C24—C25—H25	118.8
O ⁱ —Re—N1—C15	-173.72 (16)	O ⁱ —Re—N2—C25	-179.56 (17)
O—Re—N1—C15	6.28 (16)	O—Re—N2—C25	0.44 (17)
N2 ⁱ —Re—N1—C15	96.50 (16)	N1 ⁱ —Re—N2—C25	-90.05 (18)
N2—Re—N1—C15	-83.50 (16)	N1—Re—N2—C25	89.95 (18)
O ⁱ —Re—N1—C11	5.08 (14)	O ⁱ —Re—N2—C21	0.76 (16)
O—Re—N1—C11	-174.92 (14)	O—Re—N2—C21	-179.24 (16)
N2 ⁱ —Re—N1—C11	-84.70 (15)	N1 ⁱ —Re—N2—C21	90.27 (17)
N2—Re—N1—C11	95.30 (15)	N1—Re—N2—C21	-89.73 (17)
C15—N1—C11—C12	-1.0 (3)	C25—N2—C21—C22	0.8 (3)
Re—N1—C11—C12	-179.90 (15)	Re—N2—C21—C22	-179.55 (16)
N1—C11—C12—C13	0.7 (3)	N2—C21—C22—C23	-0.8 (3)
C11—C12—C13—C14	0.2 (3)	C21—C22—C23—C24	-0.2 (3)
C12—C13—C14—C15	-0.7 (3)	C22—C23—C24—C25	1.1 (4)
C11—N1—C15—C14	0.5 (3)	C21—N2—C25—C24	0.2 (3)
Re—N1—C15—C14	179.34 (16)	Re—N2—C25—C24	-179.47 (18)
C13—C14—C15—N1	0.4 (3)	C23—C24—C25—N2	-1.1 (4)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+2, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C15—H15 \cdots O	0.95	2.39	2.914 (3)	114
C25—H25 \cdots O	0.95	2.38	2.906 (3)	115
C11—H11 \cdots O ⁱ	0.95	2.39	2.913 (3)	115
C21—H21 \cdots O ⁱ	0.95	2.37	2.908 (3)	115
C22—H22 \cdots O ⁱⁱⁱ	0.95	2.41	3.309 (3)	157

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (iii) $x-1, y, z$.