

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(1,10-Phenanthroline- κ^2N,N')(triphenylphosphine- κP)silver(I) trifluoromethanesulfonateJie-Qiang Wu,^a Qiong-Hua Jin,^{a*} Ke-Yi Hu^a and Cun-Lin Zhang^b^aDepartment of Chemistry, Capital Normal University, Beijing 100048, People's Republic of China, and ^bDepartment of Physics, Capital Normal University, Beijing 100048, People's Republic of China

Correspondence e-mail: jinqh204@163.com


Received 17 June 2009; accepted 13 August 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.042; wR factor = 0.133; data-to-parameter ratio = 25.1.

The structure of the title complex, $[\text{Ag}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{C}_{18}\text{H}_{15}\text{P})]\text{CF}_3\text{SO}_3$, is based on a distorted trigonal-planar N_2P coordination of the Ag^{I} ion, provided by two N atoms of the bidentate phenanthroline ligand and one P atom of the triphenylphosphine ligand. The phenanthroline ligand and one phenyl ring of the triphenylphosphine ligand almost lie in one plane (maximum deviation = 0.014 Å from the best planes). The crystal structure may be stabilized by an intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond between the phenanthroline ligand and the O atom of the trifluoromethanesulfonate anion.

Related literature

For related structures, see: Di Nicola *et al.* (2007); Jin *et al.* (1999, 2009); Effendy *et al.* (2007a,b); Awaleh *et al.* (2005a,b); Pettinari *et al.* (2007). For general background, see: Howells & Mccown (1977); Bowmaker *et al.* (2005); Lawrance (1986).



Experimental

Crystal data

 $[\text{Ag}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{C}_{18}\text{H}_{15}\text{P})]\text{CF}_3\text{SO}_3$ $M_r = 699.42$

Triclinic, $P\bar{1}$
 $a = 10.9832$ (2) Å
 $b = 11.7533$ (2) Å
 $c = 12.2642$ (3) Å
 $\alpha = 77.711$ (1)°
 $\beta = 76.183$ (1)°
 $\gamma = 73.440$ (1)°

$V = 1455.66$ (5) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.87$ mm⁻¹
 $T = 293$ K
 $0.4 \times 0.3 \times 0.2$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\text{min}} = 0.735$, $T_{\text{max}} = 0.832$

18629 measured reflections
 9515 independent reflections
 6777 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.133$
 $S = 1.00$
 9515 reflections

379 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.59$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.66$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Ag1—N2	2.2798 (18)	Ag1—P1	2.3469 (5)
Ag1—N1	2.292 (2)		
N2—Ag1—N1	73.53 (8)	N1—Ag1—P1	138.03 (6)
N2—Ag1—P1	147.77 (6)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}20-\text{H}16\cdots\text{O}2$	0.93	2.36	3.285 (6)	173

Data collection: SMART (Bruker, 2007); cell refinement: SAINT-Plus (Bruker, 2007); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the National Science Foundation of China (grant No. 20871085), the Committee of Education of Beijing Foundation of China (grant No. KM200610028006), the Project sponsored by SRF for ROCS and SEM, the subsidy of Beijing Personnel Bureau, the National Keystone Basic Research Program (973 Program under grant Nos. 2007CB310408, No. 2006CB302901), the State Key Laboratory of Functional Materials for Informatics and the Shanghai Institute of Microsystems and Information Technology, Chinese Academy of Sciences.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2241).

References

- Awaleh, M. O., Badia, A. & Brisse, F. (2005a). *Inorg. Chem.* **44**, 7833–7845.
- Awaleh, M. O., Badia, A. & Brisse, F. (2005b). *Cryst. Growth Des.* **5**, 1897–1906.
- Bowmaker, G. A., Effendy, Marfuah, S., Skelton, B. W. & White, A. H. (2005). *Inorg. Chim. Acta*, **358**, 4371–4388.
- Bruker (2007). *SMART, SAINT-Plus and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Di Nicola, C., Effendy, Marchetti, F., Pettinari, C., Skelton, B. W. & White, A. H. (2007). *Inorg. Chim. Acta*, **360**, 1433–1450.
- Effendy, Marchetti, F., Pettinari, C., Pettinari, R., Skelton, B. W. & White, A. H. (2007a). *Inorg. Chim. Acta*, **360**, 1451–1465.
- Effendy, Marchetti, F., Pettinari, C., Skelton, B. W. & White, A. H. (2007b). *Inorg. Chim. Acta*, **360**, 1424–1432.
- Howells, R. D. & Mccown, J. D. (1977). *Chem. Rev.* **77**, 19–92.
- Jin, Q. H., Hu, K. Y. S. L. L., Wang, R., Zuo, X., Zhang, C. L. & Lu, X. M. (2009). *Polyhedron*, doi: 10.1016/j.poly.2009.06.036.
- Jin, Q. H., Xin, X. L., Zhu, F. J. & Li, Y. (1999). *Z. Kristallogr. New Cryst. Struct.* **214**, 503–504.
- Lawrance, G. A. (1986). *Chem. Rev.* **86**, 17–33.
- Pettinari, C., Ngoune, J., Skelton, B. W. & White, A. H. (2007). *Inorg. Chem. Commun.* **10**, 329–331.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m1096-m1097 [doi:10.1107/S1600536809032097]

(1,10-Phenanthroline- κ^2N,N')(triphenylphosphine- κP)silver(I) trifluoromethanesulfonate

J.-Q. Wu, Q.-H. Jin, K.-Y. Hu and C.-L. Zhang

Comment

A recent report (Di Nicola *et al.*, 2007) describes complexes between silver nitrate, a tertiary phosphine ligand and oligodentate bases, *L* that are derivatives of 2,2'-bipyridyl, which resulted in adducts with general formula $\text{AgNO}_3:\text{PR}_3:L(1:1:1)$. The silver coordination environment in these complexes is dominated by the quasi-planar N_2AgP or O_2AgP coordination. We have likewise studied mixed-ligand Ag(I) complexes of *N*-heterocyclic and PPh_3 ligands, *viz* $[\text{AgBr}(\text{phen})(\text{PPh}_3)]$ and $[\text{AgX}(2\text{-Apy})(\text{PPh}_3)]_2$ ($X = \text{Br}, \text{Cl}, \text{NO}_3$; 2-Apy = 2-aminopyridine) (Jin *et al.*, 1999, Jin *et al.*, 2009) and have synthesized the title complex $[\text{Ag}(\text{phen})(\text{PPh}_3)](\text{OTf})$. Furthermore, we have studied the role of several weakly coordinating anions (nitrate, nitrite, acetate, perchlorate trifluoroacetate and trifluoromethanesulfonate) in silver complexes.

The molecular structure of the title complex is depicted in Fig.1. The coordination polyhedron of the silver atom adopts a distorted trigonal-planar geometry, formed by two nitrogen atoms of phen with $\text{Ag}-\text{N}$ distances of 2.3469 (5) Å and 2.2797 (19) Å, and by one phosphorus atom of the PPh_3 ligand with a $\text{Ag}-\text{P}$ distance of 2.292 (2) Å. The trifluoromethanesulfonate anion is present as a counter anion and, as expected, shows no direct coordination to the metal center, in contrast to the complex $[\text{AgBr}(\text{phen})(\text{PPh}_3)]$ where the silver atom is coordinated to two nitrogen atoms of phen ($\text{Ag}-\text{N}$ 2.376 (8) Å), one phosphorus atom of PPh_3 ($\text{Ag}-\text{P}$, 2.375 (3) Å) and in addition to one bromide anion (Jin *et al.*, 1999), adopting a distorted tetrahedron as coordination polyhedron.

The molecular structure of the title complex shows little differences in comparison with the structures of compounds $\text{AgX}:\text{PPh}_3:L$, where $X = \text{nitrate}$ (Di Nicola *et al.*, 2007), nitrite (Pettinari *et al.*, 2007), acetate (Effendy *et al.*, 2007a), perchlorate (Effendy *et al.*, 2007b) and trifluoroacetate (Awaleh *et al.*, 2005a). Considering the large steric hindrance and the weak coordination ability (Awaleh *et al.*, 2005b; Howells *et al.*, 1977; Lawrance *et al.*, 1986) of the trifluoromethanesulfonate anion, there is only one $\text{C}-\text{H}\cdots\text{O}$ hydrogen-bond between the phenanthroline ligand and the O atom of the anion with the distance $\text{O}\cdots\text{H}$ of 2.609 Å and the angle $\text{C}-\text{H}\cdots\text{O}$ of 173°.

In the title complex, the $\text{P}-\text{Ag}-\text{N}_1$, $\text{P}-\text{Ag}-\text{N}_2$ and $\text{N}_1-\text{Ag}-\text{N}_2$ angles are 147.77 (6)°, 138.03 (6)° and 73.54 (8)° with a sum of 359.54°, which confirms the trigonal-planar environment around the silver atom. In the silver nitrate complex, the $\text{P}-\text{Ag}-\text{N}$ (132.66 (9)°, 131.76 (8)°) (Di Nicola *et al.*, 2007) angles are similar. However, contributing to the role of the nitrate anion, the coordination environment of silver changes from distorted trigonal planar to tetrahedral. The $\text{P}-\text{Ag}-\text{N}$ angles in the other complexes are: 136.94 (5)°, 139.60 (5)°, 71.40 (6)° in the perchlorate (Effendy *et al.*, 2007b), 129.4 (1)°, 135.7 (1)°, 71.7 (2)° in trifluoroacetate (Awaleh *et al.*, 2005a), 116.52 (6)°, 126.12 (7)°, 70.5 (1)° in acetate (Effendy *et al.*, 2007a) and 126.72 (8)°, 127.18 (9)°, 70.77 (12)° in the nitrite (Pettinari *et al.*, 2007) anion.

Hence, we should consider two types of anions in the complexes $\text{AgX}:\text{PR}_3:L$, *viz* tetrahedral or distorted trigonal-planar anions and planar or quasi-planar anions (Awaleh *et al.*, 2005a; Awaleh *et al.*, 2005b; Bowmaker *et al.*, 2005). Nitrate, nitrite and acetate belong to the former type, whereas perchlorate, trifluoroacetate and trifluoromethanesulfonate can play a role in both of them because of large steric hindrance and the weak coordination ability.

Experimental

A mixture of AgOTf, Ph₃P and phen in the molar ratio of 1:1:1 in MeOH was stirred for 1 h at ambient temperature, then filtered. Subsequent slow evaporation of the filtrate resulted in the formation of colorless crystals of the title complex. Crystals suitable for single-crystal X-ray diffraction were selected directly from the sample as prepared. Analysis found (percentage): C 53.22, H 3.29, N 4.01; calculated: C 53.19, H 3.29, N 4.02.

Refinement

All hydrogen atoms were located in the calculated sites and included in the final refinement in the riding model approximation with displacement parameters derived from the parent atoms to which they were bonded ($U_{eq}(H) = 1.2U_{eq}(C)$).

Figures




Fig. 1. Perspective view of the molecule of the title complex; hydrogen atoms are omitted for clarity. Atoms are displayed as ellipsoids at the 35% probability level.

(1,10-Phenanthroline- κ^2N,N')(triphenylphosphine- κP)silver(I) trifluoromethanesulfonate

Crystal data

[Ag(C₁₂H₈N₂)(C₁₈H₁₅P)]CF₃SO₃

$M_r = 699.42$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.9832(2) \text{ \AA}$

$b = 11.7533(2) \text{ \AA}$

$c = 12.2642(3) \text{ \AA}$

$\alpha = 77.711(1)^\circ$

$\beta = 76.183(1)^\circ$

$\gamma = 73.440(1)^\circ$

$V = 1455.66(5) \text{ \AA}^3$

$Z = 2$

$F_{000} = 704$

$D_x = 1.596 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5045 reflections

$\theta = 2.3\text{--}32.9^\circ$

$\mu = 0.87 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colourless

$0.4 \times 0.3 \times 0.2 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
diffractometer

9515 independent reflections

Radiation source: fine-focus sealed tube	6777 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.021$
$T = 293$ K	$\theta_{\text{max}} = 32.6^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -15 \rightarrow 15$
$T_{\text{min}} = 0.735$, $T_{\text{max}} = 0.832$	$k = -17 \rightarrow 16$
18629 measured reflections	$l = -18 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.133$	$w = 1/[\sigma^2(F_o^2) + (0.075P)^2 + 0.38P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
9515 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
379 parameters	$\Delta\rho_{\text{max}} = 0.59 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.66 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.09032 (2)	0.931304 (15)	0.211327 (16)	0.05664 (9)
P1	0.18488 (6)	0.77307 (5)	0.10540 (5)	0.04241 (13)
C24	-0.0779 (3)	1.12309 (19)	0.36019 (18)	0.0491 (6)
C25	-0.3066 (4)	1.2209 (4)	0.3869 (4)	0.0981 (15)
H21	-0.3787	1.2746	0.4205	0.118*
S1	0.61400 (9)	0.77152 (8)	0.31032 (8)	0.0748 (2)
C1	0.0853 (2)	0.75012 (19)	0.01682 (19)	0.0455 (5)
C3	0.1327 (3)	0.6742 (3)	-0.0638 (3)	0.0631 (7)
H6	0.2207	0.6372	-0.0786	0.076*
C2	-0.0452 (3)	0.8068 (3)	0.0342 (3)	0.0644 (7)

supplementary materials

H10	-0.0782	0.8614	0.0852	0.077*
C4	0.3400 (2)	0.77960 (19)	0.01460 (18)	0.0433 (4)
C6	0.4324 (3)	0.6788 (2)	-0.0199 (2)	0.0566 (6)
H5	0.4166	0.6031	0.0064	0.068*
C7	0.4818 (3)	0.9018 (3)	-0.0982 (3)	0.0707 (8)
H2	0.4979	0.9772	-0.1261	0.085*
C5	0.3678 (3)	0.8918 (2)	-0.0246 (2)	0.0562 (6)
H1	0.3088	0.9603	-0.0008	0.067*
C10	0.1297 (3)	0.5596 (3)	0.2319 (2)	0.0602 (7)
H15	0.0575	0.5801	0.1981	0.072*
C8	0.2170 (2)	0.63074 (19)	0.20106 (18)	0.0438 (5)
C11	0.1495 (4)	0.4583 (3)	0.3128 (3)	0.0778 (10)
H14	0.0894	0.4118	0.3342	0.093*
C9	0.3237 (3)	0.5981 (2)	0.2526 (2)	0.0544 (6)
H11	0.3828	0.6456	0.2335	0.065*
C12	0.5478 (3)	0.6894 (3)	-0.0929 (3)	0.0652 (7)
H4	0.6082	0.6213	-0.1158	0.078*
C13	0.5725 (3)	0.8001 (3)	-0.1309 (3)	0.0677 (8)
H3	0.6504	0.8071	-0.1789	0.081*
C14	-0.1276 (3)	0.7833 (3)	-0.0234 (3)	0.0766 (9)
H9	-0.2156	0.8201	-0.0094	0.092*
C16	0.0513 (3)	0.6529 (3)	-0.1223 (3)	0.0715 (8)
H7	0.0846	0.6022	-0.1768	0.086*
C15	-0.0786 (3)	0.7058 (3)	-0.1007 (3)	0.0714 (8)
H8	-0.1336	0.6889	-0.1388	0.086*
C18	0.3430 (4)	0.4952 (3)	0.3324 (3)	0.0685 (8)
H12	0.4153	0.4733	0.3662	0.082*
C17	0.2555 (4)	0.4256 (3)	0.3616 (3)	0.0763 (9)
H13	0.2688	0.3561	0.4148	0.092*
O1	0.5839 (4)	0.7776 (4)	0.4276 (3)	0.1266 (12)
O2	0.5105 (4)	0.7955 (3)	0.2518 (4)	0.1519 (17)
O3	0.7108 (4)	0.8336 (4)	0.2534 (4)	0.1455 (16)
C19	0.6933 (5)	0.6175 (4)	0.2985 (6)	0.1103 (17)
F1	0.7991 (3)	0.5804 (4)	0.3393 (4)	0.1752 (17)
F2	0.7172 (6)	0.5954 (4)	0.1989 (4)	0.218 (3)
C23	0.0474 (3)	1.1023 (2)	0.38492 (18)	0.0513 (6)
C21	0.0633 (4)	1.1640 (3)	0.4660 (2)	0.0763 (11)
C22	-0.0480 (7)	1.2472 (3)	0.5189 (3)	0.1041 (18)
H19	-0.0385	1.2879	0.5723	0.125*
C20	0.2638 (4)	1.0021 (3)	0.3561 (3)	0.0752 (9)
H16	0.3327	0.9473	0.3203	0.090*
N1	0.1478 (2)	1.02361 (18)	0.33127 (18)	0.0522 (5)
N2	-0.09368 (19)	1.06137 (17)	0.28528 (15)	0.0489 (5)
C26	-0.1838 (4)	1.2062 (2)	0.4146 (2)	0.0757 (10)
C27	-0.2110 (3)	1.0794 (3)	0.2637 (3)	0.0708 (8)
H23	-0.2217	1.0367	0.2123	0.085*
C28	-0.1636 (6)	1.2669 (3)	0.4930 (3)	0.1029 (17)
H20	-0.2329	1.3223	0.5277	0.124*
C29	-0.3190 (4)	1.1595 (4)	0.3146 (4)	0.0990 (14)

H22	-0.3996	1.1693	0.2972	0.119*
C31	0.2848 (6)	1.0623 (5)	0.4377 (4)	0.1018 (17)
H17	0.3670	1.0475	0.4536	0.122*
C30	0.1856 (6)	1.1398 (5)	0.4908 (3)	0.1008 (17)
H18	0.1988	1.1777	0.5448	0.121*
F3	0.6223 (5)	0.5475 (3)	0.3633 (5)	0.213 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.06687 (15)	0.04557 (11)	0.05563 (12)	-0.00722 (8)	-0.00064 (9)	-0.02543 (8)
P1	0.0462 (3)	0.0368 (2)	0.0428 (3)	-0.0074 (2)	-0.0011 (2)	-0.0152 (2)
C24	0.0684 (16)	0.0344 (9)	0.0358 (9)	-0.0065 (9)	0.0011 (9)	-0.0071 (7)
C25	0.072 (2)	0.074 (2)	0.094 (3)	0.0230 (17)	0.021 (2)	0.0052 (19)
S1	0.0677 (5)	0.0771 (5)	0.0829 (5)	-0.0120 (4)	-0.0169 (4)	-0.0249 (4)
C1	0.0490 (13)	0.0417 (10)	0.0443 (11)	-0.0082 (8)	-0.0041 (9)	-0.0133 (8)
C3	0.0536 (15)	0.0689 (16)	0.0681 (16)	-0.0009 (12)	-0.0068 (12)	-0.0380 (13)
C2	0.0545 (16)	0.0662 (16)	0.0732 (18)	-0.0011 (12)	-0.0090 (13)	-0.0345 (13)
C4	0.0472 (12)	0.0395 (9)	0.0426 (10)	-0.0110 (8)	-0.0056 (9)	-0.0077 (8)
C6	0.0523 (15)	0.0434 (11)	0.0633 (15)	-0.0094 (10)	0.0063 (11)	-0.0083 (10)
C7	0.075 (2)	0.0628 (16)	0.077 (2)	-0.0352 (15)	-0.0136 (16)	0.0069 (14)
C5	0.0613 (16)	0.0413 (11)	0.0690 (16)	-0.0161 (10)	-0.0162 (13)	-0.0062 (10)
C10	0.0700 (18)	0.0622 (15)	0.0540 (14)	-0.0303 (13)	-0.0077 (12)	-0.0061 (11)
C8	0.0538 (13)	0.0407 (9)	0.0377 (9)	-0.0145 (9)	-0.0004 (9)	-0.0133 (8)
C11	0.107 (3)	0.0724 (19)	0.0640 (18)	-0.0527 (19)	-0.0103 (18)	0.0020 (15)
C9	0.0615 (16)	0.0525 (12)	0.0524 (13)	-0.0168 (11)	-0.0113 (11)	-0.0106 (10)
C12	0.0510 (15)	0.0647 (16)	0.0652 (16)	-0.0060 (12)	0.0046 (12)	-0.0074 (13)
C13	0.0497 (16)	0.082 (2)	0.0651 (17)	-0.0229 (14)	-0.0056 (13)	0.0055 (14)
C14	0.0515 (17)	0.086 (2)	0.098 (2)	0.0036 (14)	-0.0254 (16)	-0.0410 (18)
C16	0.071 (2)	0.0799 (19)	0.0713 (18)	-0.0068 (15)	-0.0163 (15)	-0.0411 (15)
C15	0.071 (2)	0.0742 (18)	0.077 (2)	-0.0096 (15)	-0.0304 (16)	-0.0228 (15)
C18	0.093 (2)	0.0581 (15)	0.0577 (15)	-0.0160 (15)	-0.0278 (15)	-0.0039 (12)
C17	0.118 (3)	0.0585 (16)	0.0554 (16)	-0.0331 (17)	-0.0203 (17)	0.0037 (12)
O1	0.105 (2)	0.169 (4)	0.097 (2)	-0.019 (2)	0.0043 (18)	-0.050 (2)
O2	0.161 (3)	0.090 (2)	0.238 (5)	0.013 (2)	-0.143 (4)	-0.038 (2)
O3	0.148 (3)	0.117 (3)	0.172 (4)	-0.072 (3)	0.034 (3)	-0.043 (2)
C19	0.094 (3)	0.070 (2)	0.161 (5)	-0.011 (2)	-0.041 (3)	0.002 (3)
F1	0.092 (2)	0.166 (3)	0.236 (4)	0.038 (2)	-0.057 (2)	-0.031 (3)
F2	0.336 (7)	0.123 (3)	0.175 (4)	0.056 (3)	-0.082 (4)	-0.096 (3)
C23	0.0848 (19)	0.0382 (10)	0.0355 (10)	-0.0281 (11)	-0.0072 (10)	-0.0027 (8)
C21	0.146 (3)	0.0638 (16)	0.0408 (12)	-0.068 (2)	-0.0155 (16)	0.0002 (11)
C22	0.216 (6)	0.0626 (19)	0.0432 (15)	-0.068 (3)	0.011 (2)	-0.0225 (13)
C20	0.071 (2)	0.080 (2)	0.080 (2)	-0.0389 (17)	-0.0279 (17)	0.0172 (16)
N1	0.0608 (13)	0.0474 (10)	0.0525 (11)	-0.0202 (9)	-0.0160 (9)	-0.0014 (8)
N2	0.0510 (12)	0.0468 (10)	0.0441 (10)	-0.0026 (8)	-0.0093 (8)	-0.0096 (8)
C26	0.104 (3)	0.0428 (12)	0.0515 (14)	0.0004 (13)	0.0185 (15)	-0.0092 (10)
C27	0.0586 (18)	0.0790 (19)	0.0655 (17)	-0.0017 (14)	-0.0208 (14)	-0.0024 (14)
C28	0.173 (5)	0.0529 (16)	0.061 (2)	-0.023 (2)	0.029 (3)	-0.0266 (14)

supplementary materials

C29	0.060 (2)	0.108 (3)	0.093 (3)	0.011 (2)	-0.0072 (19)	0.009 (2)
C31	0.120 (4)	0.131 (4)	0.089 (3)	-0.094 (3)	-0.059 (3)	0.040 (3)
C30	0.171 (5)	0.109 (3)	0.061 (2)	-0.103 (4)	-0.035 (3)	0.011 (2)
F3	0.209 (5)	0.103 (2)	0.335 (7)	-0.079 (3)	-0.087 (4)	0.042 (3)

Geometric parameters (Å, °)

Ag1—N2	2.2798 (18)	C11—H14	0.9300
Ag1—N1	2.292 (2)	C9—C18	1.384 (4)
Ag1—P1	2.3469 (5)	C9—H11	0.9300
P1—C4	1.812 (2)	C12—C13	1.365 (4)
P1—C1	1.819 (3)	C12—H4	0.9300
P1—C8	1.823 (2)	C13—H3	0.9300
C24—N2	1.353 (3)	C14—C15	1.364 (5)
C24—C26	1.418 (3)	C14—H9	0.9300
C24—C23	1.422 (4)	C16—C15	1.370 (5)
C25—C29	1.308 (7)	C16—H7	0.9300
C25—C26	1.423 (7)	C15—H8	0.9300
C25—H21	0.9300	C18—C17	1.371 (5)
S1—O1	1.409 (3)	C18—H12	0.9300
S1—O2	1.417 (3)	C17—H13	0.9300
S1—O3	1.421 (4)	C19—F2	1.253 (7)
S1—C19	1.791 (5)	C19—F1	1.298 (6)
C1—C3	1.381 (3)	C19—F3	1.307 (6)
C1—C2	1.384 (4)	C23—N1	1.355 (3)
C3—C16	1.375 (4)	C23—C21	1.416 (3)
C3—H6	0.9300	C21—C30	1.385 (7)
C2—C14	1.387 (5)	C21—C22	1.442 (7)
C2—H10	0.9300	C22—C28	1.326 (7)
C4—C6	1.391 (3)	C22—H19	0.9300
C4—C5	1.397 (3)	C20—N1	1.323 (4)
C6—C12	1.387 (4)	C20—C31	1.435 (6)
C6—H5	0.9300	C20—H16	0.9300
C7—C5	1.376 (4)	N2—C27	1.327 (4)
C7—C13	1.385 (5)	C26—C28	1.403 (6)
C7—H2	0.9300	C27—C29	1.394 (5)
C5—H1	0.9300	C27—H23	0.9300
C10—C11	1.380 (4)	C28—H20	0.9300
C10—C8	1.382 (4)	C29—H22	0.9300
C10—H15	0.9300	C31—C30	1.337 (7)
C8—C9	1.384 (4)	C31—H17	0.9300
C11—C17	1.356 (5)	C30—H18	0.9300
N2—Ag1—N1	73.53 (8)	C7—C13—H3	119.9
N2—Ag1—P1	147.77 (6)	C15—C14—C2	119.6 (3)
N1—Ag1—P1	138.03 (6)	C15—C14—H9	120.2
C4—P1—C1	106.47 (11)	C2—C14—H9	120.2
C4—P1—C8	104.71 (10)	C15—C16—C3	120.4 (3)
C1—P1—C8	103.84 (10)	C15—C16—H7	119.8
C4—P1—Ag1	115.16 (7)	C3—C16—H7	119.8

C1—P1—Ag1	115.70 (7)	C14—C15—C16	120.1 (3)
C8—P1—Ag1	109.82 (7)	C14—C15—H8	119.9
N2—C24—C26	121.2 (3)	C16—C15—H8	119.9
N2—C24—C23	118.8 (2)	C17—C18—C9	120.1 (3)
C26—C24—C23	120.0 (3)	C17—C18—H12	120.0
C29—C25—C26	120.5 (3)	C9—C18—H12	120.0
C29—C25—H21	119.7	C11—C17—C18	119.9 (3)
C26—C25—H21	119.7	C11—C17—H13	120.0
O1—S1—O2	118.2 (3)	C18—C17—H13	120.0
O1—S1—O3	111.0 (3)	F2—C19—F1	109.1 (6)
O2—S1—O3	113.3 (3)	F2—C19—F3	108.5 (6)
O1—S1—C19	105.7 (3)	F1—C19—F3	102.2 (5)
O2—S1—C19	103.3 (2)	F2—C19—S1	113.4 (4)
O3—S1—C19	103.5 (3)	F1—C19—S1	112.9 (4)
C3—C1—C2	118.2 (3)	F3—C19—S1	110.1 (4)
C3—C1—P1	123.0 (2)	N1—C23—C21	121.9 (3)
C2—C1—P1	118.73 (18)	N1—C23—C24	119.3 (2)
C16—C3—C1	120.6 (3)	C21—C23—C24	118.8 (3)
C16—C3—H6	119.7	C30—C21—C23	117.7 (4)
C1—C3—H6	119.7	C30—C21—C22	123.6 (4)
C1—C2—C14	121.0 (3)	C23—C21—C22	118.7 (4)
C1—C2—H10	119.5	C28—C22—C21	121.5 (3)
C14—C2—H10	119.5	C28—C22—H19	119.2
C6—C4—C5	118.1 (2)	C21—C22—H19	119.2
C6—C4—P1	123.43 (18)	N1—C20—C31	120.8 (4)
C5—C4—P1	118.47 (19)	N1—C20—H16	119.6
C12—C6—C4	121.0 (2)	C31—C20—H16	119.6
C12—C6—H5	119.5	C20—N1—C23	119.3 (3)
C4—C6—H5	119.5	C20—N1—Ag1	126.7 (2)
C5—C7—C13	120.2 (3)	C23—N1—Ag1	113.81 (17)
C5—C7—H2	119.9	C27—N2—C24	118.7 (2)
C13—C7—H2	119.9	C27—N2—Ag1	126.7 (2)
C7—C5—C4	120.5 (3)	C24—N2—Ag1	114.57 (16)
C7—C5—H1	119.7	C28—C26—C24	119.4 (4)
C4—C5—H1	119.7	C28—C26—C25	123.8 (4)
C11—C10—C8	120.1 (3)	C24—C26—C25	116.9 (3)
C11—C10—H15	120.0	N2—C27—C29	122.8 (4)
C8—C10—H15	120.0	N2—C27—H23	118.6
C10—C8—C9	118.8 (2)	C29—C27—H23	118.6
C10—C8—P1	121.0 (2)	C22—C28—C26	121.6 (4)
C9—C8—P1	119.95 (18)	C22—C28—H20	119.2
C17—C11—C10	120.8 (3)	C26—C28—H20	119.2
C17—C11—H14	119.6	C25—C29—C27	119.9 (4)
C10—C11—H14	119.6	C25—C29—H22	120.1
C8—C9—C18	120.3 (3)	C27—C29—H22	120.1
C8—C9—H11	119.9	C30—C31—C20	119.8 (4)
C18—C9—H11	119.9	C30—C31—H17	120.1
C13—C12—C6	119.9 (3)	C20—C31—H17	120.1
C13—C12—H4	120.1	C31—C30—C21	120.4 (4)

supplementary materials

C6—C12—H4	120.1	C31—C30—H18	119.8
C12—C13—C7	120.2 (3)	C21—C30—H18	119.8
C12—C13—H3	119.9		
N2—Ag1—P1—C4	143.08 (12)	O1—S1—C19—F1	-60.6 (5)
N1—Ag1—P1—C4	-51.26 (12)	O2—S1—C19—F1	174.5 (5)
N2—Ag1—P1—C1	18.06 (13)	O3—S1—C19—F1	56.2 (5)
N1—Ag1—P1—C1	-176.27 (11)	O1—S1—C19—F3	52.9 (5)
N2—Ag1—P1—C8	-99.05 (13)	O2—S1—C19—F3	-71.9 (5)
N1—Ag1—P1—C8	66.61 (12)	O3—S1—C19—F3	169.7 (5)
C4—P1—C1—C3	40.1 (3)	N2—C24—C23—N1	1.4 (3)
C8—P1—C1—C3	-70.1 (3)	C26—C24—C23—N1	-179.3 (2)
Ag1—P1—C1—C3	169.5 (2)	N2—C24—C23—C21	-178.0 (2)
C4—P1—C1—C2	-142.9 (2)	C26—C24—C23—C21	1.3 (3)
C8—P1—C1—C2	106.9 (2)	N1—C23—C21—C30	-1.1 (4)
Ag1—P1—C1—C2	-13.6 (2)	C24—C23—C21—C30	178.3 (2)
C2—C1—C3—C16	-2.1 (5)	N1—C23—C21—C22	179.5 (2)
P1—C1—C3—C16	174.9 (3)	C24—C23—C21—C22	-1.1 (3)
C3—C1—C2—C14	3.3 (5)	C30—C21—C22—C28	-179.5 (3)
P1—C1—C2—C14	-173.8 (3)	C23—C21—C22—C28	-0.1 (5)
C1—P1—C4—C6	-75.7 (2)	C31—C20—N1—C23	-0.9 (4)
C8—P1—C4—C6	33.9 (2)	C31—C20—N1—Ag1	-176.6 (2)
Ag1—P1—C4—C6	154.6 (2)	C21—C23—N1—C20	1.0 (3)
C1—P1—C4—C5	103.2 (2)	C24—C23—N1—C20	-178.4 (2)
C8—P1—C4—C5	-147.2 (2)	C21—C23—N1—Ag1	177.19 (17)
Ag1—P1—C4—C5	-26.5 (2)	C24—C23—N1—Ag1	-2.2 (3)
C5—C4—C6—C12	-1.2 (4)	N2—Ag1—N1—C20	177.5 (2)
P1—C4—C6—C12	177.6 (2)	P1—Ag1—N1—C20	5.5 (3)
C13—C7—C5—C4	-2.4 (5)	N2—Ag1—N1—C23	1.63 (15)
C6—C4—C5—C7	2.1 (4)	P1—Ag1—N1—C23	-170.45 (11)
P1—C4—C5—C7	-176.8 (2)	C26—C24—N2—C27	-0.4 (3)
C11—C10—C8—C9	0.1 (4)	C23—C24—N2—C27	178.9 (2)
C11—C10—C8—P1	-173.6 (3)	C26—C24—N2—Ag1	-179.12 (18)
C4—P1—C8—C10	-140.7 (2)	C23—C24—N2—Ag1	0.2 (3)
C1—P1—C8—C10	-29.2 (2)	N1—Ag1—N2—C27	-179.6 (2)
Ag1—P1—C8—C10	95.1 (2)	P1—Ag1—N2—C27	-9.5 (3)
C4—P1—C8—C9	45.7 (2)	N1—Ag1—N2—C24	-0.94 (15)
C1—P1—C8—C9	157.17 (19)	P1—Ag1—N2—C24	169.11 (11)
Ag1—P1—C8—C9	-78.53 (19)	N2—C24—C26—C28	179.0 (3)
C8—C10—C11—C17	-1.3 (5)	C23—C24—C26—C28	-0.3 (4)
C10—C8—C9—C18	0.8 (4)	N2—C24—C26—C25	0.2 (4)
P1—C8—C9—C18	174.6 (2)	C23—C24—C26—C25	-179.1 (2)
C4—C6—C12—C13	0.7 (5)	C29—C25—C26—C28	-178.6 (4)
C6—C12—C13—C7	-1.0 (5)	C29—C25—C26—C24	0.1 (5)
C5—C7—C13—C12	1.9 (5)	C24—N2—C27—C29	0.2 (4)
C1—C2—C14—C15	-1.9 (6)	Ag1—N2—C27—C29	178.8 (3)
C1—C3—C16—C15	-0.6 (6)	C21—C22—C28—C26	1.2 (6)
C2—C14—C15—C16	-0.9 (6)	C24—C26—C28—C22	-1.0 (5)
C3—C16—C15—C14	2.1 (6)	C25—C26—C28—C22	177.7 (3)
C8—C9—C18—C17	-0.6 (5)	C26—C25—C29—C27	-0.3 (6)

C10—C11—C17—C18	1.5 (6)	N2—C27—C29—C25	0.1 (6)
C9—C18—C17—C11	-0.5 (5)	N1—C20—C31—C30	1.1 (5)
O1—S1—C19—F2	174.6 (5)	C20—C31—C30—C21	-1.2 (5)
O2—S1—C19—F2	49.8 (6)	C23—C21—C30—C31	1.2 (5)
O3—S1—C19—F2	-68.6 (6)	C22—C21—C30—C31	-179.4 (3)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C20—H16...O2	0.93	2.36	3.285 (6)	173

Fig. 1

