

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

6-Methyl-7,7,9-tripropargyl-7H-1,2,4-triazolo[4,3-*b*][1,2,4]triazepin-8(9H)-oneRedwan Mohamed Zemama,^a Ibtissam Amari,^a Rachid Bouhfid,^b El Mokhtar Essassi^a and Seik Weng Ng^{c*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bInstitute of Nanomaterials and Nanotechnology, Avenue de l'Armée Royale, Madinat El Irfane, 10100 Rabat, Morocco, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 7 August 2009; accepted 8 August 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.129; data-to-parameter ratio = 16.7.

The title compound, $\text{C}_{15}\text{H}_{13}\text{N}_5\text{O}$, features a triazolyl ring fused with a seven-membered triazepinyl ring; the latter ring adopts a boat conformation (with the propargyl-bearing C atom as the prow and the fused-ring C/N atoms as the stern).

Related literature

Triazepines are a class of compounds used in the treatment of neuronal disorders. They are also the reactants for the synthesis of other heterocyclic compounds; see, for example: Essassi *et al.* (1977); Richter & Sheefelot (1991).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{13}\text{N}_5\text{O}$
 $M_r = 279.30$
 Triclinic, $P\bar{1}$
 $a = 7.6710$ (2) Å
 $b = 8.2415$ (2) Å
 $c = 12.9619$ (3) Å
 $\alpha = 108.510$ (1)°
 $\beta = 90.659$ (1)°
 $\gamma = 114.726$ (1)°
 $V = 695.85$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.3 \times 0.3 \times 0.3$ mm

Data collection

Bruker APEX2 diffractometer
 Absorption correction: none
 16879 measured reflections
 3187 independent reflections
 2577 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.129$
 $S = 1.06$
 3187 reflections
 191 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2009).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2584).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2005). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Essassi, E. M., Laverigne, J. P. & Viallefont, P. (1977). *Tetrahedron*, **33**, 2807–2812.
 Richter, P. & Sheefelot, U. (1991). *Pharmazie*, **46**, 701–705.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). *pubCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o2148 [doi:10.1107/S1600536809031341]

6-Methyl-7,7,9-triisopropargyl-7*H*-1,2,4-triazolo[4,3-*b*][1,2,4]triazepin-8(9*H*)-one

R. M. Zemama, I. Amari, R. Bouhfid, E. M. Essassi and S. W. Ng

Experimental

To a solution of 6-methyl-7*H*-[1,2,4]triazolo[4,3-*b*][1,2,4]triazepin-8(9*H*)-one (1 g, 6 mmol) in *N,N*-dimethylformamide (20 ml), potassium carbonate (1.26 g, 9 mmol), propargyl bromide (0.7 ml, 9 mmol) and a catalytic amount of tetrabutylammonium bromide were added. The mixture was stirred for 12 h. After the completion of the reaction (as monitored by TLC), the solid material was removed by filtration and the solvent evaporated under vacuum. Dichloromethane (20 ml) was added and the solution filtered. The solvent was removed and the product purified by column chromatography (30% ethyl acetate/hexane) to afford colorless crystals in 15% yield; m.p. 463 K. The formulation was established by proton and carbon-13 NMR spectroscopy in DMSO-*d*₆.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for the others.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of C₁₅H₁₃N₅O at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

6-Methyl-7,7,9-triisopropargyl-7*H*-1,2,4- triazolo[4,3-*b*][1,2,4]triazepin-8(9*H*)-one

Crystal data

C₁₅H₁₃N₅O

$M_r = 279.30$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.6710$ (2) Å

$b = 8.2415$ (2) Å

$c = 12.9619$ (3) Å

$\alpha = 108.510$ (1)°

$\beta = 90.659$ (1)°

$Z = 2$

$F_{000} = 292$

$D_x = 1.333$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6818 reflections

$\theta = 2.8\text{--}32.3^\circ$

$\mu = 0.09$ mm⁻¹

$T = 293$ K

Block, colorless

supplementary materials

$\gamma = 114.726 (1)^\circ$
 $V = 695.85 (3) \text{ \AA}^3$

$0.3 \times 0.3 \times 0.3 \text{ mm}$

Data collection

Bruker APEX2 diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 293 \text{ K}$
 φ and ω scans
Absorption correction: None
16879 measured reflections
3187 independent reflections

2577 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$
 $\theta_{\text{max}} = 27.5^\circ$
 $\theta_{\text{min}} = 1.7^\circ$
 $h = -9 \rightarrow 9$
 $k = -10 \rightarrow 10$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.129$
 $S = 1.06$
3187 reflections
191 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0758P)^2 + 0.0861P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.30213 (18)	0.47985 (14)	0.37303 (8)	0.0488 (3)
N1	0.07241 (16)	-0.03916 (15)	0.11746 (9)	0.0346 (3)
N2	0.22767 (16)	0.05768 (15)	0.07150 (8)	0.0323 (3)
N3	0.4030 (2)	0.1070 (2)	-0.05735 (10)	0.0462 (3)
N4	0.46123 (19)	0.28152 (18)	0.02864 (10)	0.0446 (3)
N5	0.35151 (17)	0.38407 (15)	0.19840 (9)	0.0372 (3)
C1	0.31447 (19)	0.35728 (17)	0.29631 (11)	0.0330 (3)
C2	0.28282 (18)	0.16329 (17)	0.30330 (10)	0.0289 (3)
C3	0.09422 (18)	0.01262 (17)	0.22312 (10)	0.0303 (3)
C4	-0.0871 (2)	-0.0827 (2)	0.26582 (13)	0.0475 (4)
H4A	-0.1893	-0.1748	0.2049	0.071*
H4B	-0.1255	0.0114	0.3090	0.071*
H4C	-0.0628	-0.1461	0.3109	0.071*
C5	0.45567 (18)	0.11327 (18)	0.27425 (11)	0.0317 (3)
H5A	0.4502	0.0191	0.3055	0.038*
H5B	0.4400	0.0560	0.1947	0.038*

C6	0.6465 (2)	0.2779 (2)	0.31450 (12)	0.0412 (3)
C7	0.8004 (3)	0.4075 (3)	0.34405 (18)	0.0702 (5)
H7	0.9233	0.5109	0.3676	0.084*
C8	0.2658 (2)	0.18290 (19)	0.42552 (11)	0.0372 (3)
H8A	0.3796	0.2944	0.4728	0.045*
H8B	0.1525	0.2037	0.4431	0.045*
C9	0.2492 (2)	0.0156 (2)	0.44969 (11)	0.0398 (3)
C10	0.2456 (2)	-0.1166 (2)	0.46867 (14)	0.0512 (4)
H10	0.2427	-0.2208	0.4836	0.061*
C11	0.2649 (2)	-0.0213 (2)	-0.03028 (11)	0.0401 (3)
H11	0.1996	-0.1499	-0.0742	0.048*
C12	0.3531 (2)	0.24704 (19)	0.10322 (11)	0.0343 (3)
C13	0.3765 (2)	0.56697 (19)	0.18940 (13)	0.0451 (4)
H13A	0.4427	0.6700	0.2593	0.054*
H13B	0.4564	0.5933	0.1338	0.054*
C14	0.1885 (3)	0.5585 (2)	0.15971 (13)	0.0482 (4)
C15	0.0358 (3)	0.5466 (3)	0.13320 (17)	0.0655 (5)
H15	-0.0853	0.5372	0.1122	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0733 (8)	0.0385 (5)	0.0407 (6)	0.0331 (5)	0.0150 (5)	0.0102 (4)
N1	0.0306 (6)	0.0390 (6)	0.0356 (6)	0.0150 (5)	0.0058 (4)	0.0157 (5)
N2	0.0347 (6)	0.0390 (6)	0.0276 (5)	0.0190 (5)	0.0069 (4)	0.0138 (4)
N3	0.0575 (8)	0.0644 (8)	0.0371 (6)	0.0405 (7)	0.0202 (6)	0.0246 (6)
N4	0.0528 (8)	0.0531 (7)	0.0458 (7)	0.0315 (6)	0.0235 (6)	0.0285 (6)
N5	0.0478 (7)	0.0324 (5)	0.0395 (6)	0.0214 (5)	0.0155 (5)	0.0180 (5)
C1	0.0355 (7)	0.0307 (6)	0.0346 (7)	0.0165 (5)	0.0079 (5)	0.0114 (5)
C2	0.0318 (6)	0.0302 (6)	0.0278 (6)	0.0156 (5)	0.0082 (5)	0.0116 (5)
C3	0.0293 (6)	0.0325 (6)	0.0341 (7)	0.0158 (5)	0.0069 (5)	0.0152 (5)
C4	0.0324 (7)	0.0619 (9)	0.0511 (9)	0.0162 (7)	0.0116 (6)	0.0301 (8)
C5	0.0315 (6)	0.0366 (6)	0.0311 (6)	0.0180 (5)	0.0071 (5)	0.0132 (5)
C6	0.0358 (7)	0.0486 (8)	0.0410 (8)	0.0188 (6)	0.0084 (6)	0.0182 (6)
C7	0.0408 (10)	0.0661 (11)	0.0781 (13)	0.0050 (8)	0.0039 (9)	0.0197 (10)
C8	0.0454 (8)	0.0416 (7)	0.0290 (6)	0.0224 (6)	0.0112 (6)	0.0138 (5)
C9	0.0400 (7)	0.0508 (8)	0.0310 (7)	0.0195 (6)	0.0083 (6)	0.0183 (6)
C10	0.0515 (9)	0.0566 (9)	0.0538 (9)	0.0225 (8)	0.0074 (7)	0.0319 (8)
C11	0.0478 (8)	0.0517 (8)	0.0295 (7)	0.0313 (7)	0.0066 (6)	0.0129 (6)
C12	0.0390 (7)	0.0390 (7)	0.0361 (7)	0.0224 (6)	0.0122 (5)	0.0204 (6)
C13	0.0548 (9)	0.0323 (7)	0.0539 (9)	0.0186 (6)	0.0167 (7)	0.0236 (6)
C14	0.0677 (11)	0.0368 (7)	0.0506 (9)	0.0289 (7)	0.0161 (8)	0.0208 (7)
C15	0.0762 (13)	0.0636 (11)	0.0728 (12)	0.0435 (10)	0.0109 (10)	0.0274 (10)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.2094 (15)	C4—H4C	0.9600
N1—C3	1.2845 (17)	C5—C6	1.4583 (19)
N1—N2	1.3905 (15)	C5—H5A	0.9700

supplementary materials

N2—C12	1.3630 (17)	C5—H5B	0.9700
N2—C11	1.3653 (17)	C6—C7	1.165 (2)
N3—C11	1.293 (2)	C7—H7	0.9300
N3—N4	1.3940 (18)	C8—C9	1.4643 (19)
N4—C12	1.3035 (17)	C8—H8A	0.9700
N5—C1	1.3690 (17)	C8—H8B	0.9700
N5—C12	1.3874 (17)	C9—C10	1.180 (2)
N5—C13	1.4808 (16)	C10—H10	0.9300
C1—C2	1.5463 (16)	C11—H11	0.9300
C2—C3	1.5357 (17)	C13—C14	1.455 (2)
C2—C8	1.5532 (17)	C13—H13A	0.9700
C2—C5	1.5620 (17)	C13—H13B	0.9700
C3—C4	1.4957 (19)	C14—C15	1.172 (3)
C4—H4A	0.9600	C15—H15	0.9300
C4—H4B	0.9600		
C3—N1—N2	117.64 (11)	C2—C5—H5A	108.8
C12—N2—C11	104.18 (11)	C6—C5—H5B	108.8
C12—N2—N1	130.86 (11)	C2—C5—H5B	108.8
C11—N2—N1	124.18 (11)	H5A—C5—H5B	107.7
C11—N3—N4	107.51 (11)	C7—C6—C5	178.18 (18)
C12—N4—N3	106.47 (12)	C6—C7—H7	180.0
C1—N5—C12	124.97 (10)	C9—C8—C2	113.42 (11)
C1—N5—C13	118.17 (11)	C9—C8—H8A	108.9
C12—N5—C13	116.66 (11)	C2—C8—H8A	108.9
O1—C1—N5	120.67 (11)	C9—C8—H8B	108.9
O1—C1—C2	121.75 (12)	C2—C8—H8B	108.9
N5—C1—C2	117.55 (10)	H8A—C8—H8B	107.7
C3—C2—C1	106.03 (10)	C10—C9—C8	176.60 (16)
C3—C2—C8	112.84 (10)	C9—C10—H10	180.0
C1—C2—C8	105.71 (10)	N3—C11—N2	110.81 (13)
C3—C2—C5	110.77 (10)	N3—C11—H11	124.6
C1—C2—C5	113.58 (10)	N2—C11—H11	124.6
C8—C2—C5	107.91 (10)	N4—C12—N2	111.01 (12)
N1—C3—C4	113.98 (12)	N4—C12—N5	125.45 (12)
N1—C3—C2	125.47 (11)	N2—C12—N5	123.34 (11)
C4—C3—C2	120.51 (11)	C14—C13—N5	110.59 (12)
C3—C4—H4A	109.5	C14—C13—H13A	109.5
C3—C4—H4B	109.5	N5—C13—H13A	109.5
H4A—C4—H4B	109.5	C14—C13—H13B	109.5
C3—C4—H4C	109.5	N5—C13—H13B	109.5
H4A—C4—H4C	109.5	H13A—C13—H13B	108.1
H4B—C4—H4C	109.5	C15—C14—C13	178.00 (18)
C6—C5—C2	113.64 (11)	C14—C15—H15	180.0
C6—C5—H5A	108.8		
C3—N1—N2—C12	-40.63 (18)	C1—C2—C5—C6	37.77 (15)
C3—N1—N2—C11	151.13 (13)	C8—C2—C5—C6	-79.06 (13)
C11—N3—N4—C12	0.01 (16)	C3—C2—C8—C9	68.72 (15)
C12—N5—C1—O1	-174.64 (13)	C1—C2—C8—C9	-175.83 (11)

C13—N5—C1—O1	0.2 (2)	C5—C2—C8—C9	-53.99 (15)
C12—N5—C1—C2	3.4 (2)	C2—C8—C9—C10	72 (3)
C13—N5—C1—C2	178.27 (11)	N4—N3—C11—N2	-0.81 (16)
O1—C1—C2—C3	112.39 (14)	C12—N2—C11—N3	1.25 (15)
N5—C1—C2—C3	-65.67 (14)	N1—N2—C11—N3	172.10 (11)
O1—C1—C2—C8	-7.63 (17)	N3—N4—C12—N2	0.80 (15)
N5—C1—C2—C8	174.31 (12)	N3—N4—C12—N5	-174.12 (12)
O1—C1—C2—C5	-125.74 (14)	C11—N2—C12—N4	-1.25 (15)
N5—C1—C2—C5	56.20 (15)	N1—N2—C12—N4	-171.23 (12)
N2—N1—C3—C4	173.10 (11)	C11—N2—C12—N5	173.80 (12)
N2—N1—C3—C2	-4.67 (18)	N1—N2—C12—N5	3.8 (2)
C1—C2—C3—N1	68.84 (15)	C1—N5—C12—N4	-148.84 (14)
C8—C2—C3—N1	-175.91 (12)	C13—N5—C12—N4	36.3 (2)
C5—C2—C3—N1	-54.80 (16)	C1—N5—C12—N2	36.8 (2)
C1—C2—C3—C4	-108.79 (13)	C13—N5—C12—N2	-138.05 (13)
C8—C2—C3—C4	6.46 (16)	C1—N5—C13—C14	-84.87 (17)
C5—C2—C3—C4	127.57 (13)	C12—N5—C13—C14	90.39 (15)
C3—C2—C5—C6	156.96 (11)		

Fig. 1

