

## 3-Chloro-5-methoxy-2,6-dinitropyridine

Jian-feng Guo and Jian-long Wang\*

School of Chemical Engineering and Environment, North University of China,  
Taiyuan, People's Republic of China  
Correspondence e-mail: wangjianlong@nuc.edu.cn


Received 8 August 2009; accepted 10 August 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  
 $R$  factor = 0.048;  $wR$  factor = 0.135; data-to-parameter ratio = 14.9.

In the crystal structure of the title compound, C<sub>6</sub>H<sub>4</sub>ClN<sub>3</sub>O<sub>5</sub>, the two nitro groups are twisted with respect to the pyridine ring, making dihedral angles of 33.12 (13) and 63.66 (14)°.

### Related literature

For the synthesis, see: Bissell & Swansiger (1987); Chen *et al.* (2008).



### Experimental

#### Crystal data

C<sub>6</sub>H<sub>4</sub>ClN<sub>3</sub>O<sub>5</sub>

$M_r = 233.57$

#### Data collection

Rigaku R-AXIS RAPID IP  
diffractometer  
Absorption correction: multi-scan  
(ABSCOR; Higashi, 1995)  
 $T_{\min} = 0.808$ ,  $T_{\max} = 0.887$

5866 measured reflections  
2062 independent reflections  
1275 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.050$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.135$   
 $S = 0.99$   
2062 reflections

138 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.26$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.30$  e Å<sup>-3</sup>

Data collection: RAPID-AUTO (Rigaku, 1998); cell refinement: RAPID-AUTO; data reduction: CrystalStructure (Rigaku/MSC, 2002); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This work was supported by China North Industries Group Corporation (project No. ZC060302-15).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2586).

### References

- Bissell, E. R. & Swansiger, R. W. (1987). *J. Heterocycl. Chem.* **24**, 59–62.
- Chen, J., Li, Q.-L. & Wang, J.-L. (2008). *Chin. J. Org. Chem.* **28**, 123–126.
- Higashi, T. (1995). ABSCOR. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). PROCESS-AUTO. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2000). RAPID-AUTO. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). CrystalStructure. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2009). E65, o2154 [doi:10.1107/S1600536809031407]

## 3-Chloro-5-methoxy-2,6-dinitropyridine

Jian-feng Guo and Jian-long Wang

### S1. Comment

Pyridine derivatives are important intermediates used to synthesize pesticide, medicine and play important roles in fine chemical field. 3-Chloro-5-methoxyl-2,6-dinitro-pyridine was synthesized from 3,5-dichloropyridine N-oxide by substitution and nitration (Bissell *et al.*, 1987), and the process was improved by Chen *et al.* (2008). The crystal structure of the title compound is presented here.


The molecular structure of the title compound is shown in Fig. 1. While the methoxyl group, except H atoms, is co-planar with the pyridine ring, the two nitro groups are twisted with respect to the pyridine ring with dihedral angles of 33.12 (13) and 63.66 (14) $^{\circ}$ , respectively. Neither hydrogen bonding nor  $\pi$ - $\pi$  stacking is observed in the crystal structure.

### S2. Experimental

The title compound was prepared according to a literature method (Chen *et al.*, 2008). Crystals suitable for X-ray analysis were obtained by slow evaporation of 1,2-dichloroethane.

### S3. Refinement

H atoms were positioned geometrically and refined using a ride model with C—H = 0.93 Å for aromatic H and 0.96 Å for methyl H atoms,  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{C})$  for aromatic H atom.

**Figure 1**

The molecular structure of title compound, with 30% probability displacement ellipsoids (arbitrary spheres for H atoms).

### 3-Chloro-5-methoxy-2,6-dinitropyridine

#### Crystal data



$M_r = 233.57$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 6.6490 (13) \text{ \AA}$

$b = 10.842 (2) \text{ \AA}$

$c = 12.715 (3) \text{ \AA}$

$\beta = 95.55 (3)^\circ$

$V = 912.3 (3) \text{ \AA}^3$

$Z = 4$

$F(000) = 472$

$D_x = 1.701 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5866 reflections

$\theta = 2.5\text{--}27.5^\circ$

$\mu = 0.43 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colorless

$0.50 \times 0.40 \times 0.28 \text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID IP  
diffractometer

Radiation source: fine-focus sealed tube  
Graphite monochromator

Detector resolution: 10.00 pixels  $\text{mm}^{-1}$   
 $\omega$  scans

Absorption correction: multi-scan  
(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.808$ ,  $T_{\max} = 0.887$

5866 measured reflections

2062 independent reflections

1275 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.050$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.5^\circ$

$h = -8 \rightarrow 8$

$k = -13 \rightarrow 13$

$l = -16 \rightarrow 16$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.048$$

$$wR(F^2) = 0.135$$

$$S = 0.99$$

2062 reflections

138 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.08P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$$

Extinction correction: *SHELXTL* (Version 4.2;  
Sheldrick, 2008),

$$Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

Extinction coefficient: 0.157 (11)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.17055 (10)	0.52802 (8)	0.42365 (6)	0.0756 (3)
O1	0.8010 (3)	0.77739 (16)	0.37270 (12)	0.0536 (5)
O2	0.2241 (4)	0.3362 (2)	0.27062 (18)	0.0939 (8)
O3	0.2590 (3)	0.4071 (2)	0.11455 (16)	0.0716 (6)
O4	0.8115 (4)	0.7839 (2)	0.14363 (19)	0.0949 (8)
O5	0.9482 (4)	0.6061 (2)	0.1524 (2)	0.1051 (9)
N1	0.5510 (3)	0.55114 (17)	0.20338 (15)	0.0439 (5)
N2	0.2845 (3)	0.4126 (2)	0.2107 (2)	0.0590 (6)
N3	0.8230 (3)	0.6789 (2)	0.17477 (16)	0.0556 (6)
C1	0.4068 (3)	0.5141 (2)	0.25985 (18)	0.0440 (5)
C2	0.3749 (3)	0.5646 (2)	0.35677 (18)	0.0453 (6)
C3	0.5061 (3)	0.6544 (2)	0.39881 (17)	0.0443 (5)
H3	0.4898	0.6885	0.4645	0.053*
C4	0.6614 (3)	0.6929 (2)	0.34246 (16)	0.0407 (5)
C5	0.6705 (3)	0.6373 (2)	0.24397 (16)	0.0412 (5)
C6	0.7921 (5)	0.8349 (3)	0.47424 (19)	0.0657 (8)
H6A	0.6641	0.8754	0.4761	0.099*
H6B	0.8989	0.8944	0.4857	0.099*
H6C	0.8073	0.7732	0.5286	0.099*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0570 (4)	0.0974 (6)	0.0777 (5)	-0.0107 (4)	0.0343 (4)	0.0139 (4)

O1	0.0575 (10)	0.0580 (10)	0.0474 (9)	-0.0144 (8)	0.0153 (8)	-0.0112 (8)
O2	0.1109 (18)	0.0831 (15)	0.0882 (15)	-0.0542 (15)	0.0115 (13)	0.0088 (12)
O3	0.0704 (12)	0.0772 (13)	0.0661 (13)	-0.0183 (10)	0.0006 (10)	-0.0065 (10)
O4	0.1047 (18)	0.0908 (18)	0.0946 (16)	-0.0134 (14)	0.0377 (14)	0.0320 (13)
O5	0.0869 (16)	0.1075 (19)	0.133 (2)	0.0047 (15)	0.0717 (15)	-0.0076 (16)
N1	0.0418 (10)	0.0451 (10)	0.0457 (10)	-0.0002 (9)	0.0083 (8)	-0.0001 (8)
N2	0.0494 (12)	0.0609 (14)	0.0666 (15)	-0.0129 (11)	0.0060 (10)	0.0026 (11)
N3	0.0526 (12)	0.0688 (15)	0.0481 (11)	-0.0127 (11)	0.0182 (9)	-0.0061 (11)
C1	0.0400 (11)	0.0430 (12)	0.0492 (12)	-0.0024 (10)	0.0053 (10)	0.0060 (10)
C2	0.0388 (11)	0.0492 (13)	0.0493 (13)	0.0041 (10)	0.0121 (9)	0.0141 (10)
C3	0.0468 (12)	0.0476 (13)	0.0405 (12)	0.0064 (11)	0.0150 (9)	0.0035 (10)
C4	0.0415 (11)	0.0395 (11)	0.0422 (12)	0.0038 (10)	0.0096 (9)	0.0035 (9)
C5	0.0364 (10)	0.0460 (12)	0.0423 (11)	0.0004 (10)	0.0097 (9)	0.0037 (9)
C6	0.0845 (19)	0.0678 (17)	0.0463 (14)	-0.0182 (15)	0.0138 (13)	-0.0125 (12)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C11—C2	1.717 (2)	N3—C5	1.476 (3)
O1—C4	1.334 (3)	C1—C2	1.384 (3)
O1—C6	1.440 (3)	C2—C3	1.380 (3)
O2—N2	1.219 (3)	C3—C4	1.378 (3)
O3—N2	1.219 (3)	C3—H3	0.9300
O4—N3	1.206 (3)	C4—C5	1.396 (3)
O5—N3	1.201 (3)	C6—H6A	0.9600
N1—C5	1.300 (3)	C6—H6B	0.9600
N1—C1	1.315 (3)	C6—H6C	0.9600
N2—C1	1.472 (3)		
C4—O1—C6	117.90 (18)	C4—C3—H3	120.3
C5—N1—C1	116.89 (19)	C2—C3—H3	120.3
O3—N2—O2	124.9 (2)	O1—C4—C3	126.4 (2)
O3—N2—C1	118.7 (2)	O1—C4—C5	117.78 (18)
O2—N2—C1	116.4 (2)	C3—C4—C5	115.8 (2)
O5—N3—O4	124.5 (2)	N1—C5—C4	126.05 (19)
O5—N3—C5	118.2 (2)	N1—C5—N3	114.31 (18)
O4—N3—C5	117.3 (2)	C4—C5—N3	119.6 (2)
N1—C1—C2	123.3 (2)	O1—C6—H6A	109.5
N1—C1—N2	113.5 (2)	O1—C6—H6B	109.5
C2—C1—N2	123.2 (2)	H6A—C6—H6B	109.5
C3—C2—C1	118.6 (2)	O1—C6—H6C	109.5
C3—C2—Cl1	118.16 (18)	H6A—C6—H6C	109.5
C1—C2—Cl1	123.13 (19)	H6B—C6—H6C	109.5
C4—C3—C2	119.3 (2)		
C5—N1—C1—C2	1.9 (3)	C6—O1—C4—C5	-180.0 (2)
C5—N1—C1—N2	-177.3 (2)	C2—C3—C4—O1	-179.3 (2)
O3—N2—C1—N1	-31.5 (3)	C2—C3—C4—C5	1.0 (3)
O2—N2—C1—N1	145.0 (2)	C1—N1—C5—C4	0.9 (3)

O3—N2—C1—C2	149.3 (2)	C1—N1—C5—N3	−177.3 (2)
O2—N2—C1—C2	−34.2 (4)	O1—C4—C5—N1	178.0 (2)
N1—C1—C2—C3	−3.1 (3)	C3—C4—C5—N1	−2.3 (3)
N2—C1—C2—C3	176.0 (2)	O1—C4—C5—N3	−4.0 (3)
N1—C1—C2—Cl1	172.71 (17)	C3—C4—C5—N3	175.7 (2)
N2—C1—C2—Cl1	−8.2 (3)	O5—N3—C5—N1	−64.2 (3)
C1—C2—C3—C4	1.4 (3)	O4—N3—C5—N1	115.0 (3)
Cl1—C2—C3—C4	−174.58 (16)	O5—N3—C5—C4	117.5 (3)
C6—O1—C4—C3	0.3 (3)	O4—N3—C5—C4	−63.3 (3)