

## 5-Dichloroacetyl-4-methyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-2-one hemihydrate

K. Ravichandran,<sup>a</sup> P. Sakthivel,<sup>b</sup> S. Ponnuswamy,<sup>b</sup>  
M. Shalini<sup>b</sup> and M. N. Ponnuswamy<sup>a\*</sup>

<sup>a</sup>Centre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and <sup>b</sup>Department of Chemistry, Government Arts College (Autonomous), Coimbatore 641 018, India

Correspondence e-mail: mnpsy2004@yahoo.com


Received 29 August 2009; accepted 12 September 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.045;  $wR$  factor = 0.110; data-to-parameter ratio = 15.9.

There are two crystallographically independent organic molecules in the asymmetric unit of the title compound,  $C_{12}H_{12}Cl_2N_2O_2 \cdot 0.5H_2O$ . The benzodiazepine ring adopts a distorted boat conformation in both molecules. The crystal packing is controlled by  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{O}$  intra- and intermolecular hydrogen bonds. A graph-set motif of  $R_3^3(14)$  dimer formation by a combination of  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds stabilizes the molecules and extends along a axis.

### Related literature

For the anticonvulsant activity of benzodiazepine, see: MacDonald (2002). For their hypnotic effect, see: Gringauz (1999). For their use in the treatment of gastrointestinal and central nervous system disorders, see: Rahbaek *et al.* (1999). For other therapeutic applications, see: Albright *et al.* (1998); Lee *et al.* (1999). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For puckering and asymmetry parameters, see: Cremer & Pople (1975); Nardelli (1983). For details of the preparation of the title compound, see: Venkatraj *et al.* (2008).



### Experimental

#### Crystal data

$C_{12}H_{12}Cl_2N_2O_2 \cdot 0.5H_2O$	$V = 1364.82(8)\text{ \AA}^3$
$M_r = 592.29$	$Z = 2$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 8.5470(3)\text{ \AA}$	$\mu = 0.48\text{ mm}^{-1}$
$b = 18.0837(6)\text{ \AA}$	$T = 293\text{ K}$
$c = 8.8697(3)\text{ \AA}$	$0.26 \times 0.24 \times 0.22\text{ mm}$
$\beta = 95.405(2)^\circ$	

#### Data collection

Bruker Kappa APEXII area-detector diffractometer	14191 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 2001)	5599 independent reflections
$T_{\min} = 0.884$ , $T_{\max} = 0.901$	4873 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.110$	$\Delta\rho_{\text{max}} = 0.46\text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\text{min}} = -0.62\text{ e \AA}^{-3}$
5599 reflections	Absolute structure: Flack (1983), 2698 Friedel pairs
352 parameters	Flack parameter: 0.06 (6)
1 restraint	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1A—H1A $\cdots$ O3	0.87 (4)	2.08 (4)	2.927 (4)	164 (3)
O3—H2W $\cdots$ O1B <sup>i</sup>	0.80 (4)	2.02 (4)	2.815 (4)	173 (4)
C8A—H8A $\cdots$ O2A <sup>ii</sup>	0.93	2.51	3.268 (4)	139
C10B—H10B $\cdots$ O2B <sup>iii</sup>	0.93	2.39	3.179 (4)	143

Symmetry codes: (i)  $-x + 2, y - \frac{1}{2}, -z + 2$ ; (ii)  $x - 1, y, z$ ; (iii)  $x + 1, y, z$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

KR thanks Dr Babu Varghese, SAIF, IIT-Madras, India, for his help with the data collection and the management of Kandaswami Kandar's College, Velur, Namakkal, India, for their encouragement to pursue the programme.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5048).

## References

- Albright, J. D., Feich, M. F., Santos, E. G. D., Dusza, J. P., Sum, F.-W., Venkatesan, A. M., Coupet, J., Chan, P. S., Ru, X., Mazandarani, H. & Bailey, T. (1998). *J. Med. Chem.* **41**, 2442–2444.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Gringauz, A. (1999). *Introduction to Medicinal Chemistry*, pp. 578–580. New York: Wiley-VCH.
- Lee, J., Gauthier, D. & Rivero, R. A. (1999). *J. Org. Chem.* **64**, 3060–3064.
- MacDonald, R. L. (2002). *Benzodiazepines Mechanisms of Action*. In *Antiepileptic Drugs*, 5th ed., edited by R. H. Levy, R. H. Mattson, B. S. Meldrum & E. Perucca, pp. 179–186. Philadelphia: Lippincott Williams and Wilkins.
- Nardelli, M. (1983). *Acta Cryst. C* **39**, 1141–1142.
- Rahbaek, L., Breinholt, J., Frisvad, J. C. & Christophersen, C. (1999). *J. Org. Chem.* **64**, 1689–1692.
- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Venkatraj, M., Ponnuswamy, S. & Jeyaraman, R. (2008). *Indian J. Chem. Sect. B*, **47**, 129–135.

# supporting information

*Acta Cryst.* (2009). E65, o2551–o2552 [doi:10.1107/S1600536809036940]

## 5-Dichloroacetyl-4-methyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-2-one hemihydrate

K. Ravichandran, P. Sakthivel, S. Ponnuswamy, M. Shalini and M. N. Ponnuswamy

### S1. Comment

The anticonvulsant activity of benzodiazepines has been utilized clinically in patients to treat specific seizure types or conditions, *i.e.*, akinetic, myoclonic, absence variant seizures as well as to help terminate status epilepticus or serial seizures (MacDonald, 2002). Benzodiazepines are used for the purpose of hypnotic effects, owing to their less toxic and less severe withdrawal effects when compared with barbiturates (Gringauz, 1999). Benzodiazepines from *aspergillus* include *asperlicin*, which is used for treatment of gastrointestinal and central nervous system (CNS) disorders (Rahbaek *et al.*, 1999). The other therapeutic applications (Lee *et al.*, 1999) of benzodiazepines include vasopressin antagonists (Albright *et al.*, 1998). In view of these importance and to ascertain the molecular conformation, crystallographic study of the title compound has been carried out.

The ORTEP diagram of the title compound is shown in Fig. 1. There are two crystallographically independent molecules in the asymmetric unit. The benzodiazepine rings in the two molecules adopt a distorted boat conformation. The puckering parameters (Cremer & Pople, 1975) and the asymmetry parameters (Nardelli, 1983) for the ring in molecule A are:  $q_2 = 0.959 (3)$  Å,  $q_3 = 0.150 (3)$  Å,  $\varphi_2 = 136.1 (2)$ °,  $\varphi_3 = 359.8 (1)$ ° and  $\Delta 2(C4A) = 8.1 (3)$ °; for the ring in molecule B are:  $q_2 = 0.962 (3)$  Å,  $q_3 = 0.168 (3)$  Å,  $\varphi_2 = 141.4 (2)$ °,  $\varphi_3 = 5.3 (1)$ ° and  $\Delta 2(C4B) = 3.4 (3)$ °. The sum of the bond angles at N1A(359.0°), N1B(359.2), N5A(358.8) and N5B(359.9°) of the benzodiazepine rings in both the molecules are in accordance with  $sp^2$  hybridization.


The crystal packing is controlled by N—H···O, C—H···O and O—H···O types of intermolecular interactions in addition to van der Waals forces. The water molecule connects the molecules A and B through N1A—H1A···O3 and O3—H2W···O1B hydrogen bonds. Thus the combination of N1A—H1A···O3, O3—H2W···O1B and C3A—H3A···O2B hydrogen bonds form a graph set motif of  $R^3_3(14)$  dimer (Bernstein *et al.*, 1995) which stabilize the molecules. Atom C8A at ( $x, y, z$ ) donates a proton to O2A ( $x - 1, y, z$ ), which forms a C7 one dimensional chain running along  $a$ -axis. The intermolecular hydrogen bond C10B—H10B···O2B also connects the molecule into another C7 chain running along  $b$ -axis (Fig. 2).

### S2. Experimental


To a solution of tetrahydro-4-methyl-1,5-benzodiazepin-2-one (0.88 g) in anhydrous benzene (50 ml) was added triethylamine (2.8 ml) and dichloroacetyl chloride (1.90 ml). The contents were allowed to reflux on a water bath for 6 hrs. The reaction mixture was washed with sodium bicarbonate solution (10%), water and dried. Evaporation of the solvent results a crude mass and further crystallization from ethanol gives colorless crystals (Venkatraj *et al.*, 2008).

**S3. Refinement**

The Nitrogen and Oxygen H atoms were refined and the other H atoms positioned geometrically ( $C-H=0.93-0.98 \text{ \AA}$ ) and allowed to ride on their parent atoms, with  $1.5U_{\text{eq}}(\text{C})$  for methyl H and  $1.2U_{\text{eq}}(\text{C})$  for other H atoms.

**Figure 1**

Perspective view of the molecule showing the thermal ellipsoids are drawn at 30% probability level.

**Figure 2**

The crystal packing of the molecules viewed down c-axis. H atoms not involved in hydrogen bonding have been omitted for clarity.

**5-Dichloroacetyl-4-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-2-one hemihydrate***Crystal data*
 $M_r = 592.29$ 

Monoclinic,  $P2_1$ 

Hall symbol: P 2yb

 $a = 8.5470 (3) \text{ \AA}$ 
 $b = 18.0837 (6) \text{ \AA}$ 
 $c = 8.8697 (3) \text{ \AA}$ 
 $\beta = 95.405 (2)^\circ$ 
 $V = 1364.82 (8) \text{ \AA}^3$ 
 $Z = 2$ 
 $F(000) = 612$ 
 $D_x = 1.441 \text{ Mg m}^{-3}$ 

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$ 

Cell parameters from 4629 reflections

 $\theta = 2.3\text{--}26.5^\circ$ 
 $\mu = 0.48 \text{ mm}^{-1}$ 
 $T = 293 \text{ K}$ 

Block, colourless

 $0.26 \times 0.24 \times 0.22 \text{ mm}$ 
*Data collection*
Bruker Kappa APEXII area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  and  $\varphi$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 2001)

 $T_{\min} = 0.884$ ,  $T_{\max} = 0.901$ 

14191 measured reflections

5599 independent reflections

4873 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.022$ 
 $\theta_{\max} = 26.5^\circ$ ,  $\theta_{\min} = 2.3^\circ$ 
 $h = -10 \rightarrow 10$ 
 $k = -22 \rightarrow 22$ 
 $l = -11 \rightarrow 10$ 
*Refinement*
Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.045$ 
 $wR(F^2) = 0.110$ 
 $S = 1.04$ 

5599 reflections

352 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H atoms treated by a mixture of independent  
and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0438P)^2 + 0.7028P]$   
where  $P = (F_o^2 + 2F_c^2)/3$ 
 $(\Delta/\sigma)_{\max} = 0.006$ 
 $\Delta\rho_{\max} = 0.46 \text{ e \AA}^{-3}$ 
 $\Delta\rho_{\min} = -0.62 \text{ e \AA}^{-3}$ 

Absolute structure: Flack (1983), 2698 Friedel  
pairs

Absolute structure parameter: 0.06 (6)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1A	1.62028 (16)	-0.29670 (7)	1.77856 (13)	0.0941 (4)
Cl1B	1.04826 (15)	0.13662 (6)	0.96558 (15)	0.0906 (4)
Cl2A	1.48485 (16)	-0.15064 (6)	1.74118 (13)	0.0858 (3)

Cl2B	0.8172 (2)	0.15716 (9)	1.17529 (14)	0.1300 (7)
O1A	1.5519 (3)	-0.46993 (14)	1.2925 (3)	0.0594 (6)
O1B	1.0719 (3)	-0.07368 (15)	0.6539 (2)	0.0573 (6)
O2A	1.6504 (3)	-0.20942 (14)	1.4809 (3)	0.0635 (7)
O2B	0.7429 (3)	0.02505 (17)	1.0049 (4)	0.0852 (9)
O3	1.2224 (4)	-0.56267 (15)	1.5161 (4)	0.0677 (7)
H1W	1.279 (6)	-0.601 (3)	1.502 (6)	0.108 (19)*
H2W	1.140 (5)	-0.570 (2)	1.467 (4)	0.059 (11)*
N1A	1.3405 (3)	-0.42583 (13)	1.3919 (3)	0.0403 (5)
H1A	1.325 (4)	-0.469 (2)	1.432 (4)	0.057 (10)*
N1B	1.2097 (3)	-0.03383 (14)	0.8658 (3)	0.0399 (5)
H1B	1.261 (3)	-0.0095 (17)	0.811 (3)	0.031 (7)*
C2A	1.4518 (3)	-0.42199 (16)	1.2939 (3)	0.0420 (6)
C2B	1.1026 (3)	-0.07798 (16)	0.7907 (3)	0.0410 (6)
C3A	1.4398 (4)	-0.35821 (17)	1.1848 (3)	0.0462 (7)
H3A	1.3314	-0.3539	1.1423	0.055*
H3B	1.5028	-0.3694	1.1023	0.055*
C3B	1.0275 (4)	-0.13513 (16)	0.8847 (3)	0.0465 (7)
H3C	1.1089	-0.1594	0.9505	0.056*
H3D	0.9773	-0.1723	0.8178	0.056*
C4A	1.4920 (3)	-0.28414 (16)	1.2521 (3)	0.0441 (7)
H4A	1.6071	-0.2832	1.2637	0.053*
C4B	0.9057 (4)	-0.10270 (17)	0.9813 (4)	0.0490 (7)
H4B	0.8130	-0.0886	0.9140	0.059*
N5A	1.4357 (2)	-0.27668 (12)	1.4044 (2)	0.0368 (5)
N5B	0.9699 (3)	-0.03578 (14)	1.0580 (2)	0.0403 (5)
C6A	1.2763 (3)	-0.29648 (16)	1.4220 (3)	0.0355 (5)
C6B	1.1307 (3)	-0.03545 (15)	1.1206 (3)	0.0363 (6)
C7A	1.1664 (3)	-0.24204 (17)	1.4443 (3)	0.0457 (7)
H7A	1.1961	-0.1926	1.4462	0.055*
C7B	1.1690 (4)	-0.03548 (18)	1.2758 (3)	0.0485 (7)
H7B	1.0906	-0.0415	1.3407	0.058*
C8A	1.0148 (3)	-0.2607 (2)	1.4635 (4)	0.0548 (8)
H8A	0.9424	-0.2242	1.4820	0.066*
C8B	1.3225 (4)	-0.0267 (2)	1.3344 (4)	0.0587 (9)
H8B	1.3482	-0.0259	1.4386	0.070*
C9A	0.9695 (3)	-0.3334 (2)	1.4554 (4)	0.0530 (8)
H9A	0.8658	-0.3458	1.4675	0.064*
C9B	1.4383 (4)	-0.01891 (19)	1.2369 (4)	0.0571 (9)
H9B	1.5418	-0.0109	1.2758	0.069*
C10A	1.0749 (3)	-0.38800 (17)	1.4298 (4)	0.0467 (7)
H10A	1.0422	-0.4370	1.4237	0.056*
C10B	1.4017 (3)	-0.02294 (17)	1.0826 (4)	0.0461 (7)
H10B	1.4812	-0.0194	1.0182	0.055*
C11A	1.2304 (3)	-0.37036 (15)	1.4130 (3)	0.0353 (5)
C11B	1.2482 (3)	-0.03216 (15)	1.0231 (3)	0.0363 (5)
C12A	1.4358 (6)	-0.2213 (2)	1.1517 (4)	0.0707 (11)
H12A	1.3230	-0.2210	1.1396	0.106*

H12B	1.4751	-0.2271	1.0544	0.106*
H12C	1.4733	-0.1755	1.1964	0.106*
C12B	0.8546 (5)	-0.1577 (3)	1.0958 (5)	0.0806 (12)
H12D	0.9429	-0.1705	1.1660	0.121*
H12E	0.8147	-0.2015	1.0443	0.121*
H12F	0.7738	-0.1361	1.1498	0.121*
C13A	1.5282 (3)	-0.23943 (16)	1.5082 (3)	0.0425 (6)
C13B	0.8785 (3)	0.02480 (19)	1.0567 (3)	0.0483 (7)
C14A	1.4827 (4)	-0.2412 (2)	1.6705 (3)	0.0516 (7)
H14A	1.3773	-0.2624	1.6720	0.062*
C14B	0.9587 (5)	0.0969 (2)	1.1152 (4)	0.0651 (10)
H14B	1.0379	0.0860	1.1995	0.078*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1A	0.1134 (9)	0.0939 (8)	0.0740 (7)	0.0444 (7)	0.0047 (6)	0.0100 (6)
Cl1B	0.1093 (9)	0.0570 (6)	0.1045 (8)	-0.0225 (6)	0.0047 (7)	0.0035 (6)
Cl2A	0.1158 (9)	0.0691 (6)	0.0740 (6)	0.0202 (6)	0.0169 (6)	-0.0282 (5)
Cl2B	0.1609 (13)	0.1493 (13)	0.0748 (7)	0.1052 (12)	-0.0141 (7)	-0.0412 (8)
O1A	0.0640 (14)	0.0558 (13)	0.0609 (13)	0.0228 (12)	0.0193 (11)	0.0018 (11)
O1B	0.0562 (13)	0.0768 (16)	0.0377 (11)	-0.0032 (12)	-0.0020 (9)	-0.0029 (11)
O2A	0.0445 (12)	0.0707 (16)	0.0778 (16)	-0.0228 (11)	0.0190 (11)	-0.0222 (13)
O2B	0.0332 (12)	0.087 (2)	0.133 (3)	0.0137 (13)	-0.0012 (14)	0.0200 (19)
O3	0.0578 (16)	0.0485 (14)	0.093 (2)	0.0005 (13)	-0.0113 (15)	0.0087 (13)
N1A	0.0465 (13)	0.0319 (13)	0.0435 (13)	0.0045 (10)	0.0100 (10)	0.0037 (10)
N1B	0.0367 (12)	0.0480 (13)	0.0360 (12)	-0.0030 (11)	0.0087 (10)	0.0039 (11)
C2A	0.0474 (16)	0.0378 (15)	0.0408 (14)	0.0031 (13)	0.0035 (12)	-0.0043 (12)
C2B	0.0388 (14)	0.0439 (16)	0.0400 (15)	0.0031 (12)	0.0026 (11)	-0.0056 (12)
C3A	0.0575 (18)	0.0471 (16)	0.0348 (13)	0.0020 (14)	0.0083 (12)	-0.0023 (12)
C3B	0.0504 (17)	0.0371 (15)	0.0511 (16)	-0.0102 (13)	-0.0010 (13)	-0.0056 (12)
C4A	0.0461 (16)	0.0472 (17)	0.0404 (14)	-0.0062 (13)	0.0108 (12)	-0.0014 (13)
C4B	0.0382 (15)	0.0495 (18)	0.0588 (19)	-0.0087 (13)	0.0016 (13)	0.0007 (14)
N5A	0.0328 (11)	0.0392 (12)	0.0394 (11)	-0.0022 (9)	0.0082 (9)	-0.0032 (9)
N5B	0.0282 (11)	0.0521 (14)	0.0409 (12)	0.0018 (10)	0.0047 (9)	0.0010 (11)
C6A	0.0277 (12)	0.0417 (14)	0.0367 (13)	0.0022 (11)	0.0010 (10)	0.0018 (11)
C6B	0.0334 (13)	0.0375 (13)	0.0372 (13)	0.0031 (11)	-0.0004 (10)	0.0018 (11)
C7A	0.0420 (15)	0.0402 (16)	0.0552 (17)	0.0070 (12)	0.0057 (13)	0.0037 (13)
C7B	0.0540 (17)	0.0523 (17)	0.0390 (14)	0.0062 (14)	0.0031 (12)	0.0047 (13)
C8A	0.0372 (16)	0.057 (2)	0.071 (2)	0.0182 (14)	0.0091 (14)	0.0025 (16)
C8B	0.071 (2)	0.060 (2)	0.0419 (16)	0.0053 (18)	-0.0129 (15)	-0.0005 (15)
C9A	0.0292 (15)	0.065 (2)	0.0648 (19)	-0.0016 (14)	0.0056 (13)	0.0010 (17)
C9B	0.0434 (17)	0.057 (2)	0.066 (2)	-0.0052 (15)	-0.0182 (15)	0.0001 (16)
C10A	0.0376 (15)	0.0460 (17)	0.0563 (17)	-0.0072 (12)	0.0031 (13)	-0.0004 (13)
C10B	0.0328 (13)	0.0472 (17)	0.0568 (17)	-0.0013 (12)	-0.0033 (12)	0.0023 (14)
C11A	0.0356 (13)	0.0376 (14)	0.0326 (12)	-0.0005 (11)	0.0032 (10)	-0.0005 (11)
C11B	0.0342 (13)	0.0347 (13)	0.0393 (13)	0.0007 (11)	0.0009 (10)	0.0015 (11)
C12A	0.104 (3)	0.049 (2)	0.061 (2)	-0.0015 (19)	0.019 (2)	0.0172 (17)

C12B	0.083 (3)	0.075 (3)	0.087 (3)	-0.036 (2)	0.026 (2)	0.004 (2)
C13A	0.0333 (14)	0.0402 (15)	0.0551 (16)	-0.0060 (12)	0.0107 (12)	-0.0084 (13)
C13B	0.0353 (15)	0.0581 (19)	0.0525 (17)	0.0096 (14)	0.0086 (13)	0.0103 (15)
C14A	0.0420 (15)	0.064 (2)	0.0482 (16)	0.0011 (14)	0.0015 (12)	-0.0146 (15)
C14B	0.073 (2)	0.064 (2)	0.0539 (19)	0.0303 (18)	-0.0134 (17)	-0.0112 (16)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C11A—C14A	1.759 (3)	N5B—C13B	1.345 (4)
C11B—C14B	1.748 (4)	N5B—C6B	1.432 (3)
C12A—C14A	1.754 (3)	C6A—C7A	1.388 (4)
C12B—C14B	1.748 (4)	C6A—C11A	1.393 (4)
O1A—C2A	1.219 (4)	C6B—C7B	1.385 (4)
O1B—C2B	1.220 (3)	C6B—C11B	1.387 (4)
O2A—C13A	1.221 (3)	C7A—C8A	1.365 (4)
O2B—C13B	1.206 (4)	C7A—H7A	0.9300
O3—H1W	0.86 (6)	C7B—C8B	1.374 (5)
O3—H2W	0.80 (4)	C7B—H7B	0.9300
N1A—C2A	1.349 (4)	C8A—C9A	1.370 (5)
N1A—C11A	1.400 (4)	C8A—H8A	0.9300
N1A—H1A	0.87 (4)	C8B—C9B	1.381 (5)
N1B—C2B	1.343 (4)	C8B—H8B	0.9300
N1B—C11B	1.403 (3)	C9A—C10A	1.370 (4)
N1B—H1B	0.81 (3)	C9A—H9A	0.9300
C2A—C3A	1.503 (4)	C9B—C10B	1.376 (5)
C2B—C3B	1.508 (4)	C9B—H9B	0.9300
C3A—C4A	1.516 (4)	C10A—C11A	1.389 (4)
C3A—H3A	0.9700	C10A—H10A	0.9300
C3A—H3B	0.9700	C10B—C11B	1.378 (4)
C3B—C4B	1.527 (5)	C10B—H10B	0.9300
C3B—H3C	0.9700	C12A—H12A	0.9600
C3B—H3D	0.9700	C12A—H12B	0.9600
C4A—N5A	1.482 (3)	C12A—H12C	0.9600
C4A—C12A	1.495 (5)	C12B—H12D	0.9600
C4A—H4A	0.9800	C12B—H12E	0.9600
C4B—N5B	1.469 (4)	C12B—H12F	0.9600
C4B—C12B	1.515 (5)	C13A—C14A	1.526 (4)
C4B—H4B	0.9800	C13B—C14B	1.540 (5)
N5A—C13A	1.338 (4)	C14A—H14A	0.9800
N5A—C6A	1.431 (3)	C14B—H14B	0.9800
H1W—O3—H2W	106 (5)	C7A—C8A—C9A	119.9 (3)
C2A—N1A—C11A	125.0 (2)	C7A—C8A—H8A	120.1
C2A—N1A—H1A	117 (2)	C9A—C8A—H8A	120.1
C11A—N1A—H1A	117 (2)	C7B—C8B—C9B	119.4 (3)
C2B—N1B—C11B	126.2 (3)	C7B—C8B—H8B	120.3
C2B—N1B—H1B	114 (2)	C9B—C8B—H8B	120.3
C11B—N1B—H1B	119 (2)	C10A—C9A—C8A	120.9 (3)

O1A—C2A—N1A	120.5 (3)	C10A—C9A—H9A	119.6
O1A—C2A—C3A	123.0 (3)	C8A—C9A—H9A	119.6
N1A—C2A—C3A	116.4 (3)	C10B—C9B—C8B	120.5 (3)
O1B—C2B—N1B	121.9 (3)	C10B—C9B—H9B	119.7
O1B—C2B—C3B	122.0 (3)	C8B—C9B—H9B	119.7
N1B—C2B—C3B	116.1 (2)	C9A—C10A—C11A	120.2 (3)
C2A—C3A—C4A	115.1 (2)	C9A—C10A—H10A	119.9
C2A—C3A—H3A	108.5	C11A—C10A—H10A	119.9
C4A—C3A—H3A	108.5	C9B—C10B—C11B	120.4 (3)
C2A—C3A—H3B	108.5	C9B—C10B—H10B	119.8
C4A—C3A—H3B	108.5	C11B—C10B—H10B	119.8
H3A—C3A—H3B	107.5	C10A—C11A—C6A	118.8 (2)
C2B—C3B—C4B	113.3 (2)	C10A—C11A—N1A	120.8 (3)
C2B—C3B—H3C	108.9	C6A—C11A—N1A	120.4 (2)
C4B—C3B—H3C	108.9	C10B—C11B—C6B	119.0 (3)
C2B—C3B—H3D	108.9	C10B—C11B—N1B	120.6 (3)
C4B—C3B—H3D	108.9	C6B—C11B—N1B	120.3 (2)
H3C—C3B—H3D	107.7	C4A—C12A—H12A	109.5
N5A—C4A—C12A	111.1 (3)	C4A—C12A—H12B	109.5
N5A—C4A—C3A	109.3 (2)	H12A—C12A—H12B	109.5
C12A—C4A—C3A	111.8 (3)	C4A—C12A—H12C	109.5
N5A—C4A—H4A	108.2	H12A—C12A—H12C	109.5
C12A—C4A—H4A	108.2	H12B—C12A—H12C	109.5
C3A—C4A—H4A	108.2	C4B—C12B—H12D	109.5
N5B—C4B—C12B	110.4 (3)	C4B—C12B—H12E	109.5
N5B—C4B—C3B	109.4 (2)	H12D—C12B—H12E	109.5
C12B—C4B—C3B	112.3 (3)	C4B—C12B—H12F	109.5
N5B—C4B—H4B	108.2	H12D—C12B—H12F	109.5
C12B—C4B—H4B	108.2	H12E—C12B—H12F	109.5
C3B—C4B—H4B	108.2	O2A—C13A—N5A	123.3 (3)
C13A—N5A—C6A	123.9 (2)	O2A—C13A—C14A	119.7 (3)
C13A—N5A—C4A	116.8 (2)	N5A—C13A—C14A	116.9 (2)
C6A—N5A—C4A	118.1 (2)	O2B—C13B—N5B	122.9 (3)
C13B—N5B—C6B	122.4 (3)	O2B—C13B—C14B	120.4 (3)
C13B—N5B—C4B	118.4 (2)	N5B—C13B—C14B	116.5 (3)
C6B—N5B—C4B	119.1 (2)	C13A—C14A—C12A	108.8 (2)
C7A—C6A—C11A	119.8 (2)	C13A—C14A—C11A	108.0 (2)
C7A—C6A—N5A	120.2 (3)	C12A—C14A—C11A	110.73 (17)
C11A—C6A—N5A	120.0 (2)	C13A—C14A—H14A	109.7
C7B—C6B—C11B	120.2 (2)	C12A—C14A—H14A	109.7
C7B—C6B—N5B	120.9 (3)	C11A—C14A—H14A	109.7
C11B—C6B—N5B	118.8 (2)	C13B—C14B—C12B	109.4 (3)
C8A—C7A—C6A	120.4 (3)	C13B—C14B—C11B	107.7 (2)
C8A—C7A—H7A	119.8	C12B—C14B—C11B	109.9 (2)
C6A—C7A—H7A	119.8	C13B—C14B—H14B	109.9
C8B—C7B—C6B	120.2 (3)	C12B—C14B—H14B	109.9
C8B—C7B—H7B	119.9	C11B—C14B—H14B	109.9
C6B—C7B—H7B	119.9		

C11A—N1A—C2A—O1A	173.3 (3)	C7B—C8B—C9B—C10B	−2.6 (5)
C11A—N1A—C2A—C3A	−9.3 (4)	C8A—C9A—C10A—C11A	0.6 (5)
C11B—N1B—C2B—O1B	177.0 (3)	C8B—C9B—C10B—C11B	2.3 (5)
C11B—N1B—C2B—C3B	−5.4 (4)	C9A—C10A—C11A—C6A	−0.2 (4)
O1A—C2A—C3A—C4A	−106.8 (3)	C9A—C10A—C11A—N1A	177.8 (3)
N1A—C2A—C3A—C4A	75.9 (3)	C7A—C6A—C11A—C10A	−1.4 (4)
O1B—C2B—C3B—C4B	−107.2 (3)	N5A—C6A—C11A—C10A	−179.7 (2)
N1B—C2B—C3B—C4B	75.2 (3)	C7A—C6A—C11A—N1A	−179.4 (2)
C2A—C3A—C4A—N5A	−40.6 (4)	N5A—C6A—C11A—N1A	2.2 (4)
C2A—C3A—C4A—C12A	−164.0 (3)	C2A—N1A—C11A—C10A	138.6 (3)
C2B—C3B—C4B—N5B	−46.2 (3)	C2A—N1A—C11A—C6A	−43.4 (4)
C2B—C3B—C4B—C12B	−169.2 (3)	C9B—C10B—C11B—C6B	1.6 (4)
C12A—C4A—N5A—C13A	−90.1 (3)	C9B—C10B—C11B—N1B	177.9 (3)
C3A—C4A—N5A—C13A	146.1 (3)	C7B—C6B—C11B—C10B	−5.2 (4)
C12A—C4A—N5A—C6A	78.1 (3)	N5B—C6B—C11B—C10B	172.3 (3)
C3A—C4A—N5A—C6A	−45.7 (3)	C7B—C6B—C11B—N1B	178.5 (3)
C12B—C4B—N5B—C13B	−102.0 (3)	N5B—C6B—C11B—N1B	−3.9 (4)
C3B—C4B—N5B—C13B	134.0 (3)	C2B—N1B—C11B—C10B	141.2 (3)
C12B—C4B—N5B—C6B	83.0 (3)	C2B—N1B—C11B—C6B	−42.6 (4)
C3B—C4B—N5B—C6B	−41.1 (3)	C6A—N5A—C13A—O2A	−164.1 (3)
C13A—N5A—C6A—C7A	58.9 (4)	C4A—N5A—C13A—O2A	3.3 (4)
C4A—N5A—C6A—C7A	−108.3 (3)	C6A—N5A—C13A—C14A	21.3 (4)
C13A—N5A—C6A—C11A	−122.8 (3)	C4A—N5A—C13A—C14A	−171.3 (3)
C4A—N5A—C6A—C11A	70.0 (3)	C6B—N5B—C13B—O2B	−179.7 (3)
C13B—N5B—C6B—C7B	75.4 (4)	C4B—N5B—C13B—O2B	5.4 (4)
C4B—N5B—C6B—C7B	−109.8 (3)	C6B—N5B—C13B—C14B	4.5 (4)
C13B—N5B—C6B—C11B	−102.1 (3)	C4B—N5B—C13B—C14B	−170.4 (3)
C4B—N5B—C6B—C11B	72.7 (3)	O2A—C13A—C14A—Cl2A	53.9 (3)
C11A—C6A—C7A—C8A	2.6 (4)	N5A—C13A—C14A—Cl2A	−131.2 (2)
N5A—C6A—C7A—C8A	−179.0 (3)	O2A—C13A—C14A—Cl1A	−66.3 (3)
C11B—C6B—C7B—C8B	4.9 (5)	N5A—C13A—C14A—Cl1A	108.5 (3)
N5B—C6B—C7B—C8B	−172.6 (3)	O2B—C13B—C14B—Cl2B	27.4 (4)
C6A—C7A—C8A—C9A	−2.3 (5)	N5B—C13B—C14B—Cl2B	−156.7 (2)
C6B—C7B—C8B—C9B	−1.0 (5)	O2B—C13B—C14B—Cl1B	−92.0 (3)
C7A—C8A—C9A—C10A	0.7 (5)	N5B—C13B—C14B—Cl1B	84.0 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C4A—H4A···O2A	0.98	2.34	2.694 (4)	100
C4B—H4B···O2B	0.98	2.31	2.715 (4)	104
N1A—H1A···O3	0.87 (4)	2.08 (4)	2.927 (4)	164 (3)
O3—H2W···O1B <sup>i</sup>	0.80 (4)	2.02 (4)	2.815 (4)	173 (4)
C3A—H3A···O2B <sup>i</sup>	0.97	2.60	3.038 (4)	108

---

C8A—H8A···O2A <sup>ii</sup>	0.93	2.51	3.268 (4)	139
C10B—H10B···O2B <sup>iii</sup>	0.93	2.39	3.179 (4)	143

---

Symmetry codes: (i)  $-x+2, y-1/2, -z+2$ ; (ii)  $x-1, y, z$ ; (iii)  $x+1, y, z$ .