

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Ethoxy-2-(1,3-thiazol-2-yl)isoindolin-1-one

Wenkuan Li, Handong Yin,* Liyuan Wen, Jichun Cui and Daqi Wang

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: handongyin@163.com


Received 13 September 2009; accepted 25 September 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.042; wR factor = 0.117; data-to-parameter ratio = 13.6.

In the title compound, $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$, the dihedral angles between the isoindolone ring system and the thiazole ring and the ethoxy group are 6.50 (11) and 89.0 (2)°, respectively.

Related literature

For general background to isoindolin-1-one derivatives, see: Gai *et al.* (2003). For hybridization, see: Beddoes *et al.* (1986).



Experimental

Crystal data

 $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$ $M_r = 260.31$ Monoclinic, $P2_1/n$ $a = 8.0933$ (11) Å $b = 9.1406$ (14) Å $c = 17.2077$ (19) Å $\beta = 98.720$ (1)° $V = 1258.3$ (3) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.25$ mm⁻¹ $T = 298$ K $0.50 \times 0.49 \times 0.47$ mm

Data collection

Siemens SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.884$, $T_{\max} = 0.891$

6305 measured reflections

2236 independent reflections

1554 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.031$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$ $wR(F^2) = 0.117$ $S = 1.04$

2236 reflections

164 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge financial support by the Natural Science Foundation of China (No. 20771053) and the Natural Science Foundation of Shandong Province (Y2008B48).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2243).

References

- Beddoes, R. L., Dalton, L., Joule, T. A., Mills, O. S., Street, J. D. & Watt, C. I. F. (1986). *J. Chem. Soc. Perkin Trans. 2*, pp. 787–797.
- Gai, X., Grigg, R., Khamnaen, T., Rajviroongit, S., Sridharan, V., Zhang, L., Collard, S. & Keep, A. (2003). *Tetrahedron Lett.* **44**, 7441–7443.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). SMART and SAINT. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

supporting information

Acta Cryst. (2009). E65, o2594 [doi:10.1107/S1600536809038884]

3-Ethoxy-2-(1,3-thiazol-2-yl)isoindolin-1-one

Wenkuan Li, Handong Yin, Liyuan Wen, Jichun Cui and Daqi Wang

S1. Comment


The title compound, (I), was formed by accident, instead of the organotin compound containing schiff base which was expected. Isoindolin-1-one derivatives have been demonstrated to possess anxiolytic activity and are of interest as sedatives, hypnotics and muscle relaxants. In addition, the isoindolone moiety also features in anti-cancer drug candidates including protein kinase inhibitors (Gai *et al.* 2003). We have synthesized the title compound (I) and its crystal structure is reported herein. The molecular structure of (I) is shown in Fig.1. The dihedral angles between the isoindolone ring system and thiazole ring and oxyethyl group are 6.50 (11) and 89.0 (2)° respectively. The sum of bond angles around atom N1 (359.6 °) indicates that the atom N1 is in sp^2 hybridized state (Beddoes *et al.* 1986). The crystal packing is stabilized mainly by van der Waals interactions.

S2. Experimental

(*E*)-2-((thiazol-2-ylimino)methyl)benzoic acid (4 mmol) and sodium ethoxide (4 mmol) were added to a stirred solution of ethanol (30 ml) in a Schlenk flask and stirred for 0.5 h. Chlorotriphenyltin (4 mmol) was then added to the reactor and the reaction mixture was heated under reflux for 6 h. The resulting clear solution was evaporated under vacuum. The product was crystallized from a mixed of dichloromethane/petroleum (1:1) to afford the title compound unexpectedly. Anal. Calcd (%) for $C_{13}H_{12}N_2O_2S$ (Mr = 260.31): C, 59.98; H, 4.65; N, 10.76; O, 12.29; S, 12.32 Found (%): C, 60.00; H, 4.62; N, 10.74; O, 12.31; S, 12.30

S3. Refinement

H atoms were positioned geometrically, with C—H = 0.93, 0.96, 0.97 and 0.98 Å for aromatic, methyl, methylene and methine H atoms, respectively, and constrained to ride on their parent atoms, with $U_{iso}(H) = x U_{eq}(C)$ where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

**Figure 1**

The molecular structure of the compound, showing 50% probability displacement ellipsoids. H atoms have been omitted for clarity.

3-Ethoxy-2-(1,3-thiazol-2-yl)isoindolin-1-one

Crystal data

$C_{13}H_{12}N_2O_2S$

$M_r = 260.31$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 8.0933$ (11) Å

$b = 9.1406$ (14) Å

$c = 17.2077$ (19) Å

$\beta = 98.720$ (1)°

$V = 1258.3$ (3) Å³

$Z = 4$

$F(000) = 544$

$D_x = 1.374$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2091 reflections

$\theta = 2.5$ – 26.6 °

$\mu = 0.25$ mm⁻¹

$T = 298$ K

Block, colourless

$0.50 \times 0.49 \times 0.47$ mm

Data collection

Siemens SMART CCD area-detector

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.884$, $T_{\max} = 0.891$

6305 measured reflections

2236 independent reflections

1554 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\text{max}} = 25.1$ °, $\theta_{\text{min}} = 2.4$ °

$h = -9 \rightarrow 9$

$k = -10 \rightarrow 8$

$l = -20 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.117$

$S = 1.04$

2236 reflections

164 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0456P)^2 + 0.5551P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.40181 (10)	0.62694 (9)	0.23706 (4)	0.0734 (3)
N1	0.5832 (2)	0.5184 (2)	0.14152 (12)	0.0558 (6)
N2	0.3056 (2)	0.4365 (2)	0.11652 (11)	0.0482 (5)
O1	0.0910 (2)	0.5044 (2)	0.18270 (11)	0.0663 (5)
O2	0.44903 (19)	0.23876 (18)	0.06204 (9)	0.0515 (4)
C1	0.4324 (3)	0.5174 (3)	0.15857 (14)	0.0479 (6)
C2	0.6825 (3)	0.6082 (3)	0.19276 (15)	0.0603 (7)
H2	0.7953	0.6217	0.1899	0.072*
C3	0.6079 (4)	0.6747 (4)	0.24650 (17)	0.0713 (8)
H3	0.6608	0.7387	0.2843	0.086*
C4	0.1397 (3)	0.4400 (3)	0.12897 (15)	0.0517 (6)
C5	0.0457 (3)	0.3538 (3)	0.06545 (14)	0.0491 (6)
C6	0.1552 (3)	0.3004 (3)	0.01772 (14)	0.0484 (6)
C7	0.3314 (3)	0.3506 (3)	0.04657 (13)	0.0463 (6)
H7	0.3668	0.4169	0.0075	0.056*
C8	-0.1242 (3)	0.3255 (3)	0.04990 (16)	0.0580 (7)
H8	-0.1971	0.3625	0.0819	0.070*
C9	-0.1811 (3)	0.2406 (3)	-0.01463 (18)	0.0677 (8)
H9	-0.2947	0.2200	-0.0265	0.081*
C10	-0.0725 (4)	0.1850 (3)	-0.06238 (18)	0.0689 (8)
H10	-0.1140	0.1277	-0.1056	0.083*
C11	0.0976 (3)	0.2142 (3)	-0.04631 (16)	0.0605 (7)
H11	0.1708	0.1764	-0.0780	0.073*
C12	0.4196 (4)	0.1366 (3)	0.12151 (17)	0.0682 (8)
H12A	0.3049	0.1023	0.1115	0.082*
H12B	0.4381	0.1833	0.1727	0.082*
C13	0.5357 (4)	0.0121 (3)	0.12016 (18)	0.0747 (9)
H13A	0.5135	-0.0360	0.0701	0.112*
H13B	0.5203	-0.0559	0.1610	0.112*
H13C	0.6488	0.0473	0.1286	0.112*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0675 (5)	0.0909 (6)	0.0664 (5)	0.0092 (4)	0.0252 (4)	-0.0195 (4)
N1	0.0455 (12)	0.0655 (14)	0.0595 (13)	0.0005 (10)	0.0185 (10)	-0.0066 (11)
N2	0.0414 (11)	0.0518 (12)	0.0558 (12)	0.0053 (9)	0.0216 (9)	0.0009 (10)
O1	0.0562 (11)	0.0790 (13)	0.0707 (12)	0.0133 (10)	0.0320 (9)	-0.0043 (10)
O2	0.0436 (9)	0.0534 (10)	0.0618 (10)	0.0085 (8)	0.0218 (8)	0.0015 (8)
C1	0.0495 (14)	0.0485 (14)	0.0489 (13)	0.0101 (11)	0.0179 (11)	0.0071 (11)
C2	0.0508 (15)	0.0687 (18)	0.0615 (16)	-0.0015 (13)	0.0088 (13)	-0.0059 (14)
C3	0.0709 (19)	0.082 (2)	0.0600 (17)	0.0074 (16)	0.0064 (14)	-0.0130 (16)
C4	0.0445 (14)	0.0542 (15)	0.0612 (15)	0.0114 (12)	0.0237 (12)	0.0133 (13)
C5	0.0424 (13)	0.0491 (14)	0.0590 (15)	0.0065 (11)	0.0177 (11)	0.0150 (12)
C6	0.0435 (14)	0.0473 (14)	0.0570 (15)	0.0023 (11)	0.0164 (11)	0.0109 (12)
C7	0.0408 (13)	0.0480 (14)	0.0536 (14)	0.0065 (11)	0.0183 (11)	0.0053 (11)
C8	0.0441 (15)	0.0623 (17)	0.0712 (18)	0.0083 (13)	0.0200 (13)	0.0185 (15)
C9	0.0441 (15)	0.076 (2)	0.084 (2)	-0.0033 (14)	0.0124 (14)	0.0152 (17)
C10	0.0607 (18)	0.0698 (19)	0.0747 (19)	-0.0109 (15)	0.0059 (14)	-0.0009 (16)
C11	0.0529 (16)	0.0644 (17)	0.0672 (17)	0.0014 (13)	0.0186 (13)	0.0014 (15)
C12	0.0670 (18)	0.0639 (18)	0.0779 (19)	0.0119 (14)	0.0242 (15)	0.0180 (15)
C13	0.078 (2)	0.0633 (19)	0.078 (2)	0.0156 (16)	-0.0028 (16)	0.0035 (16)

Geometric parameters (\AA , $^\circ$)

S1—C3	1.708 (3)	C6—C11	1.377 (4)
S1—C1	1.729 (2)	C6—C7	1.509 (3)
N1—C1	1.298 (3)	C7—H7	0.9800
N1—C2	1.372 (3)	C8—C9	1.376 (4)
N2—C1	1.378 (3)	C8—H8	0.9300
N2—C4	1.392 (3)	C9—C10	1.388 (4)
N2—C7	1.478 (3)	C9—H9	0.9300
O1—C4	1.211 (3)	C10—C11	1.388 (4)
O2—C7	1.395 (3)	C10—H10	0.9300
O2—C12	1.432 (3)	C11—H11	0.9300
C2—C3	1.326 (4)	C12—C13	1.479 (4)
C2—H2	0.9300	C12—H12A	0.9700
C3—H3	0.9300	C12—H12B	0.9700
C4—C5	1.464 (4)	C13—H13A	0.9600
C5—C6	1.385 (3)	C13—H13B	0.9600
C5—C8	1.385 (3)	C13—H13C	0.9600
C3—S1—C1	88.22 (13)	O2—C7—H7	108.8
C1—N1—C2	109.8 (2)	N2—C7—H7	108.8
C1—N2—C4	124.5 (2)	C6—C7—H7	108.8
C1—N2—C7	121.80 (17)	C9—C8—C5	117.6 (2)
C4—N2—C7	113.3 (2)	C9—C8—H8	121.2
C7—O2—C12	115.58 (17)	C5—C8—H8	121.2
N1—C1—N2	122.4 (2)	C8—C9—C10	121.3 (3)

N1—C1—S1	115.1 (2)	C8—C9—H9	119.4
N2—C1—S1	122.52 (17)	C10—C9—H9	119.4
C3—C2—N1	116.0 (3)	C11—C10—C9	120.7 (3)
C3—C2—H2	122.0	C11—C10—H10	119.7
N1—C2—H2	122.0	C9—C10—H10	119.7
C2—C3—S1	111.0 (2)	C6—C11—C10	118.3 (2)
C2—C3—H3	124.5	C6—C11—H11	120.8
S1—C3—H3	124.5	C10—C11—H11	120.8
O1—C4—N2	124.2 (3)	O2—C12—C13	108.4 (2)
O1—C4—C5	129.8 (2)	O2—C12—H12A	110.0
N2—C4—C5	106.1 (2)	C13—C12—H12A	110.0
C6—C5—C8	121.7 (3)	O2—C12—H12B	110.0
C6—C5—C4	109.0 (2)	C13—C12—H12B	110.0
C8—C5—C4	129.3 (2)	H12A—C12—H12B	108.4
C11—C6—C5	120.4 (2)	C12—C13—H13A	109.5
C11—C6—C7	128.8 (2)	C12—C13—H13B	109.5
C5—C6—C7	110.8 (2)	H13A—C13—H13B	109.5
O2—C7—N2	114.23 (19)	C12—C13—H13C	109.5
O2—C7—C6	115.0 (2)	H13A—C13—H13C	109.5
N2—C7—C6	100.76 (17)	H13B—C13—H13C	109.5
