

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-[(*E*)-(5-Bromo-2-hydroxyphenyl)-methylideneamino]benzenesulfonamideZahid H. Chohan,^a Hazoor A. Shad^a and M. Nawaz Tahir^{b*}^aDepartment of Chemistry, Bahauddin Zakariya University, Multan 60800, Pakistan, and ^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan
Correspondence e-mail: dmntahir_uos@yahoo.com


Received 6 September 2009; accepted 6 September 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.037; wR factor = 0.087; data-to-parameter ratio = 8.8.

In the title compound, $\text{C}_{13}\text{H}_{11}\text{ClN}_2\text{O}_3\text{S}$, the dihedral angle between the benzene rings is 12.26 (33) $^\circ$ and an intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond helps to establish the conformation. In the crystal, $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules.

Related literature

For a related structure and background discussion, see: Chohan *et al.* (2009). For graph-set theory, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_{13}\text{H}_{11}\text{BrN}_2\text{O}_3\text{S}$ $M_r = 355.21$ Monoclinic, $P2_1$ $a = 6.1224$ (15) Å $b = 4.5263$ (13) Å $c = 23.445$ (9) Å $\beta = 94.44$ (2) $^\circ$ $V = 647.8$ (3) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 3.34$ mm⁻¹ $T = 100$ K $0.22 \times 0.20 \times 0.16$ mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

 $T_{\min} = 0.482$, $T_{\max} = 0.587$

3181 measured reflections

1653 independent reflections

1458 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.042$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.087$ $S = 1.00$

1653 reflections

188 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\max} = 0.71$ e Å⁻³ $\Delta\rho_{\min} = -1.13$ e Å⁻³

Absolute structure: Flack (1983),

279 Friedel pairs

Flack parameter: 0.025 (16)

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.82	1.86	2.583 (7)	146
$\text{N2}-\text{H21}\cdots\text{O3}^{\text{i}}$	0.73 (8)	2.26 (7)	2.950 (7)	160 (8)
$\text{N2}-\text{H22}\cdots\text{O2}^{\text{ii}}$	0.92 (7)	2.58 (8)	3.325 (7)	139 (6)
$\text{C9}-\text{H9}\cdots\text{O2}^{\text{iii}}$	0.93	2.57	3.386 (7)	146

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + 1$; (ii) $x, y + 1, z$; (iii) $x + 1, y + 1, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

HAS gratefully acknowledges the Higher Education Commission, Islamabad, Pakistan, for providing him with a scholarship under the Indigenous PhD Program (PIN 042-160410-PS2-117).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5090).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chohan, Z. H., Shad, H. A. & Tahir, M. N. (2009). *Acta Cryst.* **E65**, o57.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o2426 [doi:10.1107/S1600536809036010]

4-[(*E*)-(5-Bromo-2-hydroxyphenyl)methylideneamino]benzenesulfonamide

Z. H. Chohan, H. A. Shad and M. N. Tahir

Comment

As part of our ongoing studies of sulfonamides, we now report the structure of the title compound, (I). The crystal structure of (II) 4-[(*E*)-(5-chloro-2-hydroxyphenyl) methylidene]amino}benzenesulfonamide (Chohan *et al.*, 2009) has been reported which differs from (I) due to chloro substitution instead of bromo.

In (I), the benzene rings A (C1—C6) of 5-bromosalicylaldehyde and B (C8—C13) of sulfanilamide are oriented at a dihedral angle of 12.03 (36)°. The Br1 and S1 atoms are at a distance of -0.016 (8) and -0.159 (9) Å from the mean square planes of rings A and B, respectively. There exist two intramolecular H-bonds (Table 1, Fig. 1) forming S(5) and S(6) ring motifs (Bernstein *et al.*, 1995). Three intermolecular H-bondings (Table 1) link the molecules in polymeric nature extending along the *b* axis (Fig. 2) and forming $R_3^3(10)$ and $R_2^1(8)$ ring motifs.

Experimental

Sulfanilamide (0.344 g, 2 mmol) in ethanol (20 ml) was mixed with 5-bromosalicylaldehyde (0.402 g, 2 mmol) in ethanol (10 ml). The resultant mixture was refluxed for 4 h by monitoring through TLC. During refluxing the solution turned from colorless to orange yellow. After completion of reaction, it was cooled to room temperature, filtered and volume reduced to about one-third using rotary evaporator. It was then allowed to stand for 7 days at room temperature. After which a crystallized product was formed that was filtered, washed with ethanol (2 × 5 ml), dried and recrystallized in a mixture of methanol and ethanol (1:1) to afford shiny orange yellow prisms of (I).

Refinement

The coordinates of H-atoms of the NH₂ group were refined. The other H-atoms were positioned geometrically with O—H = 0.82, C—H = 0.93 Å for aromatic H atoms and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C, N, O})$, where $x = 1.2$ for all H atoms.

Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radius. The dotted line indicate the intramolecular H-bond.




Fig. 2. The partial packing of (I) showing how molecules form polymeric chains extending along the crystallographic *b* axis and ring motifs exist.

4-[(*E*)-(5-Bromo-2-hydroxyphenyl)methylideneamino]benzenesulfonamide

Crystal data

$C_{13}H_{11}BrN_2O_3S$

$M_r = 355.21$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 6.1224$ (15) Å

$b = 4.5263$ (13) Å

$c = 23.445$ (9) Å

$\beta = 94.44$ (2)°

$V = 647.8$ (3) Å³

$Z = 2$

$F_{000} = 356$

$D_x = 1.821$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1653 reflections

$\theta = 3.3$ – 25.5 °

$\mu = 3.34$ mm⁻¹

$T = 100$ K

Prismatic, yellow

$0.22 \times 0.20 \times 0.16$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 7.80 pixels mm⁻¹

$T = 100$ K

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.482$, $T_{\max} = 0.587$

3181 measured reflections

1653 independent reflections

1458 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\max} = 25.5$ °

$\theta_{\min} = 3.3$ °

$h = -6 \rightarrow 7$

$k = -2 \rightarrow 5$

$l = -28 \rightarrow 28$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0554P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$wR(F^2) = 0.087$	$(\Delta/\sigma)_{\max} = 0.001$
$S = 1.00$	$\Delta\rho_{\max} = 0.71 \text{ e } \text{\AA}^{-3}$
1653 reflections	$\Delta\rho_{\min} = -1.13 \text{ e } \text{\AA}^{-3}$
188 parameters	Extinction coefficient: ?
1 restraint	Absolute structure: Flack (1983), 279 Friedal pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.025 (16)
Secondary atom site location: difference Fourier map	

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R - factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.22142 (8)	1.43584 (16)	0.03133 (2)	0.0184 (2)
S1	0.2382 (2)	-0.1521 (3)	0.42877 (6)	0.0131 (4)
O1	0.8682 (5)	0.9320 (15)	0.21201 (14)	0.0184 (11)
O2	0.0460 (7)	-0.3040 (10)	0.40583 (17)	0.0175 (12)
O3	0.4146 (6)	-0.3217 (10)	0.45633 (17)	0.0148 (12)
N1	0.5538 (8)	0.6166 (11)	0.2472 (2)	0.0141 (16)
N2	0.1639 (9)	0.0737 (13)	0.4770 (2)	0.0166 (17)
C1	0.5019 (8)	0.947 (2)	0.16824 (19)	0.0124 (14)
C2	0.7192 (9)	1.0429 (15)	0.1723 (2)	0.0134 (16)
C3	0.7829 (9)	1.2590 (14)	0.1347 (2)	0.0148 (17)
C4	0.6381 (9)	1.3763 (12)	0.0932 (2)	0.0143 (19)
C5	0.4222 (9)	1.2702 (14)	0.0883 (2)	0.0120 (17)
C6	0.3552 (10)	1.0646 (14)	0.1257 (2)	0.0149 (17)
C7	0.4239 (9)	0.7346 (14)	0.2085 (2)	0.0134 (17)
C8	0.4747 (8)	0.4182 (18)	0.2878 (2)	0.0114 (14)
C9	0.6126 (9)	0.3535 (13)	0.3358 (2)	0.0147 (19)
C10	0.5457 (10)	0.1737 (14)	0.3780 (3)	0.0155 (17)
C11	0.3363 (9)	0.0532 (14)	0.3724 (2)	0.0135 (17)
C12	0.1995 (9)	0.1113 (13)	0.3242 (2)	0.0134 (17)
C13	0.2664 (9)	0.2936 (14)	0.2818 (2)	0.0145 (17)
H1	0.81278	0.79686	0.22919	0.0218*
H3	0.92711	1.32519	0.13772	0.0176*
H4	0.68212	1.52308	0.06879	0.0172*
H6	0.21007	1.00207	0.12277	0.0178*
H7	0.27623	0.68391	0.20607	0.0155*

supplementary materials

H9	0.75293	0.43345	0.33940	0.0177*
H10	0.63936	0.13250	0.41016	0.0182*
H12	0.06076	0.02660	0.32024	0.0159*
H13	0.17312	0.33314	0.24953	0.0172*
H21	0.258 (12)	0.139 (18)	0.493 (3)	0.0197*
H22	0.062 (11)	0.200 (17)	0.460 (3)	0.0197*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0216 (3)	0.0200 (3)	0.0131 (3)	0.0009 (4)	-0.0012 (2)	0.0024 (3)
S1	0.0143 (7)	0.0109 (8)	0.0142 (7)	-0.0014 (6)	0.0010 (5)	0.0006 (5)
O1	0.0173 (18)	0.019 (2)	0.0186 (18)	-0.009 (3)	0.0003 (14)	0.010 (3)
O2	0.017 (2)	0.017 (2)	0.018 (2)	-0.006 (2)	-0.0015 (17)	0.0027 (19)
O3	0.014 (2)	0.013 (2)	0.017 (2)	0.0027 (18)	-0.0014 (17)	0.0053 (17)
N1	0.013 (2)	0.013 (3)	0.017 (3)	-0.002 (2)	0.005 (2)	-0.001 (2)
N2	0.018 (3)	0.010 (3)	0.022 (3)	-0.001 (3)	0.003 (2)	-0.001 (2)
C1	0.015 (2)	0.010 (3)	0.013 (2)	0.005 (4)	0.0056 (19)	-0.006 (4)
C2	0.017 (3)	0.018 (3)	0.005 (2)	-0.001 (3)	0.000 (2)	-0.003 (2)
C3	0.011 (3)	0.011 (3)	0.023 (3)	-0.002 (3)	0.006 (2)	0.004 (3)
C4	0.020 (3)	0.008 (4)	0.016 (3)	-0.005 (3)	0.009 (2)	0.001 (2)
C5	0.016 (3)	0.010 (3)	0.010 (3)	0.005 (3)	0.002 (2)	-0.006 (2)
C6	0.015 (3)	0.015 (3)	0.015 (3)	-0.001 (3)	0.003 (2)	-0.004 (3)
C7	0.013 (3)	0.008 (3)	0.020 (3)	-0.004 (3)	0.007 (2)	-0.002 (3)
C8	0.017 (2)	0.004 (3)	0.014 (2)	0.002 (3)	0.006 (2)	0.001 (3)
C9	0.009 (3)	0.010 (4)	0.025 (3)	-0.002 (2)	0.001 (2)	-0.002 (2)
C10	0.017 (3)	0.016 (3)	0.013 (3)	0.001 (3)	-0.002 (2)	0.003 (3)
C11	0.017 (3)	0.008 (3)	0.016 (3)	0.001 (3)	0.005 (2)	0.001 (2)
C12	0.012 (3)	0.014 (3)	0.014 (3)	-0.004 (3)	0.000 (2)	-0.001 (3)
C13	0.016 (3)	0.013 (3)	0.014 (3)	0.002 (3)	-0.002 (2)	-0.002 (3)

Geometric parameters (\AA , $^\circ$)

Br1—C5	1.898 (5)	C4—C5	1.403 (8)
S1—O2	1.431 (5)	C5—C6	1.363 (8)
S1—O3	1.437 (4)	C8—C13	1.392 (8)
S1—N2	1.616 (5)	C8—C9	1.385 (7)
S1—C11	1.759 (6)	C9—C10	1.368 (9)
O1—C2	1.349 (7)	C10—C11	1.390 (8)
O1—H1	0.8200	C11—C12	1.379 (7)
N1—C8	1.421 (8)	C12—C13	1.378 (8)
N1—C7	1.276 (7)	C3—H3	0.9300
N2—H21	0.73 (8)	C4—H4	0.9300
N2—H22	0.92 (7)	C6—H6	0.9300
C1—C7	1.454 (9)	C7—H7	0.9300
C1—C6	1.395 (8)	C9—H9	0.9300
C1—C2	1.396 (8)	C10—H10	0.9300
C2—C3	1.393 (8)	C12—H12	0.9300
C3—C4	1.372 (7)	C13—H13	0.9300

O2—S1—O3	118.7 (3)	C9—C8—C13	119.5 (5)
O2—S1—N2	107.4 (3)	N1—C8—C9	117.4 (5)
O2—S1—C11	106.9 (3)	C8—C9—C10	121.0 (5)
O3—S1—N2	105.4 (3)	C9—C10—C11	119.4 (6)
O3—S1—C11	109.4 (3)	S1—C11—C10	120.2 (4)
N2—S1—C11	108.8 (3)	C10—C11—C12	120.0 (5)
C2—O1—H1	109.00	S1—C11—C12	119.7 (4)
C7—N1—C8	121.0 (5)	C11—C12—C13	120.6 (5)
S1—N2—H22	108 (5)	C8—C13—C12	119.4 (5)
S1—N2—H21	111 (6)	C2—C3—H3	119.00
H21—N2—H22	117 (8)	C4—C3—H3	119.00
C2—C1—C6	119.2 (6)	C3—C4—H4	121.00
C2—C1—C7	121.4 (5)	C5—C4—H4	121.00
C6—C1—C7	119.4 (5)	C1—C6—H6	120.00
O1—C2—C1	121.4 (5)	C5—C6—H6	120.00
C1—C2—C3	119.2 (5)	N1—C7—H7	119.00
O1—C2—C3	119.4 (5)	C1—C7—H7	119.00
C2—C3—C4	121.6 (5)	C8—C9—H9	119.00
C3—C4—C5	118.6 (5)	C10—C9—H9	119.00
C4—C5—C6	120.6 (5)	C9—C10—H10	120.00
Br1—C5—C4	118.5 (4)	C11—C10—H10	120.00
Br1—C5—C6	120.8 (4)	C11—C12—H12	120.00
C1—C6—C5	120.7 (6)	C13—C12—H12	120.00
N1—C7—C1	121.4 (5)	C8—C13—H13	120.00
N1—C8—C13	123.1 (5)	C12—C13—H13	120.00
O2—S1—C11—C10	-166.2 (5)	O1—C2—C3—C4	179.3 (5)
O2—S1—C11—C12	18.2 (6)	C1—C2—C3—C4	-1.1 (9)
O3—S1—C11—C10	-36.6 (6)	C2—C3—C4—C5	-1.2 (8)
O3—S1—C11—C12	147.9 (5)	C3—C4—C5—Br1	179.4 (4)
N2—S1—C11—C10	78.1 (6)	C3—C4—C5—C6	3.0 (8)
N2—S1—C11—C12	-97.5 (5)	Br1—C5—C6—C1	-178.8 (5)
C8—N1—C7—C1	177.5 (6)	C4—C5—C6—C1	-2.5 (9)
C7—N1—C8—C9	-165.3 (6)	N1—C8—C9—C10	177.4 (6)
C7—N1—C8—C13	13.4 (10)	C13—C8—C9—C10	-1.3 (10)
C6—C1—C2—O1	-178.8 (6)	N1—C8—C13—C12	-177.7 (6)
C6—C1—C2—C3	1.6 (10)	C9—C8—C13—C12	1.0 (9)
C7—C1—C2—O1	3.2 (10)	C8—C9—C10—C11	0.3 (9)
C7—C1—C2—C3	-176.4 (6)	C9—C10—C11—S1	-174.5 (5)
C2—C1—C6—C5	0.2 (10)	C9—C10—C11—C12	1.1 (9)
C7—C1—C6—C5	178.2 (6)	S1—C11—C12—C13	174.2 (5)
C2—C1—C7—N1	-3.7 (10)	C10—C11—C12—C13	-1.4 (9)
C6—C1—C7—N1	178.3 (6)	C11—C12—C13—C8	0.4 (9)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots N1	0.82	1.86	2.583 (7)	146
N2—H21 \cdots O3 ⁱ	0.73 (8)	2.26 (7)	2.950 (7)	160 (8)

supplementary materials

N2—H22...O2 ⁱⁱ	0.92 (7)	2.58 (8)	3.325 (7)	139 (6)
C9—H9...O2 ⁱⁱⁱ	0.93	2.57	3.386 (7)	146

Symmetry codes: (i) $-x+1, y+1/2, -z+1$; (ii) $x, y+1, z$; (iii) $x+1, y+1, z$.

Fig. 1




Fig. 2

