

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4-Acetylpyridinium hydrogen sulfate

Xue-qun Fu

Ordered Matter Science Research Center, Southeast University, Nanjing 210096, People's Republic of China

Correspondence e-mail: fuxuequn222@163.com


Received 2 July 2009; accepted 2 September 2009

 Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.053;  $wR$  factor = 0.170; data-to-parameter ratio = 16.7.

The crystal structure of the title compound,  $\text{C}_7\text{H}_8\text{NO}^+\cdot\text{HSO}_4^-$ , consists of  $\text{O}-\text{H}\cdots\text{O}$  hydrogen-bonded extended chains of hydrogen sulfate anions. Each hydrogen sulfate anion is furthermore connected to one 4-acetylpyridinium cation *via* a hydrogen bond of the  $\text{N}-\text{H}\cdots\text{O}$  type.

## Related literature

For the synthesis of 4-acetylpyridine, see: Piner *et al.* (1934). For the crystal structure of an adduct of 4-acetylpyridine with pentachlorophenol, see: Majerz *et al.* (1991). For the crystal structures of Zn and Ni complexes of 4-acetylpyridine, see: Pang *et al.* (1994); Steffen *et al.* (1977).



## Experimental

Crystal data

 $\text{C}_7\text{H}_8\text{NO}^+\cdot\text{HSO}_4^-$ 
 $M_r = 219.21$ 

 Orthorhombic,  $P2_12_12_1$ 
 $a = 4.6454$  (9) Å

 $b = 9.597$  (2) Å

 $c = 21.310$  (4) Å

 $V = 950.1$  (3) Å<sup>3</sup>
 $Z = 4$ 

 Mo  $K\alpha$  radiation

 $\mu = 0.34$  mm<sup>-1</sup>
 $T = 298$  K

 $0.20 \times 0.20 \times 0.20$  mm

## Data collection

Rigaku SCXmini diffractometer

Absorption correction: multi-scan

 (*CrystalClear*; Rigaku, 2005)

 $T_{\min} = 0.935$ ,  $T_{\max} = 0.935$ 

9843 measured reflections

2156 independent reflections

 1622 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.077$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.053$ 
 $wR(F^2) = 0.170$ 
 $S = 0.93$ 

2156 reflections

129 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.38$  e Å<sup>-3</sup>
 $\Delta\rho_{\min} = -0.19$  e Å<sup>-3</sup>

Absolute structure: Flack (1983),

860 Friedel pairs

Flack parameter: 0.2 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1B}\cdots\text{O1}^{\text{i}}$	0.86	1.92	2.772 (5)	172
$\text{O3}-\text{H3}\cdots\text{O2}^{\text{ii}}$	0.82	1.76	2.565 (5)	166

 Symmetry codes: (i)  $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x + 1, y, z$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PRPKAPPA* (Ferguson, 1999).

The authors are grateful to the starter fund of Southeast University for financial support to buy the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2126).

## References

- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Majerz, I., Malarski, Z. & Sawka-Dobrowolska, W. (1991). *J. Mol. Struct.* **249**, 109–116.  
 Pang, L., Whitehead, M. A., Bermardinelli, G. & Lucken, E. A. C. (1994). *J. Chem. Crystallogr.* **24**, 203–211.  
 Piner, R. (1934). *Ber. Deutsch Chem. Ges.* **B34**, 4250–4251.  
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Steffen, W. L. & Palenik, G. J. (1977). *Inorg. Chem.* **16**, 1119–1128.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2385 [ doi:10.1107/S1600536809035417 ]

## 4-Acetylpyridinium hydrogen sulfate

X. Fu

### Comment

4-Acetylpyridine may be used as a ligand in coordination compounds *e.g.* with Zn (Steffen & Palenik, 1977) or Ni (Pang *et al.*, 1994). The crystal structure of 4-acetylpyridine together with pentachlorophenol is also known (Majerz *et al.*, 1991).

The asymmetric unit of the title compound contains one 4-acetylpyridinium cation and one hydrogen sulfate anion (Fig 1). In the anion, the bond length of S1—O3 is 1.553 (6) Å compared to the average bond length of 1.438 (5) Å of the other S1—O bonds. It is therefore reasonable that the hydrogen atom of hydrogen sulfate is bonded to O3. The supramolecular structure consists of infinite chains of anions with one cation linked to each anion *via* an additional hydrogen bond (N1—H1B···O1 2.774 (8) Å, Fig 2).

### Experimental

4-Acetylpyridine was obtained according to the method described by Piner (1934). Reaction of equimolar amounts of 4-acetylpyridine and H<sub>2</sub>SO<sub>4</sub> produced a precipitate. This was filtered off, dried and dissolved in 96% ethanol from which single crystals were grown by slow evaporation of the solvent at room temperature.

### Refinement

The H atom connected to O3 was discernible from difference electron-density map. Nevertheless, it was placed to the ideal position, with S1—O3—H angle tetrahedral, allowing the H atom to ride on the immediately preceding atom O3 and rotate about the S1—O3 bond, refined in a riding atom approximation with a constrained bond length of O—H = 0.82 Å. Positional parameters of the other H atoms were calculated geometrically with C<sub>ar</sub>—H = 0.93 Å and C<sub>Me</sub>—H = 0.96 Å and were allowed to ride on the corresponding C atoms with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ . In the absence of significant anomalous dispersion effects, 860 Friedel pairs were merged.

### Figures




Fig. 1. Molecular structure of the title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and all H atoms have been omitted for clarity.




Fig. 2. Crystal packing of the title compound, stacking along the *b* axis. Dashed lines indicate hydrogen bonds.

## 4-Acetylpyridinium hydrogen sulfate

### Crystal data

$C_7H_8NO^+ \cdot HSO_4^-$

$M_r = 219.21$

Orthorhombic,  $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 4.6454$  (9) Å

$b = 9.597$  (2) Å

$c = 21.310$  (4) Å

$V = 950.1$  (3) Å<sup>3</sup>

$Z = 4$

$F_{000} = 456$

$D_x = 1.533$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4231 reflections

$\theta = 3.6$ – $27.6^\circ$

$\mu = 0.34$  mm<sup>-1</sup>

$T = 298$  K

Prism, colourless

$0.20 \times 0.20 \times 0.20$  mm

### Data collection

Rigaku SCXmini  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 13.6612 pixels mm<sup>-1</sup>

$T = 298$  K

$\omega$  scans

Absorption correction: multi-scan  
(CrystalClear; Rigaku, 2005)

$T_{\min} = 0.935$ ,  $T_{\max} = 0.935$

9843 measured reflections

2156 independent reflections

1622 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.077$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 3.6^\circ$

$h = -6 \rightarrow 6$

$k = -12 \rightarrow 12$

$l = -27 \rightarrow 27$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.053$

$wR(F^2) = 0.170$

$S = 0.93$

2156 reflections

129 parameters

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1249P)^2 + 1.8027P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.38$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

Extinction correction: none

Primary atom site location: structure-invariant direct methods Absolute structure: Flack (1983), 860 Friedel pairs  
 Secondary atom site location: difference Fourier map Flack parameter: 0.2 (2)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.9101 (2)	0.18577 (10)	0.16383 (5)	0.0475 (3)
O5	-0.0805 (8)	0.6665 (5)	0.03394 (17)	0.0853 (12)
O1	1.0236 (7)	0.3041 (3)	0.19787 (16)	0.0655 (9)
N1	0.6303 (8)	0.7352 (4)	0.20080 (17)	0.0573 (9)
H1B	0.7336	0.7648	0.2315	0.069*
C3	0.3084 (8)	0.6431 (4)	0.10302 (18)	0.0457 (9)
C1	0.4610 (10)	0.8250 (5)	0.1713 (2)	0.0614 (11)
H1A	0.4549	0.9175	0.1842	0.074*
C6	0.1250 (13)	0.5981 (6)	0.0476 (2)	0.0634 (13)
C5	0.6477 (10)	0.6005 (5)	0.1849 (2)	0.0548 (11)
H5A	0.7674	0.5403	0.2070	0.066*
C4	0.4852 (9)	0.5528 (5)	0.1353 (2)	0.0544 (11)
H4A	0.4949	0.4596	0.1236	0.065*
C2	0.2960 (10)	0.7819 (4)	0.1223 (2)	0.0546 (10)
H2A	0.1759	0.8446	0.1017	0.066*
O2	0.6532 (7)	0.2155 (4)	0.1291 (2)	0.0858 (12)
O3	1.1265 (8)	0.1454 (6)	0.1117 (2)	0.0968 (15)
H3	1.2838	0.1802	0.1194	0.145*
O4	0.8942 (13)	0.0646 (4)	0.2026 (2)	0.1068 (15)
C7	0.2139 (18)	0.4694 (7)	0.0132 (3)	0.106 (2)
H7A	0.0795	0.4510	-0.0200	0.160*
H7B	0.2166	0.3919	0.0417	0.160*
H7C	0.4026	0.4824	-0.0042	0.160*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0356 (4)	0.0481 (5)	0.0588 (5)	-0.0017 (4)	-0.0011 (4)	-0.0066 (5)
O5	0.058 (2)	0.121 (3)	0.077 (2)	-0.004 (3)	-0.0169 (19)	0.008 (2)
O1	0.069 (2)	0.0471 (16)	0.080 (2)	0.0051 (15)	-0.0109 (17)	-0.0183 (15)

## supplementary materials

---

N1	0.046 (2)	0.073 (2)	0.0531 (19)	-0.0074 (18)	0.0000 (17)	-0.0004 (17)
C3	0.0372 (18)	0.054 (2)	0.0459 (19)	-0.0021 (16)	0.0076 (16)	0.0062 (17)
C1	0.063 (3)	0.049 (2)	0.072 (3)	-0.003 (2)	0.002 (2)	0.000 (2)
C6	0.059 (3)	0.083 (3)	0.048 (2)	-0.016 (3)	0.009 (2)	0.003 (2)
C5	0.044 (2)	0.066 (3)	0.054 (2)	0.007 (2)	-0.0010 (19)	0.0119 (19)
C4	0.060 (3)	0.045 (2)	0.058 (2)	0.0030 (19)	0.014 (2)	0.0029 (18)
C2	0.055 (2)	0.047 (2)	0.062 (2)	0.0078 (19)	0.001 (2)	0.0075 (19)
O2	0.0428 (18)	0.096 (3)	0.118 (3)	0.0099 (17)	-0.021 (2)	-0.021 (2)
O3	0.0471 (19)	0.145 (4)	0.098 (3)	-0.010 (2)	0.001 (2)	-0.060 (3)
O4	0.137 (4)	0.073 (2)	0.110 (3)	-0.039 (3)	-0.028 (3)	0.022 (2)
C7	0.128 (6)	0.110 (5)	0.081 (4)	0.002 (5)	-0.020 (4)	-0.039 (4)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

S1—O4	1.429 (4)	C1—C2	1.360 (6)
S1—O2	1.433 (4)	C1—H1A	0.9300
S1—O1	1.447 (3)	C6—C7	1.495 (8)
S1—O3	1.547 (4)	C5—C4	1.378 (6)
O5—C6	1.194 (7)	C5—H5A	0.9300
N1—C1	1.325 (6)	C4—H4A	0.9300
N1—C5	1.339 (6)	C2—H2A	0.9300
N1—H1B	0.8600	O3—H3	0.8200
C3—C4	1.378 (6)	C7—H7A	0.9600
C3—C2	1.395 (6)	C7—H7B	0.9600
C3—C6	1.519 (6)	C7—H7C	0.9600
O4—S1—O2	114.7 (3)	C7—C6—C3	117.5 (5)
O4—S1—O1	111.6 (2)	N1—C5—C4	118.8 (4)
O2—S1—O1	113.9 (2)	N1—C5—H5A	120.6
O4—S1—O3	104.2 (3)	C4—C5—H5A	120.6
O2—S1—O3	102.7 (2)	C5—C4—C3	120.0 (4)
O1—S1—O3	108.7 (2)	C5—C4—H4A	120.0
C1—N1—C5	122.9 (4)	C3—C4—H4A	120.0
C1—N1—H1B	118.5	C1—C2—C3	119.5 (4)
C5—N1—H1B	118.5	C1—C2—H2A	120.2
C4—C3—C2	118.6 (4)	C3—C2—H2A	120.2
C4—C3—C6	123.0 (4)	S1—O3—H3	109.5
C2—C3—C6	118.5 (4)	C6—C7—H7A	109.5
N1—C1—C2	120.1 (4)	C6—C7—H7B	109.5
N1—C1—H1A	120.0	H7A—C7—H7B	109.5
C2—C1—H1A	120.0	C6—C7—H7C	109.5
O5—C6—C7	123.8 (5)	H7A—C7—H7C	109.5
O5—C6—C3	118.8 (5)	H7B—C7—H7C	109.5
C5—N1—C1—C2	-0.4 (7)	N1—C5—C4—C3	-0.1 (6)
C4—C3—C6—O5	157.9 (4)	C2—C3—C4—C5	-0.8 (6)
C2—C3—C6—O5	-21.8 (6)	C6—C3—C4—C5	179.5 (4)
C4—C3—C6—C7	-21.9 (6)	N1—C1—C2—C3	-0.5 (7)
C2—C3—C6—C7	158.4 (5)	C4—C3—C2—C1	1.1 (6)
C1—N1—C5—C4	0.7 (7)	C6—C3—C2—C1	-179.2 (4)

*Hydrogen-bond geometry* (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1B···O1 <sup>i</sup>	0.86	1.92	2.772 (5)	172
O3—H3···O2 <sup>ii</sup>	0.82	1.76	2.565 (5)	166

Symmetry codes: (i)  $-x+2, y+1/2, -z+1/2$ ; (ii)  $x+1, y, z$ .

Fig. 1




Fig. 2

