

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(4-Bromophenyl)-3-(2-thienyl-carbonyl)thiourea

Sohail Saeed,^{a*} Naghmana Rashid,^a Rizwan Hussain^b and Peter G. Jones^c

^aDepartment of Chemistry, Research Complex, Allama Iqbal Open University, Islamabad, Pakistan, ^bNational Engineering and Scientific Commission, PO Box 2801, Islamabad, Pakistan, and ^cInstitut für Anorganische und Analytische Chemie, Technische Universität Braunschweig, Postfach 3329, 38023 Braunschweig, Germany

Correspondence e-mail: sohai262001@yahoo.com


Received 28 August 2009; accepted 23 September 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.025; wR factor = 0.067; data-to-parameter ratio = 15.9.

The title compound, $\text{C}_{12}\text{H}_9\text{BrN}_2\text{OS}_2$, consists of two planar parts, *viz.* the thiophene ring including all substituents (r.m.s. deviation 0.007 Å) and the benzene ring including the respective substituents as well as the thione group (r.m.s. deviation 0.05 Å). The interplanar angle is 18.84 (6)°. An intramolecular $\text{C}_{\text{phenyl}}-\text{N}-\text{H}\cdots\text{OC}$ hydrogen bond is observed. The three-dimensional packing involves three types of interactions, *viz.* $\text{N}-\text{H}\cdots\text{S}$, $\text{C}-\text{H}\cdots\text{S}$ ($\times 2$) and $\text{Br}\cdots\text{S}$ [3.6924 (6) Å].

Related literature

For general background to the chemistry of thiourea derivatives, see: Choi *et al.* (2008); Jones *et al.* (2008); Su *et al.* (2006). For related structures, see: Saeed *et al.* (2008a,b,c); Yunus *et al.* (2008). For the cytotoxicity and genotoxicity of anticancer drugs to normal cells in cancer therapy, see: Aydemir & Bilaloglu (2003).



Experimental

Crystal data

$\text{C}_{12}\text{H}_9\text{BrN}_2\text{OS}_2$	$V = 1319.4$ (1) Å ³
$M_r = 341.24$	$Z = 4$
Monoclinic, $P2_1/n$	Cu $K\alpha$ radiation
$a = 13.1483$ (6) Å	$\mu = 7.12$ mm ⁻¹
$b = 4.4263$ (2) Å	$T = 100$ K
$c = 22.671$ (1) Å	$0.15 \times 0.05 \times 0.02$ mm
$\beta = 90.412$ (5)°	

Data collection

Oxford Diffraction Xcalibur Nova A diffractometer	20285 measured reflections
Absorption correction: multi-scan (<i>CrysAlis Pro</i> ; Oxford Diffraction, 2009)	2712 independent reflections
$T_{\text{min}} = 0.558$, $T_{\text{max}} = 1.000$	2438 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.067$	$\Delta\rho_{\text{max}} = 0.42$ e Å ⁻³
$S = 1.06$	$\Delta\rho_{\text{min}} = -0.36$ e Å ⁻³
2712 reflections	
171 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H01}\cdots\text{S2}^i$	0.85 (3)	2.74 (3)	3.5625 (16)	163 (2)
$\text{N2}-\text{H02}\cdots\text{O}$	0.85 (3)	1.89 (3)	2.624 (2)	144 (2)
$\text{C9}-\text{H9}\cdots\text{S1}^{ii}$	0.95	2.89	3.704 (2)	144
$\text{C2}-\text{H2}\cdots\text{S2}^i$	0.95	2.76	3.3193 (18)	119

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x + \frac{1}{2}, y + \frac{3}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis Pro* (Oxford Diffraction, 2009); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis Pro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Siemens, 1994); software used to prepare material for publication: *SHELXL97*.

The authors are grateful to Allama Iqbal Open University and the National Engineering & Scientific Commission, Islamabad, Pakistan, for the allocation of research and analytical laboratory facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2139).

References

- Aydemir, N. & Bilaloglu, R. (2003). *Mutat. Res.* **537**, 43–51.
 Choi, M. K., Kim, H. N., Choi, H. J., Yoon, J. & Hyun, M. H. (2008). *Tetrahedron Lett.* **49**, 4522–4525.
 Jones, C. E., Turega, S. M., Clarke, M. L. & Philp, D. (2008). *Tetrahedron Lett.* **49**, 4666–4669.
 Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, England.
 Saeed, S., Bhatti, M. H., Tahir, M. K. & Jones, P. G. (2008a). *Acta Cryst.* **E64**, o1369.
 Saeed, S., Bhatti, M. H., Yunus, U. & Jones, P. G. (2008b). *Acta Cryst.* **E64**, o1485.
 Saeed, S., Bhatti, M. H., Yunus, U. & Jones, P. G. (2008c). *Acta Cryst.* **E64**, o1566.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Siemens (1994). *XP*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Su, B.-Q., Liu, G.-L., Sheng, L., Wang, X.-Q. & Xian, L. (2006). *Phosphorus Sulfur Silicon*, **181**, 745–750.

Yunus, U., Tahir, M. K., Bhatti, M. H., Ali, S. & Wong, W.-Y. (2008). *Acta Cryst.* **E64**, o20.

supporting information

Acta Cryst. (2009). E65, o2568–o2569 [doi:10.1107/S1600536809038537]

1-(4-Bromophenyl)-3-(2-thienylcarbonyl)thiourea**Sohail Saeed, Naghmana Rashid, Rizwan Hussain and Peter G. Jones****S1. Comment**

The development of new antimicrobial and anticancer therapeutic agents is one of the fundamental goals in medicinal chemistry. Cytotoxicity and genotoxicity of anticancer drugs to normal cells are major problems in cancer therapy and engender the risk of inducing secondary malignancy (Aydemir *et al.*, 2003). A dose of an anticancer drug sufficient to kill tumor cells is often toxic to the normal tissue and leads to many side effects, which, in turn, limit the efficacy of treatment. In recent years, there has been a concerted search for novel selective antitumor agents that lack many of the unpleasant side effects of conventional agents. Thiourea and its derivatives have found extensive applications in the field of medicine, agriculture and analytical chemistry. They are known to exhibit a wide variety of biological activities such as antiviral, anti-bacterial, antifungal, anticancer, antitubercular, herbicidal and insecticidal effects, and also constitute some epoxy resin curing agents containing amino functional groups (Saeed *et al.*, 2008*a,b,c*). They have found broad areas of application *e.g.* in anion recognition, nonlinear optics and catalysis, and also display good coordination ability (Choi *et al.*, 2008; Jones *et al.*, 2008; Su *et al.*, 2006). As part of our research on thiourea coordination chemistry, we are interested in the study of the influence of non-covalent interactions, especially hydrogen bonds and π - π stacking interactions, on the coordination modes of benzothiazoles bearing the 4-nitrobenzoylthiourea group with transition metal ions. Such coordination compounds of thiourea have been studied for various biological systems in terms of their antibacterial, antifungal and anticancer activities (Yunus *et al.*, 2008). The importance of such work lies in the possibility that the next generation of thiourea derivatives might be more efficacious as antimicrobial and anticancer agents. However, a thorough investigation of their structure, activity and stability under biological conditions is required. These detailed investigations could be helpful in designing more potent antimicrobial and anticancer agents for therapeutic use. The condensation of acyl/aroil thiocyanates with primary amines affords 1,3-disubstituted thioureas in excellent yields in a single step. In the present paper, the crystal structure of the title compound is reported.

The molecule (Fig. 1) consists of two planar parts: the thiophene ring plus C5 (r.m.s. deviation 0.007 Å) and the phenyl ring plus Br,N2,C6,S2 (0.05 Å), which subtend an interplanar angle of 18.84 (6)°. An intramolecular hydrogen bond N2—H02···O is observed.

The molecular packing is determined by four intermolecular contacts, each of which involves one or other of the sulfur atoms: a surprisingly long classical H bond N1—H01···S2, two weak C—H···S interactions (Table 1) and an interaction C1—S1···Br—C10 with S1···Br 3.6924 (6) Å, C1—S1···Br 165.89 (7)° and S1···Br—C10 90.51 (6)° (operator for Br and C10: $-x + 1/2, y - 3/2, -z + 1/2$). The combined effect is to create a three-dimensional pattern, a small part of which is shown in Fig. 2.

S2. Experimental

A mixture of ammonium thiocyanate (26 mmol) and 2-thiophene carbonyl chloride (26 mmol) in anhydrous acetone (60 ml) was stirred for 45 min. 2-Bromoaniline (26 mmol) was added and the reaction mixture was refluxed for 2 h. After

cooling, the reaction mixture was poured into acidified cold water. The resulting dark yellow solid was filtered and washed with cold acetone. The title compound (I) was obtained as colourless needles and laths of several mm length by recrystallization of the solid from ethyl acetate. These tended to split lengthwise when cut, but eventually a fragment suitable for X-ray structure analysis was found.

S3. Refinement

NH H atoms were refined freely. Other H atoms were placed in calculated positions and refined using a riding model with C—H 0.95 Å; These hydrogen U values were fixed at $1.2 \times U(\text{eq})$ of the parent atom. Data are 99.4% complete to 2θ 145°.




Figure 1

Thermal ellipsoid plot (50% probability level) of the title compound.




Figure 2

Packing diagram showing the four independent contacts (dashed bonds) N—H \cdots S, C—H \cdots S ($\times 2$), S \cdots Br (see text).

1-(4-Bromophenyl)-3-(2-thienylcarbonyl)thiourea

Crystal data

$C_{12}H_9BrN_2OS_2$
 $M_r = 341.24$
 Monoclinic, $P2_1/n$
 Hall symbol: -P 2yn
 $a = 13.1483$ (6) Å
 $b = 4.4263$ (2) Å
 $c = 22.671$ (1) Å
 $\beta = 90.412$ (5) $^\circ$
 $V = 1319.4$ (1) Å 3
 $Z = 4$

$F(000) = 680$
 $D_x = 1.718$ Mg m $^{-3}$
 Melting point: 389 K
 Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å
 Cell parameters from 12273 reflections
 $\theta = 3.4\text{--}75.7^\circ$
 $\mu = 7.12$ mm $^{-1}$
 $T = 100$ K
 Needle, colourless
 $0.15 \times 0.05 \times 0.02$ mm

Data collection

Oxford Diffraction Xcalibur Nova A
 diffractometer
 Radiation source: Nova (Cu) X-ray Source
 Mirror monochromator
 Detector resolution: 10.3543 pixels mm $^{-1}$
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Oxford Diffraction, 2009)
 $T_{\min} = 0.558$, $T_{\max} = 1.000$

20285 measured reflections
 2712 independent reflections
 2438 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$
 $\theta_{\max} = 75.9^\circ$, $\theta_{\min} = 3.9^\circ$
 $h = -16 \rightarrow 16$
 $k = -5 \rightarrow 4$
 $l = -28 \rightarrow 27$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.025$ $wR(F^2) = 0.067$ $S = 1.06$

2712 reflections

171 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0399P)^2 + 0.686P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.002$ $\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Short contact:

3.6924 (0.0006) S1 - Br_\$3 165.89 (0.07) C1 - S1 - Br_\$3 90.51 (0.06) S1 - Br_\$3 - C10_\$3 Operator for generating equivalent atoms: \$3 - $x + 1/2, y - 3/2, -z + 1/2$ Least-squares planes (x, y, z in crystal coordinates) and deviations from them (* indicates atom used to define plane)
- 2.2923 (0.0090) $x + 3.4843 (0.0018) y + 13.4394 (0.0126) z = 3.7714 (0.0071)$ * -0.0064 (0.0017) C1 * -0.0022 (0.0013) C2 * -0.0022 (0.0014) C3 * 0.0106 (0.0014) C4 * -0.0078 (0.0010) S1 *
0.0079 (0.0011) C5

Rms deviation of fitted atoms = 0.0069

- 5.4364 (0.0073) $x + 3.6780 (0.0010) y + 8.5062 (0.0060) z = 1.0213 (0.0056)$

Angle to previous plane (with approximate e.s.d.) = 18.84 (0.06)

* -0.0768 (0.0015) C6 * 0.0657 (0.0011) S2 * 0.0104 (0.0018) C7 * 0.0477 (0.0017) C8 * 0.0508 (0.0017) C9 * 0.0153
(0.0017) C10 * 0.0055 (0.0018) C11 * 0.0033 (0.0020) C12 * -0.0663 (0.0010) Br * -0.0558 (0.0015) N2 - 0.2202
(0.0028) C5 - 0.1410 (0.0028) O - 0.2122 (0.0021) N1

Rms deviation of fitted atoms = 0.0479

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.629471 (15)	0.87392 (5)	0.136702 (8)	0.02773 (9)
S1	0.12627 (4)	-0.47575 (13)	0.42493 (2)	0.03066 (13)
S2	0.57481 (4)	0.15778 (13)	0.42693 (2)	0.02881 (13)
O	0.27346 (11)	-0.1628 (4)	0.34866 (6)	0.0306 (3)
N1	0.40397 (12)	-0.1458 (4)	0.41634 (7)	0.0213 (3)
H01	0.4194 (19)	-0.180 (6)	0.4521 (12)	0.031 (7)*
N2	0.44284 (13)	0.1420 (4)	0.33512 (7)	0.0238 (4)
H02	0.385 (2)	0.075 (6)	0.3245 (11)	0.032 (7)*
C1	0.25426 (14)	-0.4373 (5)	0.43690 (8)	0.0228 (4)
C2	0.28750 (14)	-0.6036 (4)	0.48600 (8)	0.0191 (4)
H2	0.3558	-0.6113	0.4998	0.023*
C3	0.20512 (17)	-0.7586 (5)	0.51212 (9)	0.0303 (4)

H3	0.2122	-0.8857	0.5457	0.036*
C4	0.11538 (17)	-0.7072 (5)	0.48445 (11)	0.0340 (5)
H4	0.0527	-0.7916	0.4969	0.041*
C5	0.31026 (14)	-0.2406 (5)	0.39652 (8)	0.0226 (4)
C6	0.47172 (14)	0.0541 (5)	0.38914 (8)	0.0215 (4)
C7	0.49109 (15)	0.3288 (4)	0.29299 (8)	0.0213 (4)
C8	0.43466 (15)	0.3754 (5)	0.24114 (9)	0.0251 (4)
H8	0.3683	0.2922	0.2375	0.030*
C9	0.47486 (15)	0.5415 (5)	0.19539 (9)	0.0268 (4)
H9	0.4365	0.5732	0.1603	0.032*
C10	0.57114 (15)	0.6608 (4)	0.20115 (8)	0.0228 (4)
C11	0.62707 (15)	0.6225 (5)	0.25232 (9)	0.0279 (4)
H11	0.6928	0.7100	0.2559	0.033*
C12	0.58713 (15)	0.4557 (5)	0.29867 (9)	0.0293 (4)
H12	0.6254	0.4288	0.3340	0.035*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.02733 (13)	0.03447 (14)	0.02145 (12)	0.00217 (8)	0.00351 (8)	0.00748 (8)
S1	0.0204 (2)	0.0323 (3)	0.0393 (3)	-0.0031 (2)	-0.00143 (19)	-0.0057 (2)
S2	0.0253 (2)	0.0430 (3)	0.0181 (2)	-0.0118 (2)	-0.00757 (18)	0.00732 (19)
O	0.0251 (7)	0.0443 (9)	0.0222 (7)	-0.0074 (6)	-0.0074 (6)	0.0054 (6)
N1	0.0211 (8)	0.0283 (9)	0.0145 (8)	-0.0019 (6)	-0.0039 (6)	0.0029 (6)
N2	0.0208 (8)	0.0319 (9)	0.0187 (8)	-0.0042 (7)	-0.0045 (6)	0.0038 (7)
C1	0.0193 (8)	0.0278 (10)	0.0212 (9)	-0.0023 (8)	0.0005 (7)	-0.0055 (8)
C2	0.0217 (9)	0.0198 (9)	0.0159 (8)	-0.0038 (7)	0.0010 (7)	-0.0037 (7)
C3	0.0391 (11)	0.0255 (10)	0.0266 (10)	-0.0025 (9)	0.0095 (8)	-0.0036 (9)
C4	0.0286 (10)	0.0260 (11)	0.0477 (13)	-0.0049 (9)	0.0173 (9)	-0.0085 (10)
C5	0.0207 (8)	0.0272 (10)	0.0199 (9)	-0.0008 (8)	-0.0020 (7)	-0.0027 (8)
C6	0.0213 (8)	0.0258 (10)	0.0174 (8)	0.0004 (7)	-0.0024 (7)	0.0002 (7)
C7	0.0230 (9)	0.0247 (10)	0.0162 (8)	0.0002 (7)	-0.0009 (7)	0.0014 (7)
C8	0.0223 (9)	0.0320 (11)	0.0210 (9)	-0.0031 (8)	-0.0047 (7)	0.0021 (8)
C9	0.0290 (10)	0.0331 (11)	0.0184 (9)	0.0002 (9)	-0.0053 (7)	0.0041 (8)
C10	0.0264 (9)	0.0232 (10)	0.0190 (9)	0.0034 (7)	0.0027 (7)	0.0035 (7)
C11	0.0222 (9)	0.0376 (12)	0.0238 (10)	-0.0027 (8)	-0.0029 (8)	0.0053 (8)
C12	0.0253 (10)	0.0411 (12)	0.0212 (9)	-0.0027 (9)	-0.0052 (7)	0.0077 (9)

Geometric parameters (Å, °)

Br—C10	1.9050 (19)	C7—C12	1.387 (3)
S1—C4	1.701 (3)	C7—C8	1.401 (3)
S1—C1	1.7111 (19)	C8—C9	1.379 (3)
S2—C6	1.6629 (19)	C9—C10	1.377 (3)
O—C5	1.234 (2)	C10—C11	1.380 (3)
N1—C5	1.374 (2)	C11—C12	1.390 (3)
N1—C6	1.402 (3)	N1—H01	0.85 (3)
N2—C6	1.338 (2)	N2—H02	0.85 (3)

N2—C7	1.417 (3)	C2—H2	0.9500
C1—C2	1.402 (3)	C3—H3	0.9500
C1—C5	1.465 (3)	C4—H4	0.9500
C2—C3	1.415 (3)	C8—H8	0.9500
C3—C4	1.352 (3)	C9—H9	0.9500
C4—S1—C1	91.28 (10)	C11—C10—Br	119.39 (15)
C5—N1—C6	128.33 (16)	C10—C11—C12	119.92 (19)
C6—N2—C7	131.42 (17)	C7—C12—C11	119.56 (18)
C2—C1—C5	130.75 (17)	C5—N1—H01	117.8 (18)
C2—C1—S1	111.98 (14)	C6—N1—H01	112.6 (18)
C5—C1—S1	117.26 (14)	C6—N2—H02	113.9 (18)
C1—C2—C3	110.59 (17)	C7—N2—H02	114.7 (18)
C4—C3—C2	113.1 (2)	C1—C2—H2	124.7
C3—C4—S1	113.04 (16)	C3—C2—H2	124.7
O—C5—N1	123.18 (18)	C4—C3—H3	123.5
O—C5—C1	121.29 (17)	C2—C3—H3	123.5
N1—C5—C1	115.51 (16)	C3—C4—H4	123.5
N2—C6—N1	114.16 (16)	S1—C4—H4	123.5
N2—C6—S2	128.06 (16)	C9—C8—H8	119.8
N1—C6—S2	117.77 (13)	C7—C8—H8	119.8
C12—C7—C8	119.64 (18)	C10—C9—H9	120.3
C12—C7—N2	125.84 (17)	C8—C9—H9	120.3
C8—C7—N2	114.50 (17)	C10—C11—H11	120.0
C9—C8—C7	120.39 (18)	C12—C11—H11	120.0
C10—C9—C8	119.34 (18)	C7—C12—H12	120.2
C9—C10—C11	121.11 (18)	C11—C12—H12	120.2
C9—C10—Br	119.48 (15)		
C4—S1—C1—C2	0.46 (16)	C5—N1—C6—N2	-5.5 (3)
C4—S1—C1—C5	-178.78 (16)	C5—N1—C6—S2	173.38 (16)
C5—C1—C2—C3	179.2 (2)	C6—N2—C7—C12	3.5 (4)
S1—C1—C2—C3	0.1 (2)	C6—N2—C7—C8	-178.4 (2)
C1—C2—C3—C4	-0.8 (3)	C12—C7—C8—C9	1.3 (3)
C2—C3—C4—S1	1.1 (2)	N2—C7—C8—C9	-177.00 (19)
C1—S1—C4—C3	-0.92 (18)	C7—C8—C9—C10	0.1 (3)
C6—N1—C5—O	2.2 (3)	C8—C9—C10—C11	-1.4 (3)
C6—N1—C5—C1	-176.30 (18)	C8—C9—C10—Br	177.04 (16)
C2—C1—C5—O	164.5 (2)	C9—C10—C11—C12	1.4 (3)
S1—C1—C5—O	-16.4 (3)	Br—C10—C11—C12	-177.09 (17)
C2—C1—C5—N1	-17.0 (3)	C8—C7—C12—C11	-1.3 (3)
S1—C1—C5—N1	162.10 (14)	N2—C7—C12—C11	176.8 (2)
C7—N2—C6—N1	-177.11 (19)	C10—C11—C12—C7	0.0 (3)
C7—N2—C6—S2	4.2 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H01 \cdots S2 ⁱ	0.85 (3)	2.74 (3)	3.5625 (16)	163 (2)
N2—H02 \cdots O	0.85 (3)	1.89 (3)	2.624 (2)	144 (2)
C9—H9 \cdots S1 ⁱⁱ	0.95	2.89	3.704 (2)	144
C2—H2 \cdots S2 ⁱ	0.95	2.76	3.3193 (18)	119

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+1/2, y+3/2, -z+1/2$.