

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4-(2-Chloroethyl)morpholinium picrate

 Rajni Kant,<sup>a\*</sup> Sabeta Kohli,<sup>a</sup> Lovely Sarmal,<sup>a</sup> B. Narayana<sup>b</sup>  
and S. Samshuddin<sup>b</sup>
<sup>a</sup>Department of Physics, University of Jammu, Jammu Tawi 180 006, India, and

<sup>b</sup>Department of Studies in Chemistry, Mangalore University, Magalagangotri 574 199, India


Correspondence e-mail: rk\_paper@rediffmail.com

Received 22 July 2009; accepted 2 September 2009

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.058;  $wR$  factor = 0.179; data-to-parameter ratio = 17.3.

 The title compound,  $\text{C}_6\text{H}_{13}\text{ClNO}^+ \cdot \text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$ , was synthesized from picric acid and 4-(2-chloroethyl)morpholine. The crystal structure is stabilized by  $\text{C}-\text{H} \cdots \text{O}$  and  $\text{N}-\text{H} \cdots \text{O}$  hydrogen-bond interactions.

## Related literature

 For the homeopathic uses of the metal derivatives of picric acid, see: Maurya *et al.* (1999). For the medical applications of ammonium picrate, see: Boericke (1982) and of morpholine derivatives, see: Lutz *et al.* (1947); Hazard *et al.* (1948); Raymond *et al.* (1999). For a related structure, see: Briggs *et al.* (2004). For a description of the Cambridge Structural Database, see: Allen (2002).


## Experimental

## Crystal data

 $\text{C}_6\text{H}_{13}\text{ClNO}^+ \cdot \text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$   
 $M_r = 378.72$   
 Triclinic,  $P\bar{1}$   
 $a = 8.2063$  (4) Å  
 $b = 9.3075$  (5) Å  
 $c = 10.2896$  (6) Å  
 $\alpha = 93.954$  (5)°  
 $\beta = 95.284$  (5)°

 $\gamma = 90.376$  (4)°  
 $V = 780.65$  (7) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.30$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.30 \times 0.24 \times 0.18$  mm

## Data collection

 Oxford Diffraction Xcalibur  
 diffractometer  
 Absorption correction: none  
 8478 measured reflections

 4960 independent reflections  
 3665 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.016$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.058$   
 $wR(F^2) = 0.179$   
 $S = 1.12$   
 4960 reflections  
 286 parameters

 H atoms treated by a mixture of  
 independent and constrained  
 refinement  
 $\Delta\rho_{\text{max}} = 0.85$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.61$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N4}-\text{H4}' \cdots \text{O1}$	0.91 (3)	1.89 (3)	2.718 (2)	151.2 (19)
$\text{N4}-\text{H4}' \cdots \text{O2}$	0.91 (3)	2.25 (2)	2.896 (2)	127.6 (17)
$\text{C2}-\text{H2} \cdots \text{O3}$	0.83 (3)	2.34 (3)	2.637 (3)	102 (2)
$\text{C2}-\text{H2} \cdots \text{O4}$	0.83 (2)	2.45 (2)	2.723 (3)	101 (2)
$\text{C4}-\text{H4} \cdots \text{O6}$	0.90 (3)	2.34 (3)	2.667 (3)	101.7 (15)
$\text{C7}-\text{H7A} \cdots \text{O7}^{\text{i}}$	0.93 (3)	2.54 (3)	3.026 (3)	113 (2)
$\text{C8}-\text{H8B} \cdots \text{O5}^{\text{ii}}$	0.94 (3)	2.47 (2)	3.290 (2)	146 (2)
$\text{C9}-\text{H9A} \cdots \text{O5}^{\text{ii}}$	0.97 (3)	2.45 (3)	3.341 (2)	154 (2)
$\text{C9}-\text{H9B} \cdots \text{O2}$	0.97 (3)	2.34 (3)	2.985 (3)	123 (2)
$\text{C12}-\text{H12A} \cdots \text{O1}$	0.96 (3)	2.57 (2)	3.254 (3)	129 (2)

 Symmetry codes: (i)  $-x, -y, -z + 1$ ; (ii)  $x - 1, y - 1, z$ .

 Data collection: *CrysAlis Pro* (Oxford Diffraction, 2007); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2007); program(s) used to solve structure: *SHELXS86* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are grateful to the Department of Science and Technology of the Government of India for funding under SR/S2/CMP-47/2003 research project.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2096).

## References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Boericke, W. (1982). *Homeopathic Materia Medica*, 9th ed., pp. 46, 148, 228, 369, 516 and 687. Calcutta: Roy Publishing House.
- Briggs, C. R. S., Allen, M. J., O'Hagan, D., Tozer, D. J., Slawin, A. M. Z., Goeta, A. E. & Howard, J. A. K. (2004). *Org. Biomol. Chem.* **2**, 732–740.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hazard, R., Corteggiani, E. & Renard, S. H. (1948). *Compt. Rend.* **95**, 227.
- Lutz, R. E., Rufus, K. A., Gilbert, A., Philip, S. B., Marion, T. C., John, F. C., Adolf, J. D., James, A. F., Robert, H. J., Norman, H. L., Tellis, A. M., Kent, C. N., Russel, J. R. Jr, Newton, H. S. Jr, Smith, J. D. & James, W. W. (1947). *J. Org. Chem.* **12**, 617–703.
- Maurya, R. C. & Sharma, P. (1999). *Indian J. Chem. Sect. A*, **38**, 509–513.
- Oxford Diffraction (2007). *CrysAlis Pro* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Raymond, G. M. & Dimitrios, A. D. (1999). *Biocontrol Sci. Technol.* **9**, 53–65.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2435 [ doi:10.1107/S1600536809035405 ]

## 4-(2-Chloroethyl)morpholinium picrate

R. Kant, S. Kohli, L. Sarmal, B. Narayana and S. Samshuddin

### Comment

The metal derivatives of picric acid are also known as medicines in homeopathy (Maurya *et al.*, 1999). Silver picrate is a good antimicrobial agent. Zinc picrate is used in Bright's disease, headache, exhaustion, facial paralysis, nymphomania, paralysis, seminal emissions, spiral weakness, loss of memory and energy.

Ammonium picrate is a remedy for malarial fever and neuralgia, whooping cough, pain in the occiput and mastoid regions (Boericke, 1982).

N-substituted morpholines were used as anti-malarials (Lutz *et al.*, 1947) and 4-methyl-4-(2-phenylethyl)-morpholinium iodide have shown marked hypertension in dogs at a dose of 5 mg./kg (Hazard *et al.*, 1948). Morpholine derivatives are also used as agricultural fungicides in cereals and are known as ergosterol biosynthesis inhibitors (Raymond *et al.*, 1999).

A search of the Cambridge Structural Database (Version 5.26; Allen, 2002) reveals that there are 90 known structures that contain the morpholinium cation. Of these there are 24 that have an *N*-ethyl chain, or longer, including the structure of 4-(2-fluoroethyl) morpholinium chloride (Briggs *et al.*, 2004).

The title compound, [C<sub>6</sub> H<sub>13</sub>Cl N O]<sup>+</sup> [C<sub>6</sub> H<sub>2</sub>N<sub>3</sub> O<sub>7</sub>]<sup>-</sup>, was synthesized from picric acid and 4-(2-chloroethyl)morpholine. All geometrical parameters are in their usual ranges. The crystal structure is stabilized by C—H···O and N—H···O hydrogen interactions.

### Experimental

Picric acid (2.29 g, 0.01 mol) was dissolved in water. 4-(2-chloroethyl) morpholine (1.49 g 0.01 mol) was dissolved in 25 ml of ethanol. The two solutions were mixed and 5 ml of 5 M HCl was added to this mixture and stirred for few minutes, filtered, dried and yellow crystals of 4-(2-Chloroethyl)morpholinium picrate were obtained by slow evaporation in ethanol. (m.p. 394 K). Analytical data: Found (Cald): C %: 36.89(37.96); H%: 4.22 (4.25); N%:14.67(14.75).

### Refinement

All hydrogen atoms were located from the difference Fourier map and refined isotropically with distance restraints 0.830–0.982 Å.

## Figures




Fig. 1. View of (I) (50% probability displacement ellipsoids).





Fig. 2. Depiction of C—H...O and N—H...O interactions in the title compound(I)

## 4-(2-Chloroethyl)morpholinium picrate

### Crystal data



$M_r = 378.72$

Triclinic,  $P\bar{1}$

$a = 8.2063$  (4) Å

$b = 9.3075$  (5) Å

$c = 10.2896$  (6) Å

$\alpha = 93.954$  (5)°

$\beta = 95.284$  (5)°

$\gamma = 90.376$  (4)°

$V = 780.65$  (7) Å<sup>3</sup>

$Z = 2$

$F_{000} = 392$

$D_x = 1.611$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3665 reflections

$\theta = 3.3$ – $32.3$ °

$\mu = 0.30$  mm<sup>-1</sup>

$T = 293$  K

Rectangular, yellow

$0.30 \times 0.24 \times 0.18$  mm

### Data collection

Oxford Diffraction Xcalibur diffractometer

$T = 293$  K

$\omega$ - $2\theta$  scans

Absorption correction: none

8478 measured reflections

4960 independent reflections

3665 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\text{max}} = 32.3$ °

$\theta_{\text{min}} = 3.3$ °

$h = -12 \rightarrow 12$

$k = -8 \rightarrow 13$

$l = -14 \rightarrow 15$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0877P)^2 + 0.2886P]$$

$$R[F^2 > 2\sigma(F^2)] = 0.058$$

$$wR(F^2) = 0.179$$

$$S = 1.12$$

4960 reflections

286 parameters

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.60 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
H8B	-0.178 (3)	-0.150 (3)	0.786 (2)	0.035 (6)*
H9B	-0.286 (3)	0.146 (3)	0.883 (3)	0.041 (6)*
H4	0.386 (3)	0.587 (2)	0.610 (2)	0.033 (6)*
H9A	-0.401 (3)	0.024 (3)	0.793 (2)	0.036 (6)*
H10A	-0.331 (3)	0.311 (3)	0.731 (2)	0.039 (6)*
H2	0.299 (3)	0.541 (3)	0.972 (3)	0.042 (7)*
H11A	-0.135 (3)	-0.023 (3)	0.548 (2)	0.040 (6)*
H12A	-0.241 (3)	0.215 (3)	0.532 (3)	0.049 (7)*
H8A	-0.102 (3)	-0.044 (2)	0.887 (3)	0.033 (6)*
H11B	-0.300 (3)	-0.076 (3)	0.597 (2)	0.037 (6)*
H7B	0.087 (4)	-0.208 (3)	0.791 (3)	0.058 (8)*
H12B	-0.355 (4)	0.101 (3)	0.435 (3)	0.055 (8)*
H7A	0.069 (3)	-0.121 (3)	0.678 (3)	0.052 (7)*
H10B	-0.504 (4)	0.259 (3)	0.757 (3)	0.052 (7)*
H4'	-0.106 (3)	0.125 (3)	0.743 (2)	0.037 (6)*
Cl	0.22010 (6)	0.00390 (7)	0.83918 (7)	0.0572 (2)
N4	-0.18045 (17)	0.05189 (15)	0.73274 (15)	0.0277 (3)
C8	-0.1068 (2)	-0.0703 (2)	0.8042 (2)	0.0333 (4)
C9	-0.3282 (2)	0.1067 (2)	0.7962 (2)	0.0343 (4)
O8	-0.4535 (2)	0.1794 (2)	0.58892 (18)	0.0535 (4)
C11	-0.2318 (2)	0.0085 (2)	0.5915 (2)	0.0365 (4)
C10	-0.4054 (3)	0.2269 (2)	0.7217 (3)	0.0469 (5)
C7	0.0592 (2)	-0.1150 (2)	0.7678 (2)	0.0405 (4)
C12	-0.3138 (3)	0.1338 (3)	0.5261 (2)	0.0465 (5)
C5	0.2198 (2)	0.43814 (18)	0.62764 (19)	0.0308 (3)
N2	0.47503 (19)	0.70058 (17)	0.84556 (19)	0.0374 (4)
C2	0.2800 (2)	0.51646 (19)	0.8933 (2)	0.0325 (4)
O1	0.0317 (2)	0.25764 (19)	0.67334 (16)	0.0543 (5)
C4	0.3325 (2)	0.54510 (18)	0.6696 (2)	0.0318 (4)
C3	0.3613 (2)	0.58362 (18)	0.80201 (19)	0.0310 (4)
C1	0.1667 (2)	0.40844 (18)	0.85050 (19)	0.0311 (4)

## supplementary materials

---

C6	0.1275 (2)	0.35914 (19)	0.71437 (19)	0.0330 (4)
N3	0.0845 (2)	0.34768 (17)	0.95407 (18)	0.0384 (4)
N1	0.1924 (2)	0.40692 (18)	0.48624 (17)	0.0389 (4)
O5	0.5335 (2)	0.76658 (17)	0.76105 (19)	0.0541 (4)
O2	-0.0347 (2)	0.2682 (2)	0.92512 (19)	0.0603 (5)
O4	0.5057 (2)	0.73018 (19)	0.96342 (19)	0.0542 (4)
O3	0.1342 (2)	0.3801 (2)	1.06775 (18)	0.0568 (5)
O7	0.0693 (3)	0.3444 (3)	0.4397 (2)	0.0808 (7)
O6	0.2896 (3)	0.4530 (3)	0.4174 (2)	0.0902 (8)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl	0.0304 (2)	0.0661 (4)	0.0743 (5)	-0.0029 (2)	0.0023 (2)	0.0011 (3)
N4	0.0240 (6)	0.0286 (6)	0.0314 (7)	-0.0047 (5)	0.0033 (5)	0.0069 (5)
C8	0.0287 (8)	0.0346 (9)	0.0378 (10)	-0.0009 (7)	0.0032 (7)	0.0116 (7)
C9	0.0269 (8)	0.0356 (9)	0.0411 (10)	-0.0010 (6)	0.0065 (7)	0.0041 (7)
O8	0.0401 (8)	0.0624 (10)	0.0567 (11)	0.0076 (7)	-0.0100 (7)	0.0140 (8)
C11	0.0333 (9)	0.0426 (10)	0.0330 (9)	-0.0036 (7)	0.0009 (7)	0.0027 (7)
C10	0.0431 (11)	0.0402 (10)	0.0573 (14)	0.0088 (9)	0.0011 (10)	0.0067 (9)
C7	0.0340 (9)	0.0425 (10)	0.0449 (12)	0.0056 (8)	0.0033 (8)	0.0012 (9)
C12	0.0446 (11)	0.0585 (13)	0.0362 (11)	-0.0044 (10)	-0.0058 (9)	0.0152 (9)
C5	0.0287 (7)	0.0271 (7)	0.0367 (9)	-0.0030 (6)	0.0012 (6)	0.0057 (6)
N2	0.0289 (7)	0.0288 (7)	0.0537 (11)	-0.0048 (6)	0.0011 (7)	0.0009 (7)
C2	0.0306 (8)	0.0285 (8)	0.0382 (10)	-0.0009 (6)	0.0009 (7)	0.0042 (7)
O1	0.0577 (10)	0.0596 (10)	0.0445 (9)	-0.0371 (8)	-0.0041 (7)	0.0103 (7)
C4	0.0265 (7)	0.0258 (7)	0.0442 (10)	-0.0024 (6)	0.0056 (7)	0.0089 (7)
C3	0.0261 (7)	0.0242 (7)	0.0425 (10)	-0.0044 (6)	0.0011 (7)	0.0036 (7)
C1	0.0293 (8)	0.0284 (7)	0.0366 (9)	-0.0038 (6)	0.0041 (7)	0.0083 (7)
C6	0.0282 (8)	0.0305 (8)	0.0409 (10)	-0.0069 (6)	0.0002 (7)	0.0103 (7)
N3	0.0397 (8)	0.0313 (7)	0.0467 (10)	-0.0014 (6)	0.0122 (7)	0.0101 (7)
N1	0.0435 (9)	0.0361 (8)	0.0369 (9)	-0.0018 (7)	0.0037 (7)	0.0020 (7)
O5	0.0504 (9)	0.0417 (8)	0.0711 (12)	-0.0223 (7)	0.0093 (8)	0.0074 (8)
O2	0.0626 (11)	0.0575 (10)	0.0627 (11)	-0.0316 (8)	0.0238 (9)	-0.0016 (8)
O4	0.0494 (9)	0.0505 (9)	0.0590 (11)	-0.0126 (7)	-0.0051 (8)	-0.0080 (8)
O3	0.0639 (11)	0.0647 (11)	0.0439 (9)	-0.0126 (9)	0.0067 (8)	0.0163 (8)
O7	0.0923 (16)	0.1001 (17)	0.0461 (11)	-0.0532 (14)	-0.0013 (10)	-0.0075 (10)
O6	0.0833 (15)	0.142 (2)	0.0462 (11)	-0.0453 (16)	0.0161 (10)	0.0066 (13)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cl—C7	1.788 (2)	C12—H12A	0.96 (3)
N4—C8	1.497 (2)	C12—H12B	0.99 (3)
N4—C11	1.503 (2)	C5—C4	1.374 (2)
N4—C9	1.504 (2)	C5—C6	1.455 (2)
N4—H4'	0.91 (2)	C5—N1	1.460 (3)
C8—C7	1.500 (3)	N2—O4	1.228 (3)
C8—H8B	0.94 (2)	N2—O5	1.229 (2)
C8—H8A	0.87 (3)	N2—C3	1.449 (2)

C9—C10	1.510 (3)	C2—C3	1.382 (3)
C9—H9B	0.98 (3)	C2—C1	1.386 (2)
C9—H9A	0.97 (2)	C2—H2	0.83 (3)
O8—C12	1.420 (3)	O1—C6	1.250 (2)
O8—C10	1.426 (3)	C4—C3	1.383 (3)
C11—C12	1.517 (3)	C4—H4	0.89 (2)
C11—H11A	0.98 (3)	C1—C6	1.450 (3)
C11—H11B	0.97 (2)	C1—N3	1.456 (2)
C10—H10A	0.98 (3)	N3—O3	1.220 (2)
C10—H10B	0.96 (3)	N3—O2	1.225 (2)
C7—H7B	0.93 (3)	N1—O7	1.206 (3)
C7—H7A	0.94 (3)	N1—O6	1.210 (3)
C8—N4—C11	112.20 (15)	Cl—C7—H7A	105.7 (17)
C8—N4—C9	110.07 (14)	H7B—C7—H7A	103 (3)
C11—N4—C9	108.38 (14)	O8—C12—C11	111.16 (18)
C8—N4—H4'	106.9 (15)	O8—C12—H12A	106.5 (17)
C11—N4—H4'	113.0 (15)	C11—C12—H12A	110.6 (16)
C9—N4—H4'	106.1 (15)	O8—C12—H12B	106.1 (17)
N4—C8—C7	115.12 (16)	C11—C12—H12B	108.8 (16)
N4—C8—H8B	107.4 (14)	H12A—C12—H12B	113 (2)
C7—C8—H8B	108.0 (14)	C4—C5—C6	124.17 (18)
N4—C8—H8A	106.8 (16)	C4—C5—N1	116.02 (16)
C7—C8—H8A	110.2 (15)	C6—C5—N1	119.80 (15)
H8B—C8—H8A	109 (2)	O4—N2—O5	123.60 (17)
N4—C9—C10	110.06 (17)	O4—N2—C3	118.95 (17)
N4—C9—H9B	105.2 (15)	O5—N2—C3	117.44 (18)
C10—C9—H9B	108.9 (15)	C3—C2—C1	118.96 (18)
N4—C9—H9A	105.1 (14)	C3—C2—H2	120.0 (18)
C10—C9—H9A	111.6 (14)	C1—C2—H2	121.0 (18)
H9B—C9—H9A	116 (2)	C5—C4—C3	119.30 (16)
C12—O8—C10	109.70 (17)	C5—C4—H4	118.7 (15)
N4—C11—C12	109.86 (17)	C3—C4—H4	122.0 (15)
N4—C11—H11A	109.0 (14)	C2—C3—C4	121.53 (16)
C12—C11—H11A	111.4 (14)	C2—C3—N2	119.26 (17)
N4—C11—H11B	102.7 (14)	C4—C3—N2	119.18 (16)
C12—C11—H11B	116.2 (14)	C2—C1—C6	124.15 (16)
H11A—C11—H11B	107 (2)	C2—C1—N3	114.57 (17)
O8—C10—C9	111.16 (18)	C6—C1—N3	121.27 (15)
O8—C10—H10A	113.2 (15)	O1—C6—C1	125.59 (17)
C9—C10—H10A	109.7 (14)	O1—C6—C5	122.46 (18)
O8—C10—H10B	105.5 (17)	C1—C6—C5	111.90 (15)
C9—C10—H10B	111.4 (17)	O3—N3—O2	121.69 (18)
H10A—C10—H10B	106 (2)	O3—N3—C1	118.95 (17)
C8—C7—C1	113.24 (15)	O2—N3—C1	119.35 (18)
C8—C7—H7B	113.2 (19)	O7—N1—O6	121.2 (2)
Cl—C7—H7B	107.2 (18)	O7—N1—C5	119.83 (18)
C8—C7—H7A	114.0 (18)	O6—N1—C5	118.78 (18)

## supplementary materials

---

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N4—H4' $\cdots$ O1	0.91 (3)	1.89 (3)	2.718 (2)	151.2 (19)
N4—H4' $\cdots$ O2	0.91 (3)	2.25 (2)	2.896 (2)	127.6 (17)
C2—H2 $\cdots$ O3	0.83 (3)	2.34 (3)	2.637 (3)	102 (2)
C2—H2 $\cdots$ O4	0.83 (2)	2.45 (2)	2.723 (3)	101 (2)
C4—H4 $\cdots$ O6	0.90 (3)	2.34 (3)	2.667 (3)	101.7 (15)
C7—H7A $\cdots$ O7 <sup>i</sup>	0.93 (3)	2.54 (3)	3.026 (3)	113 (2)
C8—H8B $\cdots$ O5 <sup>ii</sup>	0.94 (3)	2.47 (2)	3.290 (2)	146 (2)
C9—H9A $\cdots$ O5 <sup>ii</sup>	0.97 (3)	2.45 (3)	3.341 (2)	154 (2)
C9—H9B $\cdots$ O2	0.97 (3)	2.34 (3)	2.985 (3)	123 (2)
C12—H12A $\cdots$ O1	0.96 (3)	2.57 (2)	3.254 (3)	129 (2)

Symmetry codes: (i)  $-x, -y, -z+1$ ; (ii)  $x-1, y-1, z$ .

Fig. 1




Fig. 2

