

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# {4-Bromo-2-[2-(piperidin-1-ium-1-yl)-ethyliminomethyl]phenolato}diiodido-zinc(II)

Xue-Wen Zhu,\* Zhi-Gang Yin, Gang-Sen Li, Xu-Zhao Yang and Chun-Xia Zhang

Key Laboratory of Surface and Interface Science of Henan, School of Materials and Chemical Engineering, Zhengzhou University of Light Industry, Zhengzhou 450002, People's Republic of China

Correspondence e-mail: xuewen-zhu@163.com


Received 10 September 2009; accepted 14 September 2009

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.009$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.093; data-to-parameter ratio = 21.7.

In the title complex,  $[\text{ZnI}_2(\text{C}_{14}\text{H}_{19}\text{BrN}_2\text{O})]$ , the  $\text{Zn}^{\text{II}}$  atom is four-coordinated by the imine N and phenolate O atoms of the Schiff base ligand and by two iodide ions in a distorted tetrahedral coordination. In the crystal structure, molecules are linked through intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, forming chains running along the  $b$  axis.

## Related literature

For background to the chemistry of Schiff base complexes, see: Ali *et al.* (2008); Biswas *et al.* (2008); Chen *et al.* (2008); Darensbourg & Frantz (2007); Habibi *et al.* (2007); Kawamoto *et al.* (2008); Lipscomb & Sträter (1996); Tomat *et al.* (2007); Wu *et al.* (2008); Yuan *et al.* (2007). For related structures, see: Zhu (2008); Zhu & Yang (2008a,b,c); Qiu (2006a,b); Wei *et al.* (2007); Zhu *et al.* (2007).



## Experimental

### Crystal data

$[\text{ZnI}_2(\text{C}_{14}\text{H}_{19}\text{BrN}_2\text{O})]$   
 $M_r = 630.39$   
 Monoclinic,  $P2_1/n$   
 $a = 10.470$  (3) Å

$b = 12.351$  (3) Å  
 $c = 15.426$  (4) Å  
 $\beta = 102.326$  (3)°  
 $V = 1948.8$  (9) Å<sup>3</sup>

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 6.48$  mm<sup>-1</sup>

$T = 298$  K  
 $0.17 \times 0.15 \times 0.15$  mm

### Data collection

Bruker APEXII CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)  
 $T_{\text{min}} = 0.406$ ,  $T_{\text{max}} = 0.443$

14478 measured reflections  
 4193 independent reflections  
 2953 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.053$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.093$   
 $S = 1.03$   
 4193 reflections  
 193 parameters  
 1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.68$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -1.09$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

Zn1—O1	1.981 (4)	Zn1—I2	2.5428 (9)
Zn1—N1	2.047 (4)	Zn1—I1	2.5771 (9)
O1—Zn1—N1	91.61 (16)	O1—Zn1—I1	109.08 (12)
O1—Zn1—I2	114.27 (10)	N1—Zn1—I1	108.67 (12)
N1—Zn1—I2	113.47 (12)	I2—Zn1—I1	116.86 (3)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O1}^i$	0.90 (5)	1.85 (5)	2.745 (6)	176 (7)

Symmetry code: (i)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2276).

## References

- Ali, H. M., Mohamed Mustafa, M. I., Rizal, M. R. & Ng, S. W. (2008). *Acta Cryst.* **E64**, m718–m719.  
 Biswas, C., Drew, M. G. B. & Ghosh, A. (2008). *Inorg. Chem.* **47**, 4513–4519.  
 Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Chen, Z., Morimoto, H., Matsunaga, S. & Shibasaki, M. (2008). *J. Am. Chem. Soc.* **130**, 2170–2171.  
 Darensbourg, D. J. & Frantz, E. B. (2007). *Inorg. Chem.* **46**, 5967–5978.  
 Habibi, M. H., Askari, E., Chantrapromma, S. & Fun, H.-K. (2007). *Acta Cryst.* **E63**, m2905–m2906.  
 Kawamoto, T., Nishiwaki, M., Tsunekawa, Y., Nozaki, K. & Konno, T. (2008). *Inorg. Chem.* **47**, 3095–3104.  
 Lipscomb, W. N. & Sträter, N. (1996). *Chem. Rev.* **96**, 2375–2434.  
 Qiu, X.-Y. (2006a). *Acta Cryst.* **E62**, m717–m718.  
 Qiu, X.-Y. (2006b). *Acta Cryst.* **E62**, m2173–m2174.

- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tomat, E., Cuesta, L., Lynch, V. M. & Sessler, J. L. (2007). *Inorg. Chem.* **46**, 6224–6226.
- Wei, Y.-J., Wang, F.-W. & Zhu, Q.-Y. (2007). *Acta Cryst.* **E63**, m654–m655.
- Wu, J.-C., Liu, S.-X., Keene, T. D., Neels, A., Mereacre, V., Powell, A. K. & Decurtins, S. (2008). *Inorg. Chem.* **47**, 3452–3459.
- Yuan, M., Zhao, F., Zhang, W., Wang, Z.-M. & Gao, S. (2007). *Inorg. Chem.* **46**, 11235–11242.
- Zhu, X.-W. (2008). *Acta Cryst.* **E64**, m1456–m1457.
- Zhu, Q.-Y., Wei, Y.-J. & Wang, F.-W. (2007). *Acta Cryst.* **E63**, m1431–m1432.
- Zhu, X.-W. & Yang, X.-Z. (2008a). *Acta Cryst.* **E64**, m1090–m1091.
- Zhu, X.-W. & Yang, X.-Z. (2008b). *Acta Cryst.* **E64**, m1092–m1093.
- Zhu, X.-W. & Yang, X.-Z. (2008c). *Acta Cryst.* **E64**, m1094–m1095.

**supplementary materials**

*Acta Cryst.* (2009). E65, m1226-m1227 [ doi:10.1107/S1600536809037210 ]

## {4-Bromo-2-[2-(piperidin-1-ium-1yl)ethyliminomethyl]phenolato}diiodidozinc(II)

X.-W. Zhu, Z.-G. Yin, G.-S. Li, X.-Z. Yang and C.-X. Zhang

### Comment

Schiff bases have widely been used as versatile ligands in coordination chemistry (Biswas *et al.*, 2008; Wu *et al.*, 2008; Kawamoto *et al.*, 2008; Ali *et al.*, 2008; Habibi *et al.*, 2007), and their metal complexes are of great interest in many fields (Chen *et al.*, 2008; Yuan *et al.*, 2007; Tomat *et al.*, 2007; Darensbourg & Frantz, 2007). Zinc(II) is an important element in biological systems and functions as the active site of hydrolytic enzymes, such as carboxypeptidase and carbonic anhydrase where it is in a hard-donor coordination environment of nitrogen and oxygen ligands (Lipscomb & Sträter, 1996). Recently, we have reported a few Schiff base zinc complexes (Zhu, 2008; Zhu & Yang, 2008*a,b,c*). In this paper, the title new zinc(II) complex, Fig. 1, is reported.

In the title complex, the Zn<sup>II</sup> atom is four-coordinated by the imine N and phenolate O atoms of the Schiff base ligand, and by two iodide ions in a tetrahedral coordination. The coordinate bond lengths (Table 1) are typical and comparable to the corresponding values observed in the Schiff base zinc complexes we reported previously and other similar Schiff base zinc complexes (Zhu *et al.*, 2007; Wei *et al.*, 2007; Qiu, 2006*a,b*).

In the crystal structure, molecules are linked through intermolecular N—H...O hydrogen bonds, forming chains running along the *b* axis (Table 2, Fig. 2).

### Experimental

The Schiff base compound was prepared by the condensation of equimolar amounts of 5-bromosalicylaldehyde with 2-piperidin-1-ylethylamine in a methanol solution. The complex was prepared by the following method. To an anhydrous methanol solution (5 ml) of ZnI<sub>2</sub> (31.9 mg, 0.1 mmol) was added a methanol solution (10 ml) of the Schiff base compound (31.1 mg, 0.1 mmol) with stirring. The mixture was stirred for 30 min at room temperature and filtered. Upon keeping the filtrate in air for a few days, colorless block-shaped crystals were formed.

### Refinement

H2 was located from a difference Fourier map and refined isotropically, with N—H distance restrained to 0.90 (1) Å. Other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of the title complex, with ellipsoids drawn at the 30% probability level.




Fig. 2. The crystal packing of the title complex.

## [4-Bromo-2-[2-(piperidin-1-ium-1-yl)ethyliminomethyl]phenolato]diiodidozinc(II)

### Crystal data

[ZnI <sub>2</sub> (C <sub>14</sub> H <sub>19</sub> BrN <sub>2</sub> O)]	$F_{000} = 1184$
$M_r = 630.39$	$D_x = 2.149 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1 n$	Cell parameters from 2365 reflections
$a = 10.470 (3) \text{ \AA}$	$\theta = 2.5\text{--}25.1^\circ$
$b = 12.351 (3) \text{ \AA}$	$\mu = 6.48 \text{ mm}^{-1}$
$c = 15.426 (4) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 102.326 (3)^\circ$	Block, colorless
$V = 1948.8 (9) \text{ \AA}^3$	$0.17 \times 0.15 \times 0.15 \text{ mm}$
$Z = 4$	

### Data collection

Bruker APEXII CCD area-detector diffractometer	4193 independent reflections
Radiation source: fine-focus sealed tube	2953 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.053$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
$\omega$ scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	$h = -13 \rightarrow 12$
$T_{\text{min}} = 0.406$ , $T_{\text{max}} = 0.443$	$k = -15 \rightarrow 15$
14478 measured reflections	$l = -19 \rightarrow 19$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.0326P)^2 + 0.0119P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
4193 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
193 parameters	$\Delta\rho_{\text{max}} = 0.68 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -1.09 \text{ e \AA}^{-3}$

1 restraint  
 Extinction correction: none  
 Primary atom site location: structure-invariant direct methods

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.16997 (7)	0.57708 (5)	0.70705 (4)	0.03725 (18)
I1	-0.02704 (5)	0.68332 (4)	0.61843 (3)	0.05599 (16)
I2	0.16625 (5)	0.53139 (4)	0.86722 (3)	0.05989 (16)
Br1	0.46329 (7)	0.66143 (6)	0.32863 (4)	0.04966 (19)
O1	0.3336 (4)	0.6472 (3)	0.6911 (2)	0.0409 (10)
N1	0.2025 (4)	0.4456 (3)	0.6339 (3)	0.0316 (10)
N2	-0.0218 (5)	0.2189 (4)	0.6686 (3)	0.0329 (10)
C1	0.3224 (5)	0.5601 (4)	0.5493 (3)	0.0331 (13)
C2	0.3580 (6)	0.6466 (4)	0.6099 (3)	0.0354 (13)
C3	0.4260 (6)	0.7331 (5)	0.5832 (4)	0.0463 (16)
H3	0.4494	0.7909	0.6221	0.056*
C4	0.4595 (6)	0.7367 (5)	0.5032 (4)	0.0464 (15)
H4	0.5063	0.7953	0.4883	0.056*
C5	0.4238 (6)	0.6527 (5)	0.4440 (3)	0.0356 (14)
C6	0.3578 (6)	0.5660 (4)	0.4664 (3)	0.0365 (14)
H6	0.3356	0.5095	0.4261	0.044*
C7	0.2634 (5)	0.4608 (4)	0.5703 (3)	0.0321 (13)
H7	0.2697	0.4014	0.5343	0.039*
C8	0.1621 (6)	0.3353 (4)	0.6490 (4)	0.0382 (14)
H8A	0.2113	0.3096	0.7059	0.046*
H8B	0.1814	0.2880	0.6032	0.046*
C9	0.0187 (6)	0.3307 (4)	0.6481 (4)	0.0405 (14)
H9A	-0.0013	0.3811	0.6916	0.049*
H9B	-0.0305	0.3525	0.5901	0.049*
C10	-0.1500 (6)	0.2221 (5)	0.6981 (4)	0.0511 (17)
H10A	-0.2166	0.2535	0.6514	0.061*
H10B	-0.1411	0.2680	0.7501	0.061*
C11	-0.1929 (7)	0.1106 (6)	0.7199 (5)	0.0611 (19)
H11A	-0.1316	0.0830	0.7714	0.073*
H11B	-0.2780	0.1158	0.7352	0.073*

## supplementary materials

---

C12	-0.2011 (7)	0.0321 (6)	0.6442 (5)	0.068 (2)
H12A	-0.2200	-0.0400	0.6630	0.082*
H12B	-0.2711	0.0533	0.5951	0.082*
C13	-0.0735 (8)	0.0318 (5)	0.6150 (4)	0.0585 (19)
H13A	-0.0804	-0.0155	0.5641	0.070*
H13B	-0.0060	0.0030	0.6625	0.070*
C14	-0.0340 (7)	0.1429 (5)	0.5910 (4)	0.0466 (16)
H14A	-0.0989	0.1706	0.5414	0.056*
H14B	0.0490	0.1388	0.5728	0.056*
H2	0.038 (5)	0.197 (5)	0.716 (3)	0.080*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.0429 (4)	0.0396 (4)	0.0303 (3)	-0.0049 (3)	0.0101 (3)	-0.0033 (3)
I1	0.0597 (3)	0.0710 (3)	0.0357 (2)	0.0164 (2)	0.0068 (2)	0.0024 (2)
I2	0.0858 (4)	0.0614 (3)	0.0341 (2)	-0.0066 (3)	0.0165 (2)	0.0058 (2)
Br1	0.0485 (4)	0.0681 (5)	0.0363 (3)	0.0073 (3)	0.0178 (3)	0.0093 (3)
O1	0.046 (3)	0.049 (2)	0.0271 (19)	-0.013 (2)	0.0072 (17)	-0.0100 (17)
N1	0.031 (3)	0.028 (2)	0.036 (2)	-0.003 (2)	0.008 (2)	0.0024 (19)
N2	0.034 (3)	0.034 (3)	0.030 (2)	-0.001 (2)	0.008 (2)	0.004 (2)
C1	0.030 (3)	0.034 (3)	0.032 (3)	0.002 (3)	0.000 (2)	-0.004 (2)
C2	0.036 (4)	0.040 (3)	0.029 (3)	-0.002 (3)	0.004 (2)	0.002 (2)
C3	0.062 (5)	0.047 (4)	0.029 (3)	-0.011 (3)	0.009 (3)	-0.007 (3)
C4	0.049 (4)	0.042 (4)	0.050 (4)	-0.009 (3)	0.015 (3)	0.005 (3)
C5	0.037 (4)	0.047 (4)	0.025 (3)	0.010 (3)	0.012 (2)	0.012 (2)
C6	0.043 (4)	0.040 (3)	0.026 (3)	0.006 (3)	0.006 (2)	-0.003 (2)
C7	0.037 (4)	0.028 (3)	0.030 (3)	0.001 (2)	0.004 (2)	-0.008 (2)
C8	0.038 (4)	0.037 (3)	0.039 (3)	-0.001 (3)	0.007 (3)	0.003 (3)
C9	0.040 (4)	0.036 (3)	0.045 (3)	0.004 (3)	0.009 (3)	0.007 (3)
C10	0.040 (4)	0.061 (4)	0.054 (4)	0.010 (3)	0.015 (3)	0.013 (3)
C11	0.047 (4)	0.066 (5)	0.075 (5)	-0.003 (4)	0.022 (4)	0.021 (4)
C12	0.061 (5)	0.061 (5)	0.078 (5)	-0.018 (4)	0.006 (4)	0.019 (4)
C13	0.075 (5)	0.050 (4)	0.053 (4)	-0.018 (4)	0.017 (4)	-0.010 (3)
C14	0.053 (4)	0.052 (4)	0.036 (3)	-0.014 (3)	0.014 (3)	-0.002 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Zn1—O1	1.981 (4)	C6—H6	0.9300
Zn1—N1	2.047 (4)	C7—H7	0.9300
Zn1—I2	2.5428 (9)	C8—C9	1.500 (8)
Zn1—I1	2.5771 (9)	C8—H8A	0.9700
Br1—C5	1.914 (5)	C8—H8B	0.9700
O1—C2	1.329 (6)	C9—H9A	0.9700
N1—C7	1.293 (7)	C9—H9B	0.9700
N1—C8	1.460 (6)	C10—C11	1.509 (9)
N2—C9	1.498 (7)	C10—H10A	0.9700
N2—C14	1.505 (7)	C10—H10B	0.9700
N2—C10	1.508 (7)	C11—C12	1.507 (10)

N2—H2	0.90 (5)	C11—H11A	0.9700
C1—C6	1.407 (7)	C11—H11B	0.9700
C1—C2	1.417 (7)	C12—C13	1.498 (10)
C1—C7	1.441 (7)	C12—H12A	0.9700
C2—C3	1.395 (8)	C12—H12B	0.9700
C3—C4	1.354 (8)	C13—C14	1.502 (8)
C3—H3	0.9300	C13—H13A	0.9700
C4—C5	1.380 (8)	C13—H13B	0.9700
C4—H4	0.9300	C14—H14A	0.9700
C5—C6	1.359 (8)	C14—H14B	0.9700
O1—Zn1—N1	91.61 (16)	N1—C8—H8B	109.5
O1—Zn1—I2	114.27 (10)	C9—C8—H8B	109.5
N1—Zn1—I2	113.47 (12)	H8A—C8—H8B	108.0
O1—Zn1—I1	109.08 (12)	N2—C9—C8	111.1 (4)
N1—Zn1—I1	108.67 (12)	N2—C9—H9A	109.4
I2—Zn1—I1	116.86 (3)	C8—C9—H9A	109.4
C2—O1—Zn1	117.4 (3)	N2—C9—H9B	109.4
C7—N1—C8	117.6 (4)	C8—C9—H9B	109.4
C7—N1—Zn1	118.0 (3)	H9A—C9—H9B	108.0
C8—N1—Zn1	124.4 (3)	N2—C10—C11	111.7 (5)
C9—N2—C14	112.8 (4)	N2—C10—H10A	109.3
C9—N2—C10	110.5 (4)	C11—C10—H10A	109.3
C14—N2—C10	109.1 (5)	N2—C10—H10B	109.3
C9—N2—H2	106 (5)	C11—C10—H10B	109.3
C14—N2—H2	113 (4)	H10A—C10—H10B	107.9
C10—N2—H2	106 (5)	C12—C11—C10	112.6 (6)
C6—C1—C2	118.6 (5)	C12—C11—H11A	109.1
C6—C1—C7	117.2 (5)	C10—C11—H11A	109.1
C2—C1—C7	123.9 (5)	C12—C11—H11B	109.1
O1—C2—C3	119.1 (5)	C10—C11—H11B	109.1
O1—C2—C1	123.3 (5)	H11A—C11—H11B	107.8
C3—C2—C1	117.5 (5)	C13—C12—C11	109.1 (6)
C4—C3—C2	122.8 (6)	C13—C12—H12A	109.9
C4—C3—H3	118.6	C11—C12—H12A	109.9
C2—C3—H3	118.6	C13—C12—H12B	109.9
C3—C4—C5	119.5 (6)	C11—C12—H12B	109.9
C3—C4—H4	120.2	H12A—C12—H12B	108.3
C5—C4—H4	120.2	C12—C13—C14	112.5 (6)
C6—C5—C4	120.3 (5)	C12—C13—H13A	109.1
C6—C5—Br1	120.2 (4)	C14—C13—H13A	109.1
C4—C5—Br1	119.4 (4)	C12—C13—H13B	109.1
C5—C6—C1	121.3 (5)	C14—C13—H13B	109.1
C5—C6—H6	119.4	H13A—C13—H13B	107.8
C1—C6—H6	119.4	C13—C14—N2	110.8 (5)
N1—C7—C1	126.8 (5)	C13—C14—H14A	109.5
N1—C7—H7	116.6	N2—C14—H14A	109.5
C1—C7—H7	116.6	C13—C14—H14B	109.5
N1—C8—C9	110.9 (4)	N2—C14—H14B	109.5
N1—C8—H8A	109.5	H14A—C14—H14B	108.1

## supplementary materials

---

C9—C8—H8A 109.5

### *Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N2—H2...O1 <sup>i</sup>	0.90 (5)	1.85 (5)	2.745 (6)	176 (7)

Symmetry codes: (i)  $-x+1/2, y-1/2, -z+3/2$ .

Fig. 1




Fig. 2

